



Distance from the edge of forest fragments influence the abundance of aphidophagous hoverflies (Diptera: Syrphidae) in wheat fields

Orcial Ceolin Bortolotto*, Ayres de Oliveira Menezes Júnior, Adriano Thibes Hoshino and Thiago Augusto Campos

Universidade Estadual de Londrina, Rua Celso Garcia Cid, s/n.º, 86057-970, Campus Universitário, Londrina, Paraná, Brazil. *Author for correspondence. E-mail: bortolotto.orcial@gmail.com

ABSTRACT. This study aimed to evaluate the influence of the distance from the edge of native forests on the abundance of aphidophagous hoverflies in wheat fields. The study was conducted in four commercial fields in the North of Paraná State, Brazil, during the wheat crop season of 2009. Two transects were surveyed parallel to the edge of the forest at two distances: 25 (“near”) and 525 meters (“far”) from the edge. The abundance of hoverflies was compared using a χ^2 test ($p \leq 0.05$). In total, 1,845 hoverflies adults were collected, which represented 15 species and three genera. The most abundant species was *Allograpta exotica* (60.43%), followed by *Toxomerus dispar* (17.78%) and *Toxomerus watsoni* (7.26%) (Diptera: Syrphidae). An important inference was that all fields showed a higher abundance of aphidophagous hoverflies closer to the edge of the forest (25 m) during the wheat tillering stage. The initial abundance of aphidophagous hoverflies in wheat fields is likely greater near the edge because of the availability of resources in the surrounding forest that enhance hoverfly survival during periods of low aphid infestation.

Keywords: conservative biological control, predators, wheat pests, landscape preservation, refuge area.

A distância da borda de fragmentos florestais influencia a abundância de sirfídeos afidófagos (Diptera: Syrphidae) em lavouras de trigo

RESUMO. O objetivo desta pesquisa foi avaliar a influência da distância do fragmento de mata nativa sobre a abundância de sirfídeos afidófagos em lavoura de trigo. O experimento foi conduzido em quatro lavouras comerciais, localizadas no Norte do estado do Paraná, durante a safra 2009. Para isto, foram estabelecidos dois transectos paralelos à borda da mata, nas distâncias de 25 (“perto”) e 525 metros (“longe”) da borda. A abundância de sirfídeos foi comparada empregando-se o teste χ^2 ($p \leq 0,05$). No total, foram capturados 1845 sirfídeos adultos, representados por 15 espécies e três gêneros. A maior abundância de sirfídeos afidófagos foi registrada para espécie *Allograpta exotica* (60,43%), seguida por *Toxomerus dispar* (17,78%) e *Toxomerus watsoni* (7,26%) (Diptera: Syrphidae). Uma inferência de destaque foi que durante o estágio de perfilhamento do trigo, todas as lavouras apresentaram maior abundância de sirfídeos afidófagos próximo à borda da mata (25m). A abundância inicial da população de sirfídeos afidófagos na lavoura de trigo é maior na proximidade da borda da mata, provavelmente devido os recursos disponíveis para sobreviverem durante a baixa infestação de presas na lavoura.

Palavras-chave: controle biológico por conservação, predadores, pragas de trigo, preservação de paisagem, áreas de refúgio.

Introduction

Hoverflies (Diptera: Syrphidae) are notably important insects because they act as pollinators (Jun, Yibo, Jin, Fazhi, & Zhenhai, 2009; Blaauw & Isaacs, 2014) and biological control agents (Irshad, 2014). In wheat crops (*Triticum aestivum* L.), the agronomic importance is at the larval stage, where aphid predators (Schmidt et al., 2003; Bugg, Colfer, Chaney, Smith, & Cannon, 2008) can consume up to 2,000 aphids (Dib, Simon, Sauphanor, & Capowiez, 2010; Hogg, Bugg, & Daane, 2011). Meanwhile, adult hoverflies feed on nectar and

pollen, whose energy (Van Rijn, Kooijman, & Wackers, 2013) increases the reproductive rate (Laubertie, Wratten, & Hemptinne, 2012) and longevity (Pinheiro, Torres, Raimundo, & Santos, 2015). Because of the importance of floral resources to hoverflies, many studies have been conducted to promote the action of this natural enemy in crop fields using landscape management (Hickman & Wratten, 1996; Bokina, 2012; Amaral et al., 2013; Gontijo, Beers, & Snyder, 2013; Martínez-Uña, Martín, Fernández-Quintanilla, & Dorado, 2013; Haenke et al., 2014).

Moreover, the remaining forest fragment may act as an area of refuge to natural enemies during little abundance of pests in crop fields (Koh & Holland, 2015). In Brazil, although several studies have shown that natural enemies are highly efficient in suppressing aphid infestation in wheat crops (Salvadori & Salles, 2002; Alves, Prestes, Zanini, Dalmolin, & Menezes Jr., 2005; Bortolotto, Menezes Jr., Sampaio, & Hoshino, 2012; Bortolotto, Menezes Jr., & Hoshino, 2015), have no investigation about the influence of non-crop fields on these natural enemies. Among the conservative management strategies, the maintenance of non-crop habitats has been studied to understand the importance of natural enemies (Barbosa, 1998; Walton & Isaacs, 2011; Letourneau, Bothwell, & Stireman, 2012; Fahrig et al., 2015). However, it is important to consider that most studies were developed in temperate regions. Hence, the importance of the remaining forest fragment near crop fields and its contribution to natural enemies are unavailable for the Neotropical agroecosystem. In addition, some investigations were reported a devastation of native forest fragments in Brazil reducing non-crop areas (Fearnside, 2001; Nassar, 2009), and this practice caused impact on biodiversity (Vieira, Toledo, Silva, & Higuchi, 2008). In this sense, it is important to develop research to increase understanding about the importance of remaining forest fragments for natural biological control, and to encourage growers to maintain refuge areas in their farms. Thus, this study investigates the effect of the distance from the edge of a forest fragment on aphidophagous hoverflies in wheat fields.

Material and methods

Study site

The study was performed in four commercial wheat crops in northern Paraná State, Brazil, during the 2009 crop season (usually in approximately April/May to August/September). The sites located in the municipalities of Ibiporã-Santo Antônio Farm (SAF) (23° 14' 34" S 51° 27' 07" W), Ibiporã-Bonsucesso Farm (BF) (23° 12' 26" S 51° 03' 51" W), Rolândia (Gioconda Farm) (23° 23' 59" S 5° 19' 01" W), and Londrina (23° 19' 49" S 51° 08' 12" W) (Table 1).

In all of the studied fields, wheat was sown in succession to soybean [*Glycine max* (Merrill) L.] in no-tillering soil. The fields were sown on April 29 (Rolândia-GF), May 6 (Ibiporã-FBS), May 8 (Ibiporã-SAF), and May 11 (Londrina). In general, the landscape complex around each farm (with an approximately 2 km radius) was primarily composed of wheat crops (32 to 47%), followed by non-crop area [Atlantic forest fragment (compound majority by *Aspidosperma polyneuronand*; *Ficus* spp.; *Euterpe edulis*; Orchidaceae and grasses) and pasture] (26.2 to 39%), maize (7 to 26.5%) and coffee crops (2.9 to 20.5%). Exceptionally in Ibiporã-SAF oleraceous, fruits and fallow areas (predominantly infested with wild radish) were reported.

Hoverfly survey

Hoverflies were assessed in two transects (90 m in length) that demarcated each field (adapted from Murta, Ker, Costa, Espírito-Santo, & Faria, 2008). The transects were set up immediately after the emergence (phenological stage V1) of wheat crops in all fields except Londrina, which was demarcated one day after sowing. In this sense, the assessment of hoverflies began on the same day that the transect was demarcated. To compare the "edge effect", one transect was demarcated at 25 m from the edge (near), and the other was at 525 m from the forest edge (far).

The hoverfly abundance was monitored using Malaise traps (Petanidou, Vuji, & Ellis, 2011), which were placed at the center of each transect ($n = \text{two traps field}^{-1}$). The traps were made of a synthetic material and shaped like a tent, with an opening at the bottom that intercepted the insects during flight after they collided with one of the trap's septa. The traps were installed and positioned to face north, where the most sunlight was received. Each trap was approximately 1.80 m high $\times$ 1.80 m long. The collecting bottle contained 70% alcohol, which was changed weekly on the day that the number of aphids was counted. In the laboratory, the collected material was screened, and the hoverflies were identified using a stereoscope microscope according to an identification guide (Marinoni, Morales, & Spaler, 2007; Borges & Couri, 2009; Mengual, Ruiz, Rojo, Stahl, & Thompson, 2009).

Table 1. Farm description and climatic data during the study. Paraná State, crop season (May to September of 2009).

Description	Farms			
	Ibiporã (SAF)	Ibiporã (BF)	Rolândia (GF)	Londrina
Wheat field (size)	2.3 ha	16.2 ha	20.9 ha	11.4 ha
Forest fragment (size)	45 ha	24.5 ha	380 ha	25 ha
Rapid Ecological Assessment ¹	34	28	48	50
Ecological Integrity ¹	Medium	Poor	Good	Excellent
Climatic data	-	-	-	-
Temperature average (°C)	20.54	20.54	17	17.5
Rainfall total (mm)	461.00	461.00	351.7	585.9

¹Methodology and classification according to Medeiros and Torezan (2013).

Aphids survey

Aphid assessment was made for both distances (25 and 525 m from the edge of the forest). Each transect contained 10 evaluation points, where 20 tillers at each point were randomly evaluated ($n = 400$ tillers assessment in each field). All aphids were quantified and identified to the species level. To reduce interference, insecticides were not applied to a distance of 5 m from the evaluation points.

Similar to the hoverfly assessment, the aphid infestation survey began on the same day that the transect was demarcated, except in Londrina (seven days after the transect was demarcated). The assessments were performed on a weekly basis, and the aphid species were identified using a guide developed by Salvadori and Tonet (2001).

Statistical analysis

The aphid means were subjected to exploratory analyses to assess the assumptions of the normality of residuals, homogeneity of variance of the treatments, and additivity of the model to allow for parametric tests. The aphid abundance near (25 m) and far (525 m) the edge of the forest was compared using Student's *t*-test.

To compare the hoverfly abundance, the specimens were summed at each growth stage (except Rolândia-GF because of low abundance) and compared using a χ^2 (Chi-square) test. In addition, the relationship between aphid and hoverfly

abundance (density-dependence) was estimated using a quadratic regression model. The difference was considered significant only when the significance level was $p \leq 0.05$.

Results and discussion

In total, 1,845 adult hoverflies were captured, which represented 15 species and three genera (Table 2). The most abundant aphidophagous hoverfly species were *Allograpta exotica* (60.43%), *Toxomerus dispar* (17.78%) and *Toxomerus watsoni* (26.7%) (Diptera: Syrphidae). The reported major abundance of *Allograpta* and *Toxomerus* genus in the present study is consistent with that obtained in a previous report in the Neotropical region (Thompson, 1999), which indicates the adaptation of these hoverflies.

Interestingly, during the wheat tillering stage, consistently higher hoverfly abundance was reported near (25 m) the forest edge than far (525 m) from the edge of the forest (Figure 1). In this period, the predominant species was *A. exotica* (58%), which was followed by *Pseudodorus clavatus* (12%) (Diptera: Syrphidae). In Rolândia-GF farm, a low hoverfly abundance was captured (n total = 74), which did not allow us to perform a statistical test at each growth stage. However, the total abundance was higher near the forest edge (25 m) than far (525 m) from the edge of the forest fragment (Figure 1).

Table 2. Hoverflies (Diptera: Syrphidae) captured in Malaise trap during the crop season of 2009 (May to September) in the Northeast of Paraná State.

Syrphidae species	Farms				Total	
	Ibiporã (SAF)	Ibiporã (BF)	Rolândia (GF)	Londrina	Absolute	Relative
<i>Allograpta annulipes</i>	3	0	0	0	3	0.16%
<i>Allograpta exotica</i>	794	169	40	112	1,115	60.43%
<i>Allograpta falcata</i>	3	0	0	0	3	0.16%
<i>Allograpta hastata</i>	3	10	0	0	13	0.70%
<i>Allograpta neotropica</i>	3	0	0	0	3	0.16%
<i>Allograpta obliqua</i>	0	0	3	3	6	0.33%
<i>Pseudodorus clavatus</i>	6	0	13	18	37	2.01%
<i>Ocyrtamys dimidiatus</i>	3	0	0	9	12	0.65%
<i>Ocyrtamys gastrostactus</i>	0	9	0	0	9	0.49%
<i>Ocyrtamys</i> sp.1	3	0	0	0	3	0.16%
<i>Ocyrtamys</i> sp.2	0	0	0	3	3	0.16%
<i>Ocyrtamys</i> sp.3	0	0	0	3	3	0.16%
<i>Syrphus phaeostigma</i>	3	0	6	3	12	0.65%
<i>Toxomerus dispar</i>	258	15	0	55	328	17.78%
<i>Toxomerus floralis</i>	0	0	3	12	15	0.81%
<i>Toxomerus lacrymosus</i>	3	0	0	0	3	0.16%
<i>Toxomerus politus</i>	0	3	9	6	18	0.98%
<i>Toxomerus</i> sp.	52	22	0	48	122	6.61%
<i>Toxomerus watsoni</i>	134	0	0	0	134	7.26%
<i>Trichopsomyia</i> sp.	3	0	0	0	3	0.16%
Absolute abundance	1,271	228	74	272	1,845	100%
Relative abundance	68.89%	12.36%	4.01%	14.74%	-	100%
Richness	14	6	6	11	-	-




Figure 1. Hoverflies (Diptera: Syrphidae) captured in Malaise traps near (25 m) and far (525 m) from the edge of the forest fragments in different wheat growth stages (T = tillering; B = boot; F = flowering; G = grain). Parana State, wheat season (May to September) 2009. (* $p < 0.05$; ** $p < 0.01$; χ^2 test).

The highest abundance of *A. exotica* indicates a nice establishment of this specie in wheat crop. In a study developed by Greco (1995) the authors reported *A. exotica* strongly associated with the aphid abundance in wheat fields, reinforcing the importance of this specie for aphid suppression. In the current study, the aphids were represented by only two species: *Siobion avenae*, which exhibited greater abundance (85%) (Hemiptera: Aphididae), and *Rhopalosiphum padi* (15%) (Hemiptera: Aphididae). The highest aphid abundance was reported during the wheat grain stage (Figure 2), which can be explained by the largest occurrence of *S. avenae*.




Figure 2. Aphids (mean $\pm$ SE) abundance in wheat fields near (25 m) and far (525 m) from the edge of the forest fragments in different wheat growth stages (T = tillering; B = boot; F = flowering; G = grain). Parana State, crop season (May to September) 2009. (* $p < 0.05$, Student's *t* test).

Unlike the present study, Alves et al. (2005) reported highest aphid infestation before the reproductive growth of wheat. However, in that

case, *R. padi* was the predominant species. Therefore, the higher infestation of *S. avenae* in the present study is probably associated with decreased abundance of *R. padi* related to wheat growth. This promotes competition among the species, favouring *S. avenae* development during grain stage of wheat (Gianoli, 2000).

In general, the infestation of aphids was not associated with the distance from the forest fragment and varies according to the study site. For example, differences were observed only in the grain stage, and in Ibiporã (SAF), a higher abundance of aphids was recorded far (525 m) from the edge of the forest, whereas in Londrina, a higher infestation of the pest was reported near (25 m) to the edge of the forest (Figure 2). This fact indicates that aphids are not dependent of refuge area, as opposed of the initial abundance of hoverflies reported in this study.

Is important to emphasize that both aphid species present in wheat fields, *R. padi* and *S. avenae* are prey to *A. exotica* (Rojo, Gilbert, Marcos-García, Nieto, & Durante, 2003; Bokina, 2012), which can contribute to highest abundance of this hoverfly specie in wheat fields (Table 1). In addition, some studies have reported that the hoverflies *T. watsoni* and *T. dispar* usually occur less frequently than *A. exotica* in open fields (Greco, 1995; Arcaya, Mengual, Bañon-Pérez, & Rojo, 2013), and that this species are probably less adapted to agroecosystems. One important factor that needs attention is that *A. exotica* has been reported in the Brazilian agroecosystem to prey on many aphids in oleraceous and fruit gardens (Resende et al., 2006; Sturza, Dorfey, Poncio, Dequech, & Bolzan, 2011) and to have other prey, such as mites, thrips and newly hatched caterpillars (Rojo et al., 2003). Thus, we believe that the wide spectrum of preys of *A. exotica* is likely the main reason for its widespread occurrence in all areas and its adaptability to local agroecosystems because of food availability during the larval and adult stages. Secondly, it is necessary to consider the influence of climatic factors on hoverflies, because the microclimate in wheat fields can favour *A. exotica* in relation to other hoverflies species. However, there is very little information about the bioecology of these aphid predators and additional studies are necessary to verify this hypothesis.

Although flowers cultivated efficiently increases aphidophagous hoverflies around crop fields (Amaral et al., 2013; Gontijo et al., 2013; Martínez-Uña et al., 2013; Haenke et al., 2014), in the present work, is important to emphasize that we did not sow strip flowers; thus, our data strongly indicate the relevance of preserving native forest fragments to

the survival of aphidophagous hoverflies during low aphid infestation in wheat crops. This report is supported by another study in Paraná State, which was developed by Marinoni, Miranda, and Thompson (2004). Their study found a larger abundance of *Allograpta* and *Toxomerus* in hedge forest than open fields (pasture), which indicates the availability of beneficial resources to aphidophagous hoverflies.

The other factor is probably the presence of alternative prey for hoverfly larvae near the hedge. In this sense, this relation was previously reported by Koh and Holland (2015), which showed that Anthocoridae predators were in soybean crops before aphid infestation and survived because of the presence of alternative prey and floral resources or weeds. In the present study, during wheat tillering in Londrina, for example, only four aphids were quantified in the wheat fields, which indicates that the larvae of aphidophagous hoverflies can survive by feeding on other insects such as other aphid species and caterpillars, in some cases, according to other studies (Rojo et al., 2003; Sturza et al., 2011). This fact reinforces the hypothesis of the importance of forest fragments to increase or maintain hoverfly abundance near the hedge during low aphid infestation in the fields.

Furthermore, the beneficial effect of forest fragments or refuge areas can vary according to the studied taxa or the local climate. In this sense, Raymond et al. (2015) reported a higher abundance of non-aphidophagous hoverflies in the hedge forest, whereas aphidophagous hoverflies were more abundant in crops, which are associated with a high availability of prey. However, in this study, we found a prevalence of aphidophagous predators in general, which was associated with the aphid abundance, but they were initially more abundant near the hedge forest fragment. Thus, the hedge "effect" on hoverflies was reported only in wheat tillering, and the distribution varied among the fields after this wheat growth stage, likely because of the expected relation with aphid infestation (Table 3). The exception was found near (25 m) the edge in one field (Londrina) (Table 3). Although this result is not clear, we hypothesized that another wheat field, which was sown later than the assessed field at approximately a distance far (525 m) from the edge, affected the hoverfly behavior and likely contained better-quality available aphids for consumption.

The importance of hoverflies in wheat fields can be attributed due female hoverflies strongly respond to aphid infestation and can locate even small colonies of the pest (Almohamad, Verheggen, & Haubruge, 2009). In this sense, several studies have shown that

hoverflies are highly affected by the presence of infochemicals. For example, Leroy et al. (2014) found that the *honeydew* excreted by *Acyrtosiphum pisum* (Hemiptera: Aphididae) increased the oviposition of *Epyrisphus balteatus* (Diptera: Syrphidae). In addition, other studies have shown that even in the absence of *honeydew*, the presence of aphids stimulates hoverfly oviposition and caïromone emission from the infested plants (Francis, Martin, Lognay, & Haubruge., 2005; Harmel et al., 2007; Verheggen, Arnaud, Bartram, Gohy, & Haubruge, 2008). Thus, although forest fragments can aid in hoverfly survival during low aphid infestation in the field, our data indicate that when the aphid infestation increases in wheat fields, these infochemical signals stimulate aphidophagous hoverflies to forage preys and those floral resources can be less important at this time.

Table 3. Relationship between aphid abundance and hoverflies near and far from the forest edge. Regression polynomial quadratic test. Paraná, wheat season (May to September of 2009).

Farm	Distance from edge					
	Near (25 m)			Far (525 m)		
	F	p	R ²	F	p	R ²
Ibiporã (SAF)	4.39	< 0.01	0.36	5.38	< 0.01	0.42
Londrina	0.56	n.s	0.08	6.24	< 0.01	0.48
Ibiporã (BF)	2.26	0.03	0.20	2.51	0.03	0.25

*Rolândia (GF) was not possible for a statistical analysis because of the low aphid and hoverfly abundances.

Is important to consider that many natural enemies are present in agroecosystem and contribute with aphid suppression. Although the present study showed hoverflies associated with aphid abundance (Table 3), other biological control agents acted against aphid infestation also. In Brazil, after implementation of biological control of aphids of wheat (BCAW) program, parasitoids are mainly responsible for reducing aphid infestation (Salvadori & Salles, 2002; Bortolotto et al., 2012). In the present work, we report an average parasitism between 6.8 and 16.9% among the fields. Another natural enemy associated with reduced aphid abundance was the predator Dolichopodidae ($R^2 = 0.37$ to 0.56 ; $p \leq 0.05$), supporting the importance of natural biological control against this pest in wheat fields. So, these natural enemies reported in this study can justify the weak influence of hoverflies on aphid abundance.

Although this subject is not the main objective of the study, the forest fragment quality was not associated with the abundance and richness of hoverflies. This result was observed in Ibiporã (SAF), which had the largest number of individuals and the largest richness, although it contained a forest section with intermediate biotic quality

(Table 1). Our data indicate that the abundance can be explained by the fact that this area also had the highest infestation of aphids, which suggests a density-dependent relationship. However, the hoverfly richness must be linked to other factors that were not measured in the study, such as the diversity of plants on site in the agroecosystem, environmental complexity on a regional scale (Fahrig et al., 2015) and historical management of crops (input of pesticides) (Bokina, 2012). Thus, these factors must be investigated in other studies to better understand the relation between hoverflies and the environment.

Finally, this study reports for first time the importance of preserving areas of forest fragments as a refuge and likely source of supplies for aphidophagous hoverflies in Brazilian agroecosystems. There is currently a great necessity for similar studies in developing tropical regions because of the lack of such information. Thus, other aspects should be considered for further studies, such as identifying the weeds in the vicinity of the crop to select potential plants to be grown to attract these natural enemies in regions with poor natural biodiversity.

Conclusion

Hoverfly abundances are initially higher near the edge of the forest fragment. After the aphid abundance in wheat fields increases, the hoverfly population depends on the prey infestation.

Acknowledgements

We are grateful to Dra Mirian Morales for assisting in identification of insects and Capes and CNPq for providing the post-graduate scholarship.

References

- Almohamad, R., Verheggen, F. J., & Haubruge, E. (2009). Searching and oviposition behavior of aphidophagous hoverflies (Diptera: Syrphidae): a review. *Biotechnology, Agronomy, Society and Environment*, 13(3), 467-481.
- Alves, L. F. A., Prestes, T. M. V., Zanini, A., Dalmolin, M. F., & Menezes Jr., A. O. (2005). Controle biológico natural de pulgões (Hemiptera: Aphididae) em lavoura de trigo por parasitoides (Hymenoptera, Aphididae), no município de Medianeira, PR, Brasil. *Semina: Ciências Agrárias*, 26(2), 147-152.
- Amaral, D. S. S. L., Venzon, M., Duarte, M. V. A., Sousa, F. F., Pallini, A., & Harwood, J. D. (2013). Non-crop vegetation associated with chili pepper agroecosystems promote the abundance and survival of aphid predators. *Biological Control*, 64(3), 338-346.
- Arcaya, E., Mengual, X., Bañon-Pérez, C., & Rojo, S. (2013). Registros y distribución de sírfidos depredadores (Diptera: Syrphidae: Syrphinae) en el estado Lara, Venezuela. *Bioagro*, 25(2), 143-148.
- Barbosa, P. (1998). *Conservation biological control*. San Diego, CA: Academic Press.
- Blaauw, B. R., & Isaacs, R. (2014). Flower plantings increase wild bee abundance and the pollination services provided to a pollination-dependent crop. *Journal of Applied Ecology*, 51(4), 890-898.
- Bokina, I. G. (2012). Hoverflies (Diptera, Syrphidae) in agroecosystems of the forest-steppe of western Siberia and the influence of agrotechnological practice on their abundance. *Entomological Review*, 92(s/n), 1053-1060.
- Borges, Z. M., & Couri, M. (2009). Revision of *Toxomerus* Macquart, 1855 (Diptera: Syrphidae) from Brazil with synonymic notes, identification key to the species and description of three new species. *Zootaxa*, 2179(s/n), 1-72.
- Bortolotto, O. C., Menezes Jr., A., & Hoshino, A. T. (2015). Aphidophagous parasitoids can forage wheat crops before aphid infestation, Paraná State, Brazil. *Journal of Insect Science*, 15(40), 1-4.
- Bortolotto, O. C., Menezes Jr., A. O., Sampaio, M. V., & Hoshino, A. T. (2012). Aphid parasitoids that occur on wheat crops of Northeast of the Parana State with recovery of *Aphidius rhopalosiphii* in Brazil. *Semina: Ciências Agrárias*, 33(suppl. 2), 3075-3080.
- Bugg, R. L., Colfer, R. G., Chaney, W. E., Smith, H. A., & Cannon, J. (2008). *Flower flies (Syrphidae) and other biological control agents for aphids in vegetable crops*. Oakland, CA: University of California Agriculture and Natural Resources.
- Dib, H., Simon, S., Sauphanor, B., & Capowiez, Y. (2010). The role of natural enemies on the population dynamics of the rosy apple aphid, *Dysaphis plantaginea* Passerini (Hemiptera: Aphididae) in organic apple orchards in south-eastern France. *Biological Control*, 55(2), 97-109.
- Fahrig, L., Girard, J., Duro, D., Pasher, J., Smith, A., Javorek, S., ... Tishendorf, L. (2015). Farmlands with smaller crop fields have higher within-field biodiversity. *Agriculture, Ecosystems and Environment*, 200(1), 219-234.
- Földesi, R. F., & Kovács-Hostyánszki, A. (2014). Hoverfly (Diptera: Syrphidae) community of a cultivated arable field and the adjacent hedgerow near Debrecen, Hungary. *Biologia*, 69(3), 381-388.
- Fearnside, P. M. (2001). Soybean cultivation as a threat to the environment in Brazil. *Environmental Conservation*, 28(1), 23-38.
- Francis, F., Martin, T., Lognay, G., & Haubruge, E. (2005). Role of (E)- β -farnesene in systematic aphid prey location by *Episyrphus balteatus* larvae (Diptera:

- Syrphidae). *European Journal of Entomology*, 102(3), 431-436.
- Gianoli, E. (2000). Competition in cereal aphids (Homoptera: Aphididae) on wheat plants. *Environmental Entomology*, 29(2), 213-219.
- Gontijo, L. M., Beers, E. H., & Snyder, W. E. (2013). Flowers promote aphid suppression in apple orchards. *Biological Control*, 66(1), 8-15.
- Greco, C. F. (1995). Phenology and habitat selection of the aphidophagous syrphid species most frequently found in crops and pastures in the Province of Buenos Aires, Argentina. *Entomophaga*, 40, 317-320.
- Haenke, S., Kovács-Hostyánszki, A., Fründ, J., Batáry, P., Jauker, B., Tschamtker, T., & Holzschuh, A. (2014). Landscape configuration of crops and hedgerows drives local syrphid fly abundance. *Journal of Applied Ecology*, 51(2), 505-513.
- Harmel, N., Almohamad, R., Fauconnier, M. L., Du, J. P., Verheggen, F., Marlier, M., ... Francis, F. (2007). Role of terpenes from aphid-infested potato on searching and oviposition behavior of the hoverfly predator *Episyrphus balteatus*. *Insect Science*, 14(1), 57-63.
- Hickman, J. M., & Wratten, S. D. (1996). Use of *Phacelia tanacetifolia* strips to enhance biological control of aphids by hoverfly larvae in cereal fields. *Journal of Economic Entomology*, 89(4), 832-840.
- Hogg, B. N., Bugg, R. L., & Daane, K. M. (2011). Attractiveness of common insectary and harvestable floral resources to beneficial insects. *Biological Control*, 56(1), 76-84.
- Irshad, M. (2014). Review: role of syrphids (Diptera: Syrphidae) as biotic agents and pollinators in Pakistan. *Journal of Bioresource Management*, 1(2), 1-9.
- Jun, S., Yibo, L., Jin, C., Fazhi, S., & Zhenhai, D. (2009). The pollination of *Paphiopedilum hirsutissimum*. *Orchid Review*, 117(1286), 78-81.
- Koh, I., & Holland, J. D. (2015). Grassland plantings and landscape natural areas both influence insect natural enemies. *Agriculture, Ecosystems and Environment*, 199(s/n), 109-199.
- Laubertie, E. A., Wratten, S. W., & Hemptinne, J.-L. (2012). The contribution of potential beneficial insectary plant species to adult hoverfly (Diptera: Syrphidae) fitness. *Biological Control*, 61(1), 1-6.
- Leroy, P. D., Almohamad, R., Attia, S., Capella, Q., Verhegg, F. J., Haubruge, E., & Francis, F. (2014). Aphid honeydew: An arrestant and a contact kairomone for *Episyrphus balteatus* (Diptera: Syrphidae) larvae and adults. *European Journal of Entomology*, 111(2), 237-242.
- Letourneau, D., Bothwell, A. S., & Stireman, J. (2012). Perennial habitat fragments, parasitoid diversity and parasitism in ephemeral crops. *Journal of Applied Ecology*, 49(6), 1405-1416.
- Marinoni, L., Miranda, G. F. G., & Thompson, F. C. (2004). Abundância e riqueza de espécies de Syrphidae (Diptera) em áreas de borda e interior de floresta no Parque Estadual de Vila Velha, Ponta Grossa, Paraná, Brasil. *Revista Brasileira de Entomologia*, 48(4), 554-559.
- Marinoni, L., Morales, M. N., & Spaler, I. (2007). Chave de identificação ilustrada para os gêneros de Syrphinae (Diptera, Syrphidae) de ocorrência no sul do Brasil. *Biota Neotropica*, 7(1), 145-160.
- Martínez-Uña, A., Martín, J. M., Fernández-Quintanilla, C., & Dorado, J. (2013). Provisioning floral resources to attract aphidophagous hoverflies (Diptera: Syrphidae) useful for pest management in central Spain. *Journal of Economic Entomology*, 106(6), 2327-2335.
- Medeiros, H. R., & Torezan, J. M. (2013). Evaluating the ecological integrity of Atlantic forest remnants by using rapid ecological assessment. *Environmental Monitoring and Assessment*, 185(5), 4373-4382.
- Mengual, X., Ruiz, C., Rojo, S., Stahl, G., & Thompson, F. C. (2009). A conspectus of the flower fly genus *Allograpta* (Diptera: Syrphidae) with description of a new subgenus and species. *Zootaxa*, 2214(s/n), 1-28.
- Murta, A. F., Ker, F. T. O., Costa, D. B., Espírito-Santo, M. M., & Faria, M. L. (2008). Efeitos de remanescentes de Mata Atlântica no controle biológico de *Euselasia apisaon* (dahman) (Lepidoptera: Riodinidae) por *Trichogramma maxacalii* (Voegelé e Pointel) (Hymenoptera: Trichogrammatidae). *Neotropical Entomology*, 37(2), 229-232.
- Nassar, A. M. (2009). Brazil as an Agricultural and Agroenergy Superpower. In L. Brainard, & L. Martínez-Díaz (Eds.), *Brazil as an economic superpower?* (p. 55-80). Washington, DC: The Brookings Institution.
- Petanidou, T., Vuji, A., & Ellis, W. N. (2011). Hoverfly diversity (Diptera: Syrphidae) in a Mediterranean scrub community near Athens, Greece. *Annales de la Société entomologique de France (N.S.)*. *International Journal of Entomology*, 47(1-2), 168-175.
- Pinheiro, L. A., Torres, L. M., Raimundo, J., & Santos, S. A. P. (2015). Effects of pollen, sugars and honeydew on lifespan and nutrient levels of *Episyrphus balteatus*. *Biocontrol*, 60(1), 47-57.
- Raymond, L., Sarthou, J.-P., Plantegenest, M., Gauffre, B., Ladet, S., & Vialatte, A. (2015). Immature hoverflies overwinter in cultivated fields and may significantly control aphid populations in autumn. *Agriculture, Ecosystems and Environment*, 185(1), 99-105.
- Resende, A. L. S., Silva, E. E., Silva, V. B., Ribeiro, R. L. D., Guerra, J. G. M., & Aguiar-Menezes, E. L. (2006). Primeiro registro de *Lipaphis pseudobrassicae* Davis (Hemiptera: Aphididae) e sua associação com insetos predadores, parasitóides e formigas em couve (Cruciferae) no Brasil. *Neotropical Entomology*, 35(4), 551-555.
- Rojo, S., Gilbert, F., Marcos-García, M. A., Nieto, J. M., & Durante, M. P. M. (2003). *A world review of predatory hoverflies (Diptera, Syrphidae: Syrphinae) and their prey*. Alicante, ES: Centro Iberoamericano de la Biodiversidad.

- Salvadori, J. R., & Salles, L. A. B. (2002). Controle biológico dos pulgões do trigo. In J. R. P. Parra, P. S. M. Botelho, B. S. C. Ferreira, & J. M. S. Bento (Eds.), *Controle biológico no Brasil: parasitóides e predadores* (p. 427-447). São Paulo, SP: Manole.
- Salvadori, J. R., & Tonet, G. E. L. (2001). *Manejo integrado dos pulgões de trigo*. Passo Fundo, RS: Embrapa- CNPT.
- Schmidt, M. H., Lauer, A., Purtauf, T., Thies, C., Schaefer, M., & Tschardt, T. (2003). Relative importance of predators and parasitoids for cereal aphid control. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 270(1527), 1905-1909.
- Sturza, V. S., Dorfey, C., Poncio, S., Dequech, S. T. B., & Bolzan, A. (2011). First record of larvae of *Allograpta exotica* Wiedemann (Diptera, Syrphidae) preying on *Aphis gossypii* Glover (Hemiptera, Aphididae) in watermelon in Brazil. *Revista Brasileira de Entomologia*, 55(2), 272-274.
- Thompson, F. C. (1999). A key to the genera of the flower flies (Diptera: Syrphidae) of the neotropical region including descriptions of new genera and species and a glossary of taxonomic terms. *Contributions on Entomology International*, 3, 319-378.
- Van Rijn, P. C. J., Kooijman, J., & Wackers, F. L. (2013). The contribution of floral resources and honeydew to the performance of predatory hoverflies (Diptera: Syrphidae). *Biological Control*, 67(1), 32-38.
- Verheggen, F. J., Arnaud, L., Bartram, S., Gohy, M., & Haubruge, E. (2008). Aphid and plant volatiles induce oviposition in an aphidophagous hoverfly. *Journal of Chemical Ecology*, 34(3), 301-307.
- Vieira, I. C. G., Toledo, P. M., Silva, J. M. C., & Higuchi, H. (2008). Deforestation and threats to the biodiversity of Amazonia. *Brazilian Journal of Biology*, 68(4), 949-956.
- Walton, N. J., & Isaacs, R. (2011). Influence of native flowering plant strips on natural enemies and herbivores in adjacent blueberry fields. *Environmental Entomology*, 40(3), 697-705.

Received on May 7, 2015.

Accepted on August 10, 2015.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.