

In vitro germination and disinfestation of sweet cactus (*Nopalea cochenillifera* (L.) Salm Dyck)

Juliana Pereira de Castro¹, Emmanuelle Rodrigues Araújo^{2*}, Mailson Monteiro do Rêgo¹ and Elizanilda Ramalho do Rêgo¹

¹Departamento de Fitotecnia, Centro de Ciências Agrárias, Universidade Federal da Paraíba, Areia, Paraíba, Brazil.

²Universidade Federal de Pernambuco, Rua Dom Manoel de Medeiros, s/n, 52171-900, Recife, Pernambuco, Brazil.

*Author for correspondence. E-mail: manucg@gmail.com

ABSTRACT. This work aimed to promote *in vitro* seed germination and disinfestation of the sweet cactus. Seeds were submerged in alcohol at 70% for 1 min. and then treated with sodium hypochlorite solution at different concentrations: 0.0; 0.5; 1.0 and 1.5%. After this treatment, the seeds were washed four times in distilled, deionized and autoclaved water. Seeds were inoculated in MS medium with different concentrations of sucrose (0.0, 2.5, 5.0, 7.5 and 10.0 g L⁻¹). The experimental design was completely randomized in a factorial experiment 4 x 5, with five replicates for treatments. Contamination, germination and seedling growth were evaluated. The results of the analysis of variance indicate that there were no significant interactions among factors. There was no significance for sucrose concentrations on contamination but these treatments showed significant differences for seed germination and seedling growth ($p \leq 0.01$). At higher sucrose concentrations there was lower germination and smaller seedlings. There were significant differences between the treatments with sodium hypochlorite for all traits analysed ($p \leq 0.01$). There was low contamination with the increase in sodium hypochlorite concentrations with higher germination percentage and taller seedlings.

Keywords: Cactaceae, forage, tissue culture.

RESUMO. Desinfestação e germinação de sementes de palma doce (*Nopalea cochenillifera* (L.) Salm Dyck) *in vitro*. Este trabalho objetivou desinfestar e promover a germinação de sementes de palma doce “*in vitro*”. As sementes foram mergulhadas em álcool a 70% por 1 min. e, posteriormente, imersas em solução de hipoclorito de sódio nas concentrações de 0,0; 0,5; 1,0 e 1,5% com três gotas de Tween 20 por 10 min. e lavadas em quatro águas bidestiladas estéreis. As sementes foram cultivadas em meio MS suplementado com 0,0; 2,5; 5,0; 7,5 e 10,0% de sacarose. O delineamento experimental foi inteiramente casualizado, em esquema fatorial 4 x 5 (quatro concentrações de hipoclorito de sódio e cinco concentrações de sacarose), com cinco repetições. Avaliou-se a contaminação, germinação e crescimento da plântula. Não houve interação significativa entre os fatores, havendo porém, diferenças significativas entre os tratamentos de hipoclorito e concentrações de sacarose ($p \leq 0,01$). As concentrações de sacarose não tiveram efeito significativo na contaminação do meio, porém com o aumento dos teores de sacarose houve tendência ao decréscimo na germinação e crescimento da plântula ($p \leq 0,01$). Quanto maior a concentração de hipoclorito no meio menor foi a contaminação e maior a germinação e comprimento da plântula ($p \leq 0,01$).

Palavras-chave: Cactaceae, forrageira, cultura de tecido.

Introduction

Nopalea cochenillifera (L.) Salm Dyck is known in Brazil as Small Palm or Sweet Palm and is cultivated in the Brazilian semi-arid northeast, mainly by the dairy cattle ranchers, and the largest crop areas are found in the States of Alagoas, Pernambuco and Paraíba. It has been used as a food supply, in the drought period by small farmers (MAIA NETO, 2003).

In long-term drought, the ranchers cultivate cacti due to its high water content vital to livestock rising, and ranchers feed them through all year round, as in the same way as *Opuntia ficus-indica* (SANTOS et al., 1996).

Maintenance collections of succulent plants can be problematic since many of these species are very susceptible to rots by bacteria and fungi. Although their seeds could be stored at low relative humidity and temperature, these microorganisms

can affect the seeds germination vigor and can promote the anormal growing of plantlets in tissue culture. Methods for seed disinfestation in micropropagation of cacti have been developed to overcoming these problems and to obtain healthy plants (COUTO et al., 2004).

Disinfestations of *Opuntia* seeds are necessary when the germination is performed *in vitro*. In such case, cultures are supplemented by a source of sugar and salts in order to obtain energy and other requirements. The nutritional requirements by explants *in vitro* depend on the species, varieties and tissue (CALDAS et al., 1998). Reis et al. (2009) showed the influence of the culture medium in the content and chemical composition of the essential oil of *Melissa officinalis*. Sucrose is included in culture media to promote rapid growth (CALDAS et al., 1998; MALDANER et al., 2006). In tissue culture, the use of sugar is necessary, sometimes due to the low photosynthetic rate as presented by orchids and cactus. However, the presence of sugar and salts easily allows the development of unwanted microorganisms. To avoid such phenomenon, an efficient procedure must be done as a pretreatment of the seeds with liquid or gaseous substances (ALVAREZ-PARDO et al., 2006).

The liquid disinfectant should remove or kill all microorganisms from the seeds' surface without damaging the embryos. For the purpose of disinfesting seeds, the calcium hypochlorite is the most used one, besides hydrogen peroxide and sodium hypochlorite. Sodium and calcium hypochlorite are prescribed to be used at different concentrations and times of exposure to disinfest seeds (ARDITTI; ERNST, 1984; PIERRIK, 1997).

The present investigation was carried out with the objectives to promote *in vitro* seed germination, disinfestation and plant growth of the sweet cactus.

Material and methods

Plant Material

Seeds were obtained from mature fruits from Galant District, Paraíba State, Brazil. Surface sterilization was realized in the Plant Biotechnology Laboratory at Universidade Federal da Paraíba.

Surface Sterilization Treatments

On laminar flowhood using a Becker, the ripe fruits were surface sterilized for 1 min. in 70% ethanol; followed rinsing with sterile deionized water and dipped in a commercial sodium hypochlorite solution at different concentrations:

0.0; 0.5; 1.0 and 1.5% for 10 minutes, established based on previous experiment. After this treatment, the seeds were washed four times in distilled, deionized and autoclaved water and placed on a paper filter to dry off the excess water.

Culture Medium Treatments

Seeds were inoculated in MS medium (MURASHIGE; SKOOG, 1962) with different concentrations of sucrose (0.0; 2.5; 5.0; 7.5 and 10.0 g L⁻¹) plus 0.8 % of agar. The pH was adjusted to 5.7 with NaOH. Ten milliliters of medium were autoclaved for 20 minutes at 120°C in glass tubes. The tubes were tilted; thus, when the medium cooled down a larger surface was formed, the seeds were sown and kept in the dark during 15 days at 25±2°C and then moved to a room with 16 hours of light exposure with a luminosity of 30 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and 8 hours under darkness. The swollen seeds with green embryos were observed with a stereomicroscope and they were considered as germinated when presented plantlets (HAILES; SEATON, 1989).

Data analysis

The data were submitted to the square root transformation like proposed by Bartlett (1936). The experimental design was completely randomized and the data were analysed in a factorial experiment 4 x 5, with five replicates for treatments. The following variables were evaluated: contamination, germination seed frequency and seedling growth. The results were submitted to statistical ANOVA analysis and followed by regression analysis ($p \leq 0.01$). The statistical analyses were done using Genes software (CRUZ, 2001).

Results and discussion

The results of ANOVA showed that there was no significantly interaction between sucrose concentrations and sodium hypochlorite concentrations (Table 1). This indicates that the factors affect the response independently and should be analyzed separately. Thus, the regressions analysis for all treatments in a factor should be examined as resulted from a single-factor experiment (Figures 1 and 2).

There were significant differences between the treatments with sodium hypochlorite for all traits analyzed ($p \leq 0.01$) (Table 1). The disinfestations treatments 0.5; 1.0 and 1.5% were effective in to control the contamination. The germination

increased with the disinfection with sodium hypochlorite and started at seventh day after the inoculation of the seeds (Figure 1).

Table 1. Analysis of variance of contamination, germination and seedling growth in Sweet cactus (*Nopalea cochenillifera*) *in vitro*.

Source of variation	D.F.	Mean Square		
		Contamination	Germination	Seedling growth
Sucrose (S)	4	0.004 ^{ns}	0.258 ^{**}	32.43 ^{**}
Sodium Hypochlorite (NaHClO)	3	1.424 ^{**}	0.338 ^{**}	21.58 ^{**}
(NaHClO) x S	12	0.01 ^{ns}	0.037 ^{ns}	3.453 ^{ns}
Residue	80	0.01	0.042	3.944
Total	99			

^{**}Significant ($p \leq 0.01$) by F statistics; ^{ns}no significant.




Figure 1. Regression analysis of contamination, germination and seedling growth in Sweet cactus (*Nopalea cochenillifera*) *in vitro*, with different sodium hypochlorite concentrations.




Figure 2. Regression analysis of germination and seedling growth in Sweet cactus (*Nopalea cochenillifera*) *in vitro*, with different sucrose concentrations.

We verified a higher germination percentage and taller seedlings with the increase in sodium hypochlorite concentrations (Figure 1). The beginning of contamination was detected at third day after the inoculation of seeds. According to Barrueto Cid and Zimmermann (2006), the sodium hypochlorite is the most efficient product on disinfection of explants. Golle et al. (2010) working with *Pinus taeda* L. seeds and using sodium hypochlorite at several concentrations in different periods for *in vitro* disinfection founded the combination of NaOCl at 3% for 5 minutes provided a good disinfection but inhibited the seeds germination.

There was no significance for sucrose concentrations on contamination but these treatments showed significant differences for seed germination and seedling growth ($p \leq 0.01$) (Table 1). The germination and the seedling growth decreased with the increase in sucrose concentration (Figure 2). The same was observed by Leal et al. (2005), studying the germination and development of *Pitaya* with different concentrations of sucrose. Nevertheless, Dávila-Figueroa et al. (2005) observed significant effects of 3% sucrose for *in vitro* propagation of *Turbinicarpus*. Unlike, Da Luz et al. (2008) found no differences for *in vitro* pollen germination using different sucrose concentrations.

Conclusion

Sodium hypochlorite was able to decrease the contamination in sweet cactus *in vitro* germination, and seedling growth; and the higher sucrose concentration reduced the seed germination and seedling growth. Data are also presented regarding the disinfection, and asymbiotic seed germination represents an efficient way to propagate this species.

Acknowledgements

The research present in this paper was funded by research grants from CAPES, CNPq and FINEP. We would like to thank to Elizabel de Souza Ramalho Viana and Fernando Luiz Finger for their insightful feedback on a preliminary version of this paper.

References

- ALVAREZ-PARDO, V. M.; FERREIRA, A. G.; NUNES, V. F. Seed disinfection methods for *in vitro* cultivation of epiphyte orchids from Southern Brazil. **Horticultura Brasileira**, v. 24, n. 2, p. 217-220, 2006.
- ARDITTI, J.; ERNST, R. Physiology of germinating orchid seeds. In: ARDITTI, J. (Ed.). **Orchid biology: reviews and perspectives III**. New York: Cornell University Press, 1984. p. 177-222.
- BARRUETO CID, L. P.; ZIMMERMANN, M. J. **A contaminação in vitro de plantas**. Brasília: Embrapa Recursos Genéticos e Biotecnologia, 2006. (Boletim de Pesquisa e Desenvolvimento).
- BARTLETT, M. S. The square root transformation in analysis of variance. **Journal of Statistics Society**, v. 3, suppl., p. 68-78, 1936.
- CALDAS, L. S.; HARIDASAN, P.; FERREIRA, M. E. Meios Nutritivos. In: TORRES, A. C.; CALDAS, L. S.; BUSO, J. A. (Ed.). **Cultura de tecido e transformação genética**. Brasília: Embrapa-SPI; Embrapa-CNPq, 1998. v. 1. p. 87-132.
- COUTO, J. M. F.; OTONI, W. C.; PINHEIRO, A. L.; FONSECA, E. P. Desinfestação e germinação *in vitro* de

- sementes de mogno (*Swietenia macrophylla* King). **Revista Árvore**, v. 28, n. 5, p. 633-642, 2004.
- CRUZ, C. D. **Programa GENES** (versão Windows), aplicativo computacional em genética e estatística. Viçosa: UFV, 2001.
- DA LUZ, C. L.; SCHUELTER, A. R.; DA LUZ, C. L.; DAL'MASOL, A.; VIEIRA, E. S. N.; BARRETO, R. R. Germinação *in vitro* de grãos de pólen e efeito da proteção das plantas na frutificação de cubiu (*Solanum sessiliflorum* anum Dunal). **Acta Scientiarum. Agronomy**, v. 30, n. 4, p. 539-545, 2008.
- DÁVILA-FIGUEROA, C. A.; ROSA-CARRILLO, M. L.; PÉREZ-MOLFHE-BALCH, E. *In vitro* propagation of eight species or subspecies of *Turbinicaps* (Cactaceae). **In Vitro Cellular and Developmental Biology – Plant**, v. 41, n. 4, p. 540-545, 2005.
- GOLLE, D. P.; REINIGER, L. R. S.; CURTI, A. R.; HANAUER, J. G.; WALDOW, D. A. G. Substratos alternativos e tratamentos pré-germinativos na germinação *in vitro* de sementes de *Pinus taeda* L. **Revista Árvore**, v. 34, n. 1, p. 39-48, 2010.
- HAILES, N. S. J.; SEATON, P. T. The effects of composition of the atmosphere on the growth of seedlings of *Cattleya aurantica*. In: PRITCHARD, H. W. (Ed.). **Modern methods in orchid conservation: the role of physiology, ecology and management**. Cambridge: Cambridge University Press, 1989. p. 73-85.
- LEAL, E. A.; SANTOS, A. M.; LINS, C. Q. G.; SOUZA, A. D. Germinação e desenvolvimento de Pitaya (*Hylocereus undatos*) *in vitro*. **Arquivos do Instituto de Biologia**, v. 72, supl. 1, p. 1-63, 2005.
- MAIA NETO, A. L. Utilização da palma forrageira para produção de leite no semi-árido nordestino. **Bahia Agrícola**, v. 5, n. 3, p. 45-49, 2003.
- MALDANER, J.; NICOLOSO, F. T.; FLORES, R.; SKREBSKY, E. C. Sacarose e nitrogênio na multiplicação de *Pfaffia glomerata* (spreng) Pedersen. **Ciência Rural**, v. 6, n. 4, p. 1201-1206, 2006.
- MURASHIGE, T.; SKOOG, F. A. A revised medium for rapid growth and bioassays with tobacco tissue culture. **Physiologia Plantarum**, v. 15, n. 3, p. 473-497, 1962.
- PIERRIK, R. L. M. **In vitro cultures of higher plants**. Dordrecht: Martinus Nijhoff Publishers, 1997.
- REIS, E. S.; PINTO, J. E. B. P.; ROSADO, L. D. S.; CORRÊAL, R. M. Teor e composição química do óleo essencial de *Melissa officinalis* L. *in vitro* sob influência do meio de cultura. **Acta Scientiarum. Agronomy**, v. 31, n. 2, p. 331-335, 2009.
- SANTOS, D. C.; FARIAS, L.; LIRA, M. A.; FERNANDES, A. P. M.; FREITAS, E. V.; MORENO, J. A. Produção e composição química da palma forrageira cv. Gigante (*Opuntia ficus-indica* Mill) sob adubação e calagem, no Agreste Semiárido de Pernambuco. **Pesquisa Agropecuária Pernambucana**, v. esp., n. 9, p. 69-78, 1996.

Received on February 4, 2009.

Accepted on August 2, 2009.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.