

## Numerical Solution of Non-linear Volterra Integral Equation of the First Kind

Boutheina Tair, Mourad Ghait, Hamza Guebbai, Mohemd Zine Aissaoui

**ABSTRACT:** In this paper, we focus on the numerical solution of a nonlinear Volterra equation of the first kind. The existence and uniqueness of the exact solution are ensured under a necessary condition which we present next. We develop a numerical method based on two essential parts which are linearization and discretization. We start with the discretization of the equations using the concept of Nystrom’s method and for the linearization we apply Newton’s method. We present theorems that show the convergence of the proposed method. At the end, numerical examples are provided to show the efficiency of our method.

**Key Words:** Volterra integral equation, non-linear integral equation, numerical integration, Newton method.

### Contents

|  |          |
|--|----------|
| <b>1 Introduction</b>  | <b>1</b> |
| <b>2 Problem Position</b>  | <b>2</b> |
| <b>3 Analytical Study</b>  | <b>2</b> |
| <b>4 Numerical Study</b>   | <b>3</b> |
| 4.1 Discretization . . . . .   | 3        |
| 4.1.1 Approximate solution convergence of discrete problem . . . . . | 5        |
| 4.2 Linearization . . . . .  | 6        |
| <b>5 Numerical Examples</b>  | <b>8</b> |
| <b>6 Conclusion</b>  | <b>9</b> |

### 1. Introduction

Volterra integral equations are the good choice for scientists to model many evolution problems. Historically, the emergence of this equation was to express population dynamics through Volterra’s most famous work, as it constituted a qualitative leap in modeling and applied mathematics [15], and since then, the use of this type of equations have expanded year after year. In year 1976 [9], it was the best solution for modeling population evolution, in 1988 it was used to express vibrational motion and was applied to express the sorption kinetics of mixtures in 1990, then we find it a year 1996 in semi conductor devices and then to express viscosity in 1999. The above was a slight review of some of the works presented in the last century and to date it has countless applications like the tumor growth [10,12], modeling the system related to leukemia [13], birth-death process [11] and relativistic quantum physics [14].

Because these equations forms varied according to the domain where they are applied, there are different types and shapes. In this manuscript, we focus on one particular type, which is the non-linear Volterra equation. This has been presented in many works in the following general form:

$$\forall x \in [0, X], \int_0^x \mathcal{K}(x, t, u(t)) dt = f(x), \quad X < +\infty, \quad (1.1)$$

where  $f$  is a given function in the Banach space  $C^1[0, X]$ .

---

2010 *Mathematics Subject Classification:* 34A12, 47H30, 65D30, 49M15.  
 Submitted April 08, 2022. Published June 29, 2022

The above equation first appeared in the work of Abel to express the motion of a particle along a series curve. For our part, we present some of the recent work in order to shed light on its great importance. The importance of this equation was not limited to a specific field or domain. We find it in engineering [1,3,4], in physics [6], in artificial network [7], in meteorology to express the amount of precipitation [8], in medicine, where it has been used recently to study the evolution of the Corona virus or to find the best model of it [2,5], the population dynamics and spread of epidemics [16]. The equation (1.1) is considered to be one of the problems that are ill-posed, according to Hadamard's definition [20]. Therefore, we find many researches concerned with the development of the regularization methods [21,22]. On the other side, there are many scientific papers that deal with the search for a numerical solution to this type of equation, in which many numerical methods have been invented, and we mention among them: Petryshyn's fixed point theorem [24], homotopy perturbation [25], discrete operational vector scheme [25], hp-vertion collocation method [27] and Nyström method [28,29,30,31,32,33]. In this manuscript, we recall sufficient conditions to prove the existence and uniqueness of the solution in the Banach space  $C^0[0, X]$ . For the method approach, we develop a numerical method based on two principle ideas of Nyström and Newton procedures. Therefore, we use some concepts of linearization and discretization that have been applied in the treatment of the nonlinear Fredholm integral equation of the second kind [34,35,36].

## 2. Problem Position

In this paper, we are interested in developing a method for finding the best possible approximate solution. But, we do not deal directly with the equation (1.1) for the aforementioned reason. Instead, we suggest that the kernel  $\mathcal{K}$  is derivable with respect to  $x$ . So, that we can convert the equation into a non-linear Volterra equation of second kind:

$$\forall x \in [0, X], \mathcal{K}(x, x, u(x)) = f'(x) - \int_0^x \frac{\partial \mathcal{K}}{\partial x}(x, t, u(t)) dt. \quad (2.1)$$

In order to make the presentation clearer and easy to read, let  $C^0[0, X]$  be a Banach space equipped with next norm

$$\forall v \in C^0[0, X], \|v\|_{C^0[0, X]} = \max_{0 \leq x \leq X} |u(x)|.$$

## 3. Analytical Study

Since we cannot invent a numerical solution without ensuring the existence and uniqueness of the exact solution, we assume that the kernel verifies the below hypotheses which are sufficient for ensuring this.

$$(H_1) \left\{ \begin{array}{l} (1) \quad f(0) = 0, \\ (2) \quad f \text{ and } f' \text{ in } C^0[0, X], \\ (3) \quad \mathcal{K} \text{ and } \frac{\partial \mathcal{K}}{\partial x} \in C^0\left([0, X]^2 \times \mathbb{R}\right), \\ (4) \quad \exists M > 0, \max_{0 \leq x, t \leq X} \left( |\mathcal{K}(x, t, v)|, \left| \frac{\partial \mathcal{K}}{\partial x}(x, t, v) \right| \right) \leq M, \\ (5) \quad \exists L > 0, \forall x, t \in [0, X], \forall v, \bar{v} \in \mathbb{R}: \left| \frac{\partial \mathcal{K}}{\partial x}(x, t, v) - \frac{\partial \mathcal{K}}{\partial x}(x, t, \bar{v}) \right| \leq L |v - \bar{v}|, \\ (6) \quad \forall y \in \mathbb{R}, \exists ! v : \mathcal{K}(x, x, v) = y, \\ (7) \quad \exists \theta > 0, \forall x, t, \forall v, \bar{v} \in \mathbb{R} : |\mathcal{K}(x, t, v) - \mathcal{K}(x, t, \bar{v})| \geq \theta |v - \bar{v}|. \end{array} \right.$$

Let us apply the successive approximation method or Picard's method [16] to equation (2.1). This consists in the construction of sequences  $\{u_n\}_{n \in \mathbb{N}}$  and  $\{\psi_n\}_{n \in \mathbb{N}}$  which are defined by

$$\begin{cases} u_0(x) = f'(x), \\ \mathcal{K}(x, x, u_n(x)) = f'(x) - \int_0^x \frac{\partial \mathcal{K}}{\partial x}(x, t, u_{n-1}(t)) dt, \quad \forall n \geq 1, \end{cases} \quad (3.1)$$

and

$$\begin{cases} \psi_0(x) = f'(x), \\ \psi_n(x) = u_n(x) - u_{n-1}(x), \quad \forall n \geq 1. \end{cases} \quad (3.2)$$

**Theorem 3.1.** *Under the hypotheses  $(H_1)$ , the equation (2.1) has a unique solution in  $C^0[0, X]$ .*

*Proof.* Details can be found in [16]. □

#### 4. Numerical Study

In the search for a numerical solution of the equation (1.1), we follow two essential steps: The first one is to discretize our problem (1.1) by replacing the integral sign by a numerical integration, whereas the second step is the linearization of the new approximate problem using Newton's method. In order to estimate the numerical error, we use the continuity module  $\kappa_0(\cdot, h)$  by

$$\forall v \in C^0[0, X], \kappa_0(v, h) = \sup_{|x-y| \leq h} |v(x) - v(y)|.$$

##### 4.1. Discretization

We define  $\Delta_n$ ,  $n \geq 1$  the uniform discretization of the interval  $[0, X]$  as:

$$\Delta_n = \left\{ n \geq 1, 0 = x_0 < x_1 < \dots < x_{n-1} < x_n = X, h = x_{j+1} - x_j, 0 \leq j \leq n \right\}. \quad (4.1)$$

By choosing the collocation points  $x = x_i$ , we obtain

$$\mathcal{K}(x_i, x_i, u(x_i)) + \int_0^{x_i} \frac{\partial \mathcal{K}}{\partial x}(x_i, t, u(t)) dt = f(x_i), \quad 0 \leq i \leq n. \quad (4.2)$$

Our main goal is searching  $u_i$  as an approximation of  $u(x_i)$ . So, we replace the integral sign in (4.2) by the following numerical Gaussian scheme:

$$\forall v \in C^0[0, X], \int_0^{x_i} v(x) dx \approx h \sum_{j=0}^i \omega_j v(x_j), \quad 0 \leq i \leq n,$$

such that  $\{\omega_i\}_{0 \leq i \leq n}$  is called the weights and verify:

$$\exists W > 0, \sup_{0 \leq i \leq n} |\omega_i| \leq W < +\infty, \quad n \geq 1.$$

We get the following discrete problem: For all  $i$  fixed, find  $u_i$  solution of the equation:

$$\mathcal{K}(x_i, x_i, u_i) + h \sum_{j=0}^i \omega_j \frac{\partial \mathcal{K}}{\partial x}(x_i, x_j, u_j) = f'(x_i), \quad (4.3)$$

which is equivalent for all  $i = 0, \dots, n$  and  $n \geq 1$ , to non-linear equation:

$$\begin{cases} \mathcal{K}(x_i, x_i, u_i) + h \omega_i \frac{\partial \mathcal{K}}{\partial x}(x_i, x_i, u_i) = S_i, \\ S_i = f'(x_i) - h \sum_{j=0}^{i-1} \omega_j \frac{\partial \mathcal{K}}{\partial x}(x_i, x_j, u_j). \end{cases} \quad (4.4)$$

Before studying the convergence of approximate solution, we need to show that the system (4.4) has a unique solution.

**Theorem 4.1.** *For  $h$  sufficiently small, the system (4.4) has a unique solution.*

*Proof.* For  $i \geq 1$  fixed, suppose that  $\{u_i\}_{0 \leq i \leq n}$  are known. Now, we present two sequences: the first is  $\{u_i^p\}_{p \in \mathbb{N}}$  and verifies the problem

$$\begin{cases} \mathcal{K}(x_i, x_i, u_i^{p+1}) = S_i - hw_i \frac{\partial \mathcal{K}}{\partial x}(x_i, x_i, u_i^p), & p \geq 1, \\ u_i^0 = u_{i-1}, \end{cases} \quad (4.5)$$

and the second is  $\{\phi_i^p\}_{p \in \mathbb{N}}$  and is defined by

$$\begin{cases} \phi_i^p = u_i^p - u_i^{p-1}, & p \geq 1, \\ \phi_i^0 = u_{i-1}. \end{cases} \quad (4.6)$$

It is clear that  $\sum_{q=0}^p \phi_i^q = u_i^p$ . Then, we prove that  $u_i^p$  converge to  $u_i$ . For all  $p \geq 1$ , by using  $(H_1)$ , 5

$$\begin{aligned} |\mathcal{K}(x_i, x_i, u_i^p) - \mathcal{K}(x_i, x_i, u_i^{p-1})| &\leq h|\omega_i| \left| \frac{\partial \mathcal{K}}{\partial x}(x_i, x_i, u_i^{p-1}) - \frac{\partial \mathcal{K}}{\partial x}(x_i, x_i, u_i^{p-2}) \right|, \\ &\leq LhW |u_i^{p-1} - u_i^{p-2}|. \end{aligned} \quad (4.7)$$

On the other hand, under the assumption  $(\mathcal{H})$  (7), we have

$$|\mathcal{K}(x_i, x_i, u_i^p) - \mathcal{K}(x_i, x_i, u_i^{p-1})| \geq \theta |u_i^p - u_i^{p-1}|. \quad (4.8)$$

According to equalities (4.7) and (4.8), we obtain

$$|u_i^p - u_i^{p-1}| \leq \frac{LhW}{\theta} |u_i^{p-1} - u_i^{p-2}|, \quad (4.9)$$

which gives

$$|\phi_i^p| \leq \frac{LhW}{\theta} |\phi_i^{p-1}|. \quad (4.10)$$

By induction, we can prove that

$$|\phi_i^p| \leq \left( \frac{LhW}{\theta} \right)^p |\phi_{i-1}^0|. \quad (4.11)$$

Finally, we get

$$\sum_{q=0}^p |\phi_i^q| = \sum_{q=0}^p \left( \frac{LhW}{\theta} \right)^q |u_i^0|.$$

Assuming that  $h$  is small enough so that  $\frac{LhW}{\theta} < 1$ , we can show that  $\sum_{q \geq 1} \left( \frac{LhW}{\theta} \right)^q$  is convergent.

Therefore,  $\sum_{q=0}^p \phi_i^q$  is convergent. So,  $\lim_{p \rightarrow +\infty} u_i^p = u_i$ . It remains to check whether this limit satisfies the required our equation.

Recalling (4.5)

$$\mathcal{K}(x_i, x_i, u_i^{p+1}) = S_i - hw_i \frac{\partial \mathcal{K}}{\partial x}(x_i, x_i, u_i^p), \quad p \geq 1.$$

Then,

$$\lim_{p \rightarrow +\infty} \mathcal{K}(x_i, x_i, u_i^{p+1}) = S_i - hw_i \lim_{p \rightarrow +\infty} \frac{\partial \mathcal{K}}{\partial x}(x_i, x_i, u_i^p),$$

since,  $\mathcal{K}$  and  $\frac{\partial \mathcal{K}}{\partial x} \in C^0([a, b]^2 \times \mathbb{R})$ . This leads

$$\mathcal{K}(x_i, x_i, u_i) = S_i - hw_i \frac{\partial \mathcal{K}}{\partial x}(x_i, x_i, u_i),$$

which gives the result.

It remains to show the existence of the solution. Let  $\{u_i\}_{0 \leq i \leq n}$  and  $\{v_i\}_{0 \leq i \leq n}$  be solutions of system (4.7) and (4.8) respectively. Then, for all  $i \geq 1$  fixed

$$|\mathcal{K}(x_i, x_i, u_i) - \mathcal{K}(x_i, x_i, v_i)| \geq \theta |u_i - v_i|. \quad (4.12)$$

On the other hand, we have

$$\begin{aligned} |\mathcal{K}(x_i, x_i, u_i) - \mathcal{K}(x_i, x_i, v_i)| &\leq hw \left| \frac{\partial \mathcal{K}}{\partial x}(x_i, x_i, u_i) - \frac{\partial \mathcal{K}}{\partial x}(x_i, x_i, v_i) \right|, \\ &\leq hWL |u_i - v_i|. \end{aligned} \quad (4.13)$$

Next, from (4.12) and (4.13) it follows that

$$|u_i - v_i| \leq \frac{hWL}{\theta} |u_i - v_i|. \quad (4.14)$$

Since  $\frac{hWL}{\theta} < 1$ , we get  $u_i = v_i$ . This proves the uniqueness of the solution of the system (4.4) and completes the proof of Theorem 4.1.  $\square$

*4.1.1. Approximate solution convergence of discrete problem.* First, we define the local consistency error as

$$\delta_n(h, x_i) = \int_0^{x_i} \mathcal{K}(x_i, t, u(t)) dt - h \sum_{j=0}^i \omega_j \frac{\partial \mathcal{K}}{\partial x}(x_i, x_j, u(x_j)).$$

We say the numerical method is consistent if

$$\lim_{h \rightarrow 0} \left( \max_{0 \leq i \leq n} |\delta_n(h, x_i)| \right) = 0.$$

Now, we establish the convergence of the approximate solution  $u_i$ .

**Theorem 4.2.** *If the approximation method is consistent, then*

$$\lim_{h \rightarrow 0} \left( \max_{0 \leq i \leq n} |u_i - u(x_i)| \right) = 0.$$

*Proof.* For  $n$  large enough and  $i = 0, 1, \dots, n$ , we have

$$\begin{aligned} |\mathcal{K}(x_i, x_i, u_i) - \mathcal{K}(x_i, x_i, u(x_i))| &\leq h \sum_{j=0}^i |\omega_j| \left| \frac{\partial \mathcal{K}}{\partial x}(x_i, x_j, u_j) - \frac{\partial \mathcal{K}}{\partial x}(x_i, x_j, u(x_j)) \right| + |\delta_n(h, x_i)|, \\ &\leq Lh |\omega_i| |u_i - u(x_i)| + Lh \sum_{j=0}^{i-1} |\omega_j| |u_j - u(x_j)| + |\delta_n(h, x_i)|. \end{aligned}$$

But, using  $H)_{1,7}$  we get

$$\theta |u_i - u(x_i)| \leq |\mathcal{K}(x_i, x_i, u_i) - \mathcal{K}(x_i, x_i, u(x_i))|.$$

Then, we obtain

$$\theta|u_i - u(x_i)| \leq Lh|\omega_i| |u_i - u(x_i)| + Lh \sum_{j=0}^{i-1} |\omega_j| |u_j - u(x_j)| + |\delta_n(h, x_i)|.$$

Thus

$$\theta|u_i - u(x_i)| \leq LhW |u_i - u(x_i)| + LhW \sum_{j=0}^{i-1} |u_j - u(x_j)| + |\delta_n(h, x_i)|,$$

so that

$$|u_i - u(x_i)| \leq \frac{LhW}{\theta - LhW} \sum_{j=0}^{i-1} |u_j - u(x_j)| + \frac{\max_{0 \leq i \leq n} |\delta_n(h, x_i)|}{\theta - LhW}.$$

Applying Gronwel's lemma [16], we obtain

$$|u_i - u(x_i)| \leq \frac{1}{\theta - LhW} \left(1 + \frac{LhW}{\theta - LhW}\right)^{i-1} \left(\max_{0 \leq i \leq n} |\delta_n(h, x_i)| + LhW|u_0 - u(x_0)|\right).$$

Moreover, we have

$$\left(1 + \frac{LhW}{\theta - LhW}\right)^{i-1} \leq \left(1 + \frac{LhW}{\theta - LhW}\right)^n,$$

and

$$\lim_{n \rightarrow \infty} \left(1 + \frac{LhW}{\theta - LhW}\right)^n < +\infty.$$

This means that there is a constant  $\mu > 0$  such that

$$\forall n \geq 1, \max_{0 \leq i \leq n} \frac{1}{\theta - LhW} \left(1 + \frac{LhW}{\theta - LhW}\right)^{i-1} \leq \mu.$$

This implies that

$$\max_{0 \leq i \leq n} |u_i - u(x_i)| \leq \mu \left(\max_{0 \leq i \leq n} |\delta_n(h, x_i)| + LhW|u_0 - u(x_0)|\right),$$

and when  $h$  converges to 0, we get convergence of  $n$  to  $\infty$ . The desired result follows.  $\square$

## 4.2. Linearization

We start by presenting the process of Newton's method. Let us recalling that en general, Newton's method is applied to solve many problems of applied mathematics which have the general form:

$$\begin{cases} \text{Find the solution } v \in \mathbb{R} \text{ of} \\ g(v) = 0, \end{cases} \quad (4.15)$$

where  $g$  is a non linear function. For  $i \geq 1$  fixed, we define the non-linear functional  $\psi_i$  as:

$$\begin{aligned} \psi_i : \mathbb{R} &\longrightarrow \mathbb{R} \\ v &\longmapsto \psi_i(v) = \mathcal{K}(x_i, x_i, v) + hw_i \frac{\partial \mathcal{K}}{\partial x}(x_i, x_i, v) - S_i. \end{aligned} \quad (4.16)$$

Assume that  $\frac{\partial \mathcal{K}}{\partial x}$  verifies the following hypotheses

$$(H_2) \quad \left\| \frac{\partial^2 \mathcal{K}}{\partial v \partial x} \right\| \in C^0([0, X]^2 \times \mathbb{R}, \mathbb{R})$$

Then, we get  $\psi_i \in C^2(\mathbb{R}, \mathbb{R})$ .

The system to solve has the following form:

$$\begin{cases} \text{Find the solution } u_i \in \mathbb{R} \text{ of} \\ \psi_i(u_i) = 0. \end{cases} \quad (4.17)$$

Applying Newton's method, we obtain the following system:

$$\begin{cases} u_i^0 \in \mathbb{R}, \\ u_i^{k+1} = u_i^k - \frac{\psi_i(u_i^k)}{\psi_i'(u_i^k)}. \end{cases} \quad (4.18)$$

**Theorem 4.3.** For  $i$  fixed, let  $\psi_i$  be a non-linear function defined by (4.16) of class  $C^2$ . Let  $R > 0$ ,  $J_{i,R}$  be closed ball such that  $J_{i,R} = [u_i - R, u_i + R] \subseteq I_i$  and  $C = \frac{\max |\psi_i''(v)|}{2 \min |\psi_i'(v)|}$ . If  $CR < 1$  and if the starting approximation  $u_i^0$  in  $J_{i,R}$ , then for all  $u_i^k$  in  $J_{i,R}$  and

$$|u_i^{k+1} - u_i| \leq C|u_i^k - u_i|^2,$$

where

$$C|u_i^k - u_i|^2 \leq \left( C|u_i^0 - u_i| \right)^{2k} \leq (CR)^{2k}.$$

*Proof.* See [37] and [38]. □

**Theorem 4.4.** Let  $u(x_i)$  be a solution of (4.2) and  $u_i^k$  is an approximate solution of the iterative system (4.18), then

$$\lim_{n \rightarrow +\infty} \lim_{k \rightarrow +\infty} \left( \max_{0 \leq i \leq n} |u(x_i) - u_i^k| \right) = 0.$$

*Proof.* For  $n$  and  $k$  large enough, we have

$$\max_{0 \leq i \leq n} |u(x_i) - u_i^k| \leq \max_{0 \leq i \leq n} |u(x_i) - u_i| + \max_{0 \leq i \leq n} |u_i - u_i^k|.$$

Theorem 4.2 gives

$$\max_{0 \leq i \leq n} |u_i - u(x_i)| \leq \mu \left( \max_{0 \leq i \leq n} |\delta_n(h, x_i)| + LhW|u_0 - u(x_0)| \right),$$

and from Theorem 4.3 we get

$$\max_{0 \leq i \leq n} |u_i - u_i^k| \leq \frac{(CR)^{2k}}{C}.$$

Then,

$$\max_{0 \leq i \leq n} |u(x_i) - u_i^k| \leq \mu \left( \max_{0 \leq i \leq n} |\delta_n(h, x_i)| + LhW|u_0 - u(x_0)| \right) + \frac{(CR)^{2k}}{C}.$$

So, when  $n \rightarrow +\infty$  and  $k \rightarrow +\infty$ , we get the desired result. □

### 5. Numerical Examples

In this section, we give two numerical examples to illustrate the efficiency and accuracy of proposed method. In the following examples, we calculate  $u_i$  according the scheme (4.18), and we define the discrete error as:

$$err_n = \max_{0 \leq i \leq n} |u(x_i) - u_i^k|.$$

Let us start with the equation:

$$\forall x \in [0, 1], \int_0^x \frac{1}{x+t+9+\exp(u(t))} dx = \frac{1}{2} \log \left( \frac{3x+10}{x+10} \right), \quad (5.1)$$

where the exact solution is  $u(x) = \log(x+1)$ . Table 1 shows the error magnitudes between the approximate and exact solutions. The results we present in the table were obtained using the MATLAB program.

| $n \backslash k$ | 10         | 20         | 30         | 50          |            |             |            |             |
|------------------|------------|------------|------------|-------------|------------|-------------|------------|-------------|
| 10               | 9.6047e-04 | 7.799171   | 9.6047e-04 | 10.994035   | 9.6047e-04 | 14.677863   | 9.6047e-04 | 25.032011   |
| 50               | 2.0446e-04 | 27.800183  | 2.0446e-04 | 69.370806   | 2.0446e-04 | 71.267143   | 2.0446e-04 | 273.740221  |
| 100              | 1.0290e-04 | 48.720499  | 1.0290e-04 | 91.716690   | 1.0290e-04 | 158.136187  | 1.0290e-04 | 5526.485901 |
| 500              | 7.9802e-04 | 225.026614 | 1.0705e-04 | 429.065173  | 1.8603e-05 | 2.0682e-05  | 7.3098e-06 | 3082.171232 |
| 1000             | 4.0433e-04 | 840.972380 | 5.3099e-05 | 1400.235219 | 8.2444e-06 | 1562.552823 | 1.8275e-06 | 4724.151322 |

Table 1: Errors of exact and approximate solutions.

In the table 1, in each row, we choose  $n$  and then change the value of  $k$ . We notice that when  $n$  is equal to 10 and 50, the error does not change. But, after having chosen a large  $n$ , the more the number of iterations  $k$  increases, the less the error becomes. So the approximate solution converges to the exact solution. In the next figure, we plot the approximate and exact solutions to observe the difference between them.




Figure 1: Approximate and exact solutions with  $h = 0.1$  of equation (5.1).

We give another equation

$$\forall x \in [0, 3], \int_0^x \frac{\exp(x)t^2 + 1}{\cos(t)^2 + 1 + u(t)^2} dt = \frac{x}{2} + \frac{x^3 \exp(x)}{6}, \tag{5.2}$$

and the exact solution  $u(x) = \sin(x)$ . Let introduce this table, which explain the error between the numerical and exact solution in all points  $x_i$ . What we are going to present in the table are calculated using MATLAB.

| n \ k | 10         | time(s)     | 20         | time(s)     | 30         | time(s)     | 50         | time(s)     |
|-------|------------|-------------|------------|-------------|------------|-------------|------------|-------------|
| 10    | 0.0260     | 11.175920   | 0.0176     | 22.756830   | 0.0176     | 26.460829   | 0.0176     | 37.429666   |
| 50    | 0.0068     | 38.840905   | 0.0013     | 87.966106   | 7.3009e-04 | 109.458969  | 7.3009e-04 | 158.506082  |
| 100   | 0.0037     | 59.144494   | 5.7459e-04 | 119.415460  | 1.8268e-04 | 174.320379  | 1.8268e-04 | 231.792491  |
| 500   | 7.9802e-04 | 249.483759  | 1.0705e-04 | 597.724081  | 1.8603e-05 | 5109.798293 | 7.3098e-06 | 1204.280340 |
| 1000  | 4.0433e-04 | 1003.764503 | 5.3099e-05 | 3112.587714 | 8.2444e-06 | 23388.40    | 1.8275e-06 | 2560.750899 |

Table 2: Errors of exact and approximate solutions.

In each row of the table 2, we will set  $n$  to a certain value and then change to  $k$  until we get the best possible error. From one row to the next, we increase the value of  $n$ , we notice that the error decreases each time. Our goal through the two tables 1 and 2 is to specify everywhere the number of divisions  $n$  of the interval  $[a, b]$  the larger. We obtain an error close to zero. Thus, we guarantee the convergence of the numerical solution to the exact solution.




Figure 2: Approximate and exact solution with  $h = 0.1$  of equation (5.2).

### 6. Conclusion

In this paper, we are interested in the analytical and numerical study of non-linear Volterra integral equation of the first kind. A sufficient condition has been introduced in order to prove the existence and uniqueness of the analytical solution. Then, for the numerical approach, we started with the discretization of our problem by applying the principle of Nyström’s method. Next, we obtained at each iteration  $i$  a non-linear equation to solve. For this reason, we applied the principle of Newton’s method to reformulate

the discrete nonlinear equation to a linear equation. We have presented two convergence theorems: the first one for the convergence of the discrete solution  $u_i$  to  $u(x_i)$  and the second one for the convergence of the iterative solution  $u_i^k$  to the discrete solution  $u_i$ . At the end, we have given two numerical examples to illustrate the efficiency of the proposed numerical method.

### References

1. D. N. Karamov, D. N. Sidorov, I. R. Muftahov, A. V. Zhukov, F. Liu, *Optimization of isolated power systems with renewables and storage batteries based on nonlinear Volterra models for the specially protected natural area of lake Baikal*. J. Phys.: Conf. Ser, 1847, 012037, (2021).
2. S., Noeiaghdam, S., Micula, J. J. Nieto, *A Novel Technique to Control the Accuracy of a Nonlinear Fractional Order Model of COVID-19: Application of the CESTAC Method and the CADNA Library*. Mathematics, 9, 1321, (2021).
3. E. Markova, I. Sidler, S. Solodusha, *Integral Models Based on Volterra Equations with Prehistory and Their Applications in Energy*. Mathematics , 9, 1127,(2021).
4. S. Solodusha, M. Bulatov, M. *Integral Equations Related to Volterra Series and Inverse Problems: Elements of Theory and Applications in Heat Power Engineering*. Mathematics, 9, 1905(2021).
5. W. Gao, P. Veerasha, C.Cattani, C. Baishya, H. M. Baskonus, *Modified Predictor–Corrector Method for the Numerical Solution of a Fractional-Order SIR Model with 2019-nCoV*. Fractal Fract, 6, 92, (2022).
6. T. Luo, Y. Xie, R. Wang, X. Yu, *An Analytic Solution to Precipitation Attenuation Expression with Spaceborne Synthetic Aperture Radar Based on Volterra Integral Equation*. Remote Sens, 14, 357(2022).
7. A. Jafarian, R. Rezaei, A. Golmankhaneh, *On Solving Fractional Higher-Order Equations via Artificial Neural Networks*. Iran. J. Sci. Technol. Trans. Sci. (2022).
8. T., Luo, Y., Xie, R., Wang, X., Yu,*An Analytic Solution to Precipitation Attenuation Expression with Spaceborne Synthetic Aperture Radar Based on Volterra Integral Equation* . Remote Sensing, 14(2), 357, (2022).
9. F. Brauer, *Constant Rate Harvesting of Populations Governed by Volterra Integral Equations*, J of mathematical analysis and applications, 56, IS-27 (1976).
10. M. C. De Bonis, C. Laurita, V. Sagaria, *A numerical method for linear Volterra integral equations on infinite intervals and its application to the resolution of metastatic tumor growth models*. Applied Numerical Mathematics, 172, 475-496,(2022).
11. V. Giorno, A. G. Nobile, *A numerical method for linear Volterra integral equations on infinite intervals and its application to the resolution of metastatic tumor growth models*. Applied Mathematics and Computation, 422, 126993,(2022).
12. P. Romàn-Romàn, S. Romàn-Romàn , *Juan José Serrano-Pérez and Francisco Torres-Rui:Using First-Passage Times to Analyze Tumor Growth Delay*. Mathematics, 9, 642, (2021).
13. I.S. Halpea, G.L. Parajdi, R. Precup, *On the controllability of a system modeling cell dynamics related to Leukemia*. Symmetry, 13, 1867, (2021).
14. M. Lienert, R. Tumulka, *A new class of Volterra-type integral equations from relativistic quantum physics*. J. Integral Equations Applications, 31, 4, 535-569, (2019).
15. Wazwaz, A.M., *Linear and Nonlinear Integral Equations: Methods and Applications*, Springer Berlin Heidelberg, (2011).
16. Linz P., *Analytical and Numerical Methods for Volterra Equations*, Society for Industrial Mathematics, (1987).
17. Sh. Lal, S. Kumar, *Approximation of functions by Bernoulli wavelet and its applications in solution of Volterra integral equation of second kind*. Arab. J. Math. (2021).
18. I. Dimo, S. Maire, V. Todorov, *An unbiased Monte Carlo method to solve linear Volterra equations of the second kind, Neural Computing and Applications*,(2021).
19. R. Katani, *A Numerical Method for Proportional Delay Volterra Integral Equations*. Int. J. Appl. Comput. Math. 7-17, (2021).
20. Kabanikhin, S. I.,*Inverse and Ill-posed Problems: Theory and Applications*. De Gruyter, (2011).
21. Z. Masouri, *A Regularization-direct Method to Numerically Solve First Kind Fredholm Integral Equation*. Kyungpook Math. J. 60, 869-88, (2020).
22. T. T. Karakeev, T. M. Imanaliev,*Regularization of Volterra Linear Integral Equations of the First Kind with the Smooth Data*. Lobachevskii Journal of Mathematics. 41,1, 34-45, (2020).
23. Atkinson, K., Han, W., *Theoretical Numerical Analysis: A Functional Analysis Framework*. Springer-Verlag, New York, (2001).
24. A. Deepa, A. Kumarb, S. Abbasc, M. Rabbanid,*Solvability and numerical method for non-linear Volterra integral equations by using Petryshyn's fixed point theorem*. Int. J. Nonlinear Anal. Appl. 13,1, 1-28,(2022).

25. A. Fawze, B. Juma'a, W. Al Hayani, *Homotopy perturbation technique to solve nonlinear systems of Volterra integral equations of 1st kind*. Journal of Soft Computing and Artificial Intelligence JSCAI, 2, 19-26, (2021).
26. R. Dehbozorgi, K. Maleknejad, K. *Direct Operational Vector Scheme for First-Kind Nonlinear Volterra Integral Equations and Its Convergence Analysis*. Mediterr. J. Math. 18-31, (2021).
27. K. Nedaiasl, R. Dehbozorgi, K. Maleknejad, *Hp-version collocation method for a class of nonlinear Volterra integral equations of the first kind*. Applied Numerical Mathematics, 150, 452-477,(2019).
28. H. Guebbai, S. Lemita, S. Segni, W. Merchela, *Difference derivative for an integro-differential nonlinear volterra equation*. Vestnik Udmurtskogo Universiteta Matematika Mekhanika Komp'uternye Nauki, 30(2), 176-188,(2020).
29. S. Segni, M. Ghiat, H. Guebbai, *New approximation method for Volterra nonlinear integro-differential aquation*. Asian-European Journal of Mathematics, 12(1), 1950016, (2019).
30. M. Ghiat, H. Guebbai, M. Kurulay, S. Segni, *On the weakly singular integro-differential nonlinear Volterra equation depending in acceleration term*. Computational and Applied Mathematics, 39 (2), 206,(2020).
31. S. Touati, S. Lemita, M. Ghiat, M. Z. Aissaoui M.Z. *Solving a non-linear Volterra-Fredholm integro-differentail equation with weakly singular kernels*. Fasciculi Mathematics, 62, 155-168, (2019).
32. M. Ghiat, H. Guebbai, *Analytical and numerical study for an integro-differential nonlinear volterra equation with weakly singular kernel*. Computational and Applied Mathematics, 37(4), 4661-4974, (2018).
33. M. Ch. Bounaya, S. Lemita, M. Ghiat, H. Guebbai, M. Z. Aissaoui, *On a nonlinear integro-differential equation of Fredholm type*. Int. J. Comput. Sci. Math. 13(2), 194, (2021).
34. Grammont, L., Ahues, M., D'Almeida, F.D.: For nonlinear infinite dimensional equations, which to begin with: linearization or discretization? Dedicated to the memory of Alain Largillier. Journal of Integral Equations and Applications Volume. 26(2) (2014). <https://10.1216/JIE-2014-26-3-413>
35. L. Grammont, *Nonlinear Integral Equations of the Second Kind: A New Version of Nyström Method*. Numerical Functional Analysis and Optimization. 34(5), (2013).
36. A. Khellaf, W. Merchela, S. Benarab, *New numerical process solving nonlinear infinite-dimensional equations*. Computational and Applied Mathematics. 39(93),16 (2020).
37. C. Villani, *Le fabuleux destin de la méthode de Newton*. Inauguration de la Fondation Hadamard, 17, (2011).
38. I.K Argyros, *Convergence and Applications of Newton-type Iterations*. Springer Science+Business, Media, LLC, New York, (2008).
39. Lions, J. L., *Exact Controllability, Stabilizability, and Perturbations for Distributed Systems*, Siam Rev. 30, 1-68, (1988).
40. C. M. Dafermos, C. M., *An abstract Volterra equation with application to linear viscoelasticity*. J. Differential Equations 7, 554-589, (1970).

*Boutheina Tair,*  
*Department of Mathematics,*  
*08 Mai 1945 Guelma university ,*  
*Algeria.*  
*E-mail address: tairboutheina2@gmail.com*

and

*Mourad Ghiat,*  
*Department of Mathematics,*  
*08 Mai 1945 Guelma university ,*  
*Algeria.*  
*E-mail address: mourad.ghi24@gmail.com*

and

*Hamza Guebbai,*  
*Department of Mathematics,*  
*08 Mai 1945 Guelma university ,*  
*Algeria.*  
*E-mail address: guebaihamza@yahoo.fr*

and

*Mohamed Zine Aissaoui,*  
*Department of Mathematics,*  
*08 Mai 1945 Guelma university ,*  
*Algeria.*  
*E-mail address: aissaoui.zine@univ-guelma.dz*