



Nörlund Difference Sequence Spaces in 2-Normed Spaces Using Orlicz Function

Sunil K. Sharma, Ravi Kumar, Ayhan Esi and Ajay K. Sharma

ABSTRACT: In this paper, we introduce the concept of paranormed Nörlund-difference sequence spaces with Orlicz function over 2-normed spaces denoted by $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ with Δ -transform are in c_0, c and ℓ_∞ . Additionally, we investigate various topological characteristics of these spaces and analyze various inclusion relationships between these spaces. Furthermore, we establish and construct the α –, β – and γ –duals of these spaces, along with establishing the Schauder basis for $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$.

Keywords: Difference sequence space, Nörlund matrix, paranormed sequence space, Orlicz function, α –, β – and γ – duals.

Contents

| | |
|---|----------|
| 1 Introduction and Preliminaries | 1 |
| 2 The Nörlund-Difference Sequence Spaces with Orlicz Function | 4 |
| 3 The α–, β– and γ– Duals of the Spaces | 8 |

1. Introduction and Preliminaries

In the paper, we will use ω to represent the class of all sequences formed by real or complex numbers, while ℓ_∞, c and c_0 will respectively denote the set of all bounded, convergent and null sequences. Additionally, we will use cs, bs, ℓ_1 and ℓ_p to represent the sequence spaces of all convergent, bounded, absolutely and p -absolutely convergent series, respectively. Gähler [4] introduced the concept of a 2-normed space as an innovative linear extension of a normed linear space. Subsequently, this concept has been further investigated by numerous researchers, as highlighted in the literature (refer to [3, 5]).

Suppose Y be a real vector space having dimension d , where $2 \leq d < \infty$. Consider the function $\|\cdot, \cdot\| : Y \times Y \rightarrow \mathbb{R}$, defined on Y , which acts as a 2-norm and satisfying the following properties:

1. $\|x, y\| = 0$ if and only if x and y are linearly dependent in Y ;
2. $\|x, y\| = \|y, x\|$;
3. $\|\alpha x, y\| = |\alpha| \|x, y\|, \alpha \in \mathbb{R}$;
4. $\|x, y + z\| \leq \|x, y\| + \|x, z\|$ for all $x, y, z \in Y$.

The pair $(Y, \|\cdot, \cdot\|)$ is known as 2-normed space, as in [5]. For instance, consider $Y = \mathbb{R}^2$ equipped with a 2-norm defined by $\|x, y\|$, where x and y are vectors in \mathbb{R}^2 , representing the area of the parallelogram spanned by vectors x and y . This 2-norm is expressed as follows:

$$\|y_1, y_2\|_E = \text{abs} \begin{vmatrix} y_{11} & y_{12} \\ y_{21} & y_{22} \end{vmatrix}$$

Then, $(Y, \|\cdot, \cdot\|)$ is a 2-normed space. In a 2-normed space $(Y, \|\cdot, \cdot\|)$, a sequence (χ_k) is said to be converge to $L \in Y$ if

$$\lim_{k \rightarrow \infty} \|\chi_k - L, z\| = 0$$

for every $z \in Y$.

In a 2-normed space $(Y, \|\cdot, \cdot\|)$, a sequence (χ_k) is called Cauchy if

$$\lim_{\substack{k \rightarrow \infty \\ p \rightarrow \infty}} \|\chi_k - \chi_p, z\| = 0$$

for every $z \in Y$. When every Cauchy sequence in Y has limit $L \in Y$, then Y is termed as complete with respect to the 2-norm. Moreover, any 2-normed space that is complete is termed as a 2-Banach space.

Definition 1.1 *Given that Y be a linear metric space, then $P : Y \rightarrow \mathbb{R}$ is a paranormed space if*

1. $P(y) \geq 0$
2. $P(-y) = P(y)$
3. $P(y + z) \leq P(y) + P(z)$ for all $y, z \in Y$
4. if α_k be sequence of scalars such that $\alpha_k \rightarrow \alpha$ as $k \rightarrow \infty$ and y_k be the sequence of vectors such that $P(y_k - y) \rightarrow 0$ as $k \rightarrow \infty$, then $P(\alpha_k y_k - \alpha y) \rightarrow 0$ as $k \rightarrow \infty$.

A total paranorm, denoted as P , is a paranorm such that $P(y) = 0$ implies $y = 0$. A total paranormed space is defined as a space Y equipped with a total paranorm P , i.e., (Y, P) . The metric of every linear metric space is uniquely determined by some total paranorm see [25]. For additional details regarding sequence spaces, we refer to [1,15,16,17,20,21,22] and references therein.

Consider φ_1 and φ_2 as arbitrary sequence spaces. We define matrix $A = (a_{nk})$ as any infinite matrix, where $a_{nk} \in \mathbb{R}$ and $n, k \in \mathbb{N}$. Thus the matrix mapping from φ_1 into φ_2 defines by A and is represented by $A : \varphi_1 \rightarrow \varphi_2$, if for any $\xi = (\xi_n) \in \varphi_1$, the resulting sequence $A\xi = \{(A\xi)_n\} \in \varphi_2$, where

$$(A\xi)_n = \sum_k a_{nk} \xi_k, \text{ for } n \in \mathbb{N}. \quad (1.1)$$

By (φ_1, φ_2) , we represent the collection of all matrices A , for each $n \in \mathbb{N}$ and every $\xi \in \varphi_1$ the series defined in equation (1.1) converges. We define that a sequence ξ is said to be A -summable to u if the sequence of matrix-vector products $A\xi$ converges to u . This convergence of Ax to u is termed as the A -limit of ξ .

Assuming (p_n) be a bounded sequence where each term is strictly positive, with $\sup p_n = Q$ and $M = \max\{1, Q\}$. Maddox in ([10,11,12]), Nakano [18] and Simons [23] defined the linear spaces $\ell_\infty, c(p)$ and $c_0(p)$ in the following manner:

$$\begin{aligned} \ell_\infty &= \{\xi = (\xi_n) \in \omega : \sup_{n \in \mathbb{N}} |\xi_n|^{p_n} < \infty\}, \\ c(p) &= \{\xi = (\xi_n) \in \omega : \lim_{n \rightarrow \infty} |\xi_n - L|^{p_n} = 0\} \text{ for some } L > 0, \\ c_0(p) &= \{\xi = (\xi_n) \in \omega : \lim_{n \rightarrow \infty} |\xi_n|^{p_n} = 0\}. \end{aligned}$$

In paranormed, they are complete sequence spaces

$$g(\xi) = \sup_{n \in \mathbb{N}} |\xi_n|^{\frac{p_n}{M}}, \text{ iff } \inf_{k \in \mathbb{N}} p_k > 0 \quad (1.2)$$

The set $S(u, v)$ is defined for the sequence space u and v as

$$S(u, v) = \{y \in \omega : \xi y \in v, \forall \xi \in u\} \quad (1.3)$$

The duals $\alpha-, \beta-, \gamma-$ of k , represented by $k^\alpha, k^\beta, k^\gamma$, respectively and defined as follows: $k^\alpha = S(k, \ell_1)$, $k^\beta = S(k, cs)$, $k^\gamma = S(k, bs)$.

Definition 1.2 The sequence x_n is known as Schauder basis of the paranormed sequence space (η, f) , if $\xi \in \eta$, there exist a sequence (α_n) such that

$$\lim_{k \rightarrow \infty} f\left(\xi - \sum_{n=0}^k \alpha_n x_n\right) = 0.$$

The Nörlund means, introduced by Peyerimhoff [19] and Mears [14], are a concept in mathematics that involve manipulating sequences. Given a sequence $T_n = \sum_{k=0}^n t_k$ for all $n \in \mathbb{N}$, where $t_k \geq 0$ and $t_0 > 0$, the representation of Nörlund means for $t = (t_k)$ as a matrix is $\mathfrak{N}^t = (a_{nk}^t)$, where

$$a_{nk}^t = \begin{cases} \frac{t_{n-k}}{T_n}, & 0 \leq k \leq n \\ 0, & k > n \end{cases} \text{ for all } n \in \mathbb{N}.$$

Singh and Malik [24] introduced paranormed Norlund \mathfrak{N}^t -difference sequence spaces. The Nörlund matrix \mathfrak{N}^t transforms to the matrix \mathcal{C}_1 when $t = e = (1, 1, 1, \dots)$, which corresponds to the matrix of arithmetic means. Now, let us consider $t_n = A_n^{r-1}$, where $r > -1$ and A_n^{r-1} denotes the n -th Abel mean of order $r - 1$. Then, the method \mathfrak{N}^t gives the Cesàro method \mathcal{C}_r for $r > -1$. In other words, the matrix \mathfrak{N}^t for $t_n = A_n^{r-1}$ can be represented as follows:

$$A_n^r = \begin{cases} \frac{(r+1)(r+2)\cdots(r+n)}{n!}, & n \in \mathbb{N} \\ 1, & n = 0 \end{cases} \text{ for } n \in \mathbb{N}.$$

For $n \in \mathbb{N}$, D_n denotes the determinant of square matrix, where $t_0 = D_0 = 1$ given, as

$$D_n = \begin{vmatrix} t_1 & 1 & 0 & 0 & \dots & 0 \\ t_2 & t_1 & 1 & 0 & \dots & 0 \\ t_3 & t_2 & t_1 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ t_{n-1} & t_{n-2} & t_{n-3} & t_{n-4} & \dots & 1 \\ t_n & t_{n-1} & t_{n-2} & t_{n-3} & \dots & t_1 \end{vmatrix}$$

If $V^t = (r_{nk}^t)$ is the inverse of $\mathfrak{N}^t = (a_{nk}^t)$, [14], then

$$r_{nk}^t = \begin{cases} (-1)^{n-k} D_{n-k} T_k, & 0 \leq k \leq n, \\ 0, & k > n, \end{cases} \text{ for all } n, k \in \mathbb{N}.$$

Also for all $k \in \mathbb{N}$, we have

$$D_k = \sum_{i=0}^{k-1} (-1)^{i-1} t_i D_{k-i} + (-1)^{k-1} t_k.$$

An Orlicz function \mathfrak{M} is a function that is continuous, convex and nondecreasing with $\mathfrak{M}(0) = 0$, $\mathfrak{M}(u) > 0$ for $u > 0$ and $\mathfrak{M}(u) \rightarrow \infty$ as $u \rightarrow \infty$. This Orlicz function is used to define a sequence space, which was introduced by Lindenstrauss and Tzafriri [9]. The concept involves using the Orlicz function to characterize the properties of the sequence space. In other words, the sequence space $X_{\mathfrak{M}}$ associated with the Orlicz function \mathfrak{M} consists of all sequences $\xi = (\xi_n)$ such that

$$\sum_{n=1}^{\infty} \mathfrak{M}(|\xi_n|) < \infty.$$

Alternatively, we can define $X_{\mathfrak{M}}$ as the set of all sequences such that the norm

$$\|\xi\|_{\mathfrak{M}} = \inf \left\{ \lambda > 0 : \sum_{n=1}^{\infty} \mathfrak{M}\left(\frac{|\xi_n|}{\lambda}\right) < \infty \right\}$$

is finite, where $\|\cdot, \cdot\|$ denotes the Orlicz norm induced by the Orlicz function \mathfrak{M} .

Furthermore, within every Orlicz sequence space $\ell_{\mathfrak{M}}$, there exists a subspace which is isomorphic to ℓ_p for $p \geq 1$ [9]. The Δ_2 -condition states that $\mathfrak{M}(Ku) \leq K\mathfrak{M}(u)$ for all $u \geq 0$ and $0 < K < 1$. This condition verifies the boundedness of the Orlicz function.

Furthermore, the Orlicz function $\mathfrak{M}(u)$ is expressed in terms of an integral as:

$$\mathfrak{M}(u) = \int_0^u \eta(y) dy.$$

Here, $\eta(y)$ denotes the kernel of \mathfrak{M} . The kernel η exhibits right differentiability for $y \geq 0$ with $\eta(0) = 0$, $\eta(y) > 0$, η is nondecreasing and $\eta(y) \rightarrow \infty$ as $y \rightarrow \infty$. This representation provides another perspective on the Orlicz function and its properties.

Kızılmaz [7] introduced several distinct difference spaces, namely $\ell_{\infty}(\Delta)$, $c(\Delta)$ and $c_0(\Delta)$ which were extensively investigated by Et and Colak [2]. Let m (where $m \geq 0$) be an integer. Given any sequence space λ , we define $\lambda(\Delta^m) = \{t = (t_n) \in \omega : (\Delta^m \xi_n) \in \lambda\}$ for $\lambda = c_0, c$ and ℓ_{∞} , where $\Delta^m \xi = (\Delta^m \xi_n) = (\Delta^{m-1} \xi_n - \Delta^{m-1} \xi_{n+1})$. This can also be expressed as

$$\Delta^m \xi_n = \sum_{u=0}^m (-1)^u \binom{m}{u} \xi_{n+u}.$$

2. The Nörlund-Difference Sequence Spaces with Orlicz Function $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_{\infty}}$ Over 2-Normed Spaces

In this section, we introduce new paranormed Nörlund-difference sequence spaces equipped with Orlicz function defined over 2-normed spaces denoted as $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_{\infty}}$. We construct paranormed structures over these spaces and demonstrate their linear isomorphism.

The Nörlund sequence spaces $\mathfrak{N}^t(p)$ were introduced by Yeşilkayagil and Başar [26] as

$$\mathfrak{N}^t(p) = \{\xi = (\xi_n) \in \omega : \sum_n \left\| \frac{1}{T_k} \sum_{i=0}^n t_{k-i} x_i \right\|^{p_n} < \infty\}$$

where $0 < p_n \leq Q < \infty$.

Now, the Δ -Nörlund-difference Orlicz sequence spaces over 2-normed spaces are defined as follows

$$\begin{aligned} \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0} &= \left\{ \xi \in \omega : \lim_{n \rightarrow \infty} \left[\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i} \Delta \xi_i}{\rho}, z \right\| \right) \right]^{p_n} = 0 \right\} \\ \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c &= \left\{ \xi \in \omega : \lim_{n \rightarrow \infty} \frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i} \Delta \xi_i - L}{\rho}, z \right\| \right)^{p_n} = 0 \right\} \\ \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_{\infty}} &= \left\{ \xi \in \omega : \sup_n \left[\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i} \Delta \xi_i}{\rho}, z \right\| \right) \right]^{p_n} < \infty \right\}. \end{aligned}$$

where $\mathfrak{M} = (\mathfrak{M}_k)$ be the sequence of Orlicz functions and $p = p_n$ be the sequence of bounded positive real numbers.

Let $\chi = (\chi_n)$ be defined as the $\mathfrak{N}^t(\Delta)$ -transform of the sequence $\xi = (\Delta \xi_n)$, then

$$\chi = (\chi_n) = \frac{1}{T_n} \sum_{i=0}^n t_{n-i} \Delta \xi_i \quad \forall n \in \mathbb{N}.$$

Theorem 2.1 For $0 \leq p_n \leq Q < \infty$ and $M = \max\{1, Q\}$, the spaces $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_{\infty}}$ are the complete linear metric spaces paranormed by

$$g(\xi) = \sup_n \left[\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i} \Delta \xi_i}{\rho}, z \right\| \right) \right]^{\frac{p_n}{M}}.$$

Proof: The result has been proven for $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$. Since every bounded sequence has a finite supremum, the proof for the other spaces follows the similar method. Suppose that $\xi, \chi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, then

$$\begin{aligned} \sup_n \left[\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i} \Delta (\xi_i + \chi_i)}{\rho}, z \right\| \right) \right]^{\frac{p_n}{M}} \\ \leq \sup_n \left[\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i} \Delta \xi_i}{\rho}, z \right\| \right) \right]^{\frac{p_n}{M}} \\ + \sup_n \left[\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i} \Delta \chi_i}{\rho}, z \right\| \right) \right]^{\frac{p_n}{M}} \end{aligned}$$

and for any $\alpha \in \mathbb{R}$, we have

$$\alpha^{p_k} \leq \{\max(1, \alpha^M)\}. \quad (2.1)$$

Given that $\mathfrak{M}(\theta) = 0$, it follows that $g(\theta) = 0$ and $g_1(\xi) = g_1(-\xi) \forall \xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$. Therefore, inequalities (2.1) imply sub-additivity of the function g and $g(\alpha\xi) \leq \max(1, \alpha^M)g(\xi)$. Moreover, suppose $(\xi^{(n)}) \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, then $g(\xi^{(n)} - \xi) \rightarrow 0$ and let (α_n) is the sequence of scalars converging to α . Thus,

$$\begin{aligned} g(\alpha_n \xi^{(n)} - \alpha \xi) &= \sup_n \left[\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i} \Delta (\alpha_n \xi_i^{(n)} - \alpha \xi_i)}{\rho}, z \right\| \right) \right]^{\frac{p_n}{M}} \\ &\leq \alpha_n - \alpha^{\frac{p_n}{M}} g(\xi^{(n)}) + \alpha^{\frac{p_n}{M}} g(\xi^{(n)} - \xi) \\ &\rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Hence, g is paranorm.

Given a Cauchy sequence $\{\xi^i\} \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, where $\xi^i = \{\xi_0^{(i)}, \xi_1^{(i)}, \xi_2^{(i)}, \dots\}$. For $\epsilon > 0$ be given, there exists $n_0(\epsilon)$ such that $g(\xi^i - \xi^j) < \epsilon$ for $i, j \geq n_0(\epsilon)$. Then, for any $k \in \mathbb{N}$, we get

$$\begin{aligned} (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^i)_k - (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^j)_k^{\frac{p_n}{M}} \\ \leq \sup_n (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^i)_k - (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^j)_k^{\frac{p_n}{M}} \\ < \frac{\epsilon}{2}, \forall i, j > n_0(\epsilon) \end{aligned} \quad (2.2)$$

it follows that the sequence of real numbers forms a Cauchy sequence

$\{(\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^0)_k, (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^1)_k, \dots\}$, for $k \in \mathbb{N}$. Therefore,

$\{(\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^i)_k \rightarrow (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_k$ as $i \rightarrow \infty$.

For the sequences $(\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_0, (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_1, \dots$ with infinitely many limits, there exist a sequence

$\{(\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_0, (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_1, (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_2, \dots\}$. By applying inequality (2.2) as $j \rightarrow \infty$, we get

$$(\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^i)_k - \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_k < \frac{\epsilon}{2}, i \geq n_0(\epsilon).$$

Since $\xi^i = (\xi_n^i) \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ for each $i \in \mathbb{N}$, there exists $n_0(\epsilon) \in \mathbb{N}$ such that

$$(\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^i)_k^{\frac{p_n}{M}} < \frac{\epsilon}{2}, \text{ for every } i \geq n_0(\epsilon) \text{ and } k \in \mathbb{N}. \quad (2.3)$$

For fixed $i \geq n_0(\epsilon)$ and from (2.3), we get

$$\begin{aligned} (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_k^{\frac{p_n}{M}} &\leq (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^i)_k - (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_k^{\frac{p_n}{M}} \\ &\quad + (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi^i)_k^{\frac{p_n}{M}} < \epsilon \end{aligned}$$

for every $i \geq n_0(\epsilon)$. Hence, $\xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$. \square

Remark 2.1 The space $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ does not satisfy the absolute property, it follows that $\|g(\xi)\| \neq \|g(|\xi|)\|$. This indicates that $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ is non-absolute type.

Theorem 2.2 For $0 < p_n \leq Q < \infty$, the sequence spaces $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$, which are of non-absolute type, can be either paranorm or norm isomorphic to the spaces $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_c$, $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ and $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ respectively.

Proof: Consider the linear transformation $T : \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0} \rightarrow \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ defined as $T\xi = (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0})\xi$. If $T\xi = 0$ only when $\xi = 0$, then T is injective.

Let $\chi \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ and $\xi = (\xi_n) = (\Delta\xi_n)$ be the sequence which is defined as

$$\xi = (\xi_n) = \sum_{j=0}^n (-1)^{n-j} D_{n-j} T_j \Delta \chi_j, \text{ for all } n \in \mathbb{N}. \text{ Then, we have}$$

$$\begin{aligned} g(\xi) &= \sup_n \left[\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{n-i} \Delta \xi_i}{\rho}, z \right\| \right) \right]^{\frac{p_n}{M}} \\ &= \sup_n \left[\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{n-i} (\sum_{j=0}^n (-1)^{n-j} D_{n-j} T_j \Delta \chi_j)}{\rho}, z \right\| \right) \right]^{\frac{p_n}{M}} \\ &= \sup_n |\chi_n|^{\frac{p_n}{M}} < \infty. \end{aligned}$$

Thus, $\xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, implying that T is surjective and its preservation under paranorm. Therefore $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ and $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ are linearly isomorphic. In the same manner, we can establish that $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ are isomorphic to $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ respectively. \square

Theorem 2.3 Consider the sequence $\Psi^{(n)}(t) = \{\Psi_k^{(n)}(t)\}$ which is defined as

$$\Psi_k^{(n)}(t) = \begin{cases} (-1)^{(k-n)D_{k-n}T_n}, & 0 \leq n \leq k \\ 0, & n > k. \end{cases}$$

Then

1. For $0 < p_n \leq Q < \infty$, the sequence $\{\Psi^{(n)}(t)\}_{n \in \mathbb{N}}$ forms a basis for $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ and $\xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ can be uniquely expressed as $\xi = \sum_n \alpha_n(t) \Psi^{(n)}(t)$, where $\alpha_n(t) = (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_n$, for all $n \in \mathbb{N}$.
2. The set $\{e, \Psi^{(n)}(t)\}$ forms a basis of $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and every $\xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ can be uniquely expressed as $\xi = \mu e + \sum_n [\alpha_n(t) - \mu] \Psi^{(n)}(t)$, where $\mu = \lim_{n \rightarrow \infty} (\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)\xi)_n$.

Proof: (1) Clearly, $\{\Psi^{(n)}(t)\} \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, also

$$\mathfrak{N}^t \Psi^{(n)}(t) = e^{(n)} \in \ell(c_0, \Delta), \forall n \in \mathbb{N}, 0 < p_n \leq Q < \infty. \quad (2.4)$$

For given $\xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ and for every non-negative integer m , we have

$$\xi^{[m]} = \sum_{n=0}^m \alpha_n(t) \Psi^{(n)}(t). \quad (2.5)$$

Thus by using \mathfrak{N}^t and above equations, we have

$$\mathfrak{N}^t \xi^{[m]} = \sum_{n=0}^m \alpha_n(t) \mathfrak{N}^t \Psi^{(n)}(t) = \sum_{n=0}^m (\mathfrak{N}^t \xi)_n e^{(n)}.$$

Now, for all $i, m \in \mathbb{N}$, we have

$$\{\mathfrak{N}^t(\xi - \xi^{[m]})\}_i = \begin{cases} 0, & 0 \leq i \leq m \\ (\mathfrak{N}^t \xi)_i, & i > m \end{cases}.$$

For given $\epsilon > 0$, there exists a number $m_0 \in \mathbb{N}$ such that

$$\left[\sum_{i=m}^{\infty} (\mathfrak{N}^t \xi)_i^{p_n} \right]^{\frac{1}{M}} < \frac{\epsilon}{2}, \text{ for all } m \geq m_0.$$

Thus,

$$\begin{aligned} g[\mathfrak{N}^t(\xi - \xi^{[m]})] &= \left[\sum_{i=m}^{\infty} (\mathfrak{N}^t \xi)_i^{p_n} \right]^{\frac{1}{M}} \\ &\leq \left[\sum_{i=m_0}^{\infty} (\mathfrak{N}^t \xi)_i^{p_n} \right]^{\frac{1}{M}} \\ &< \epsilon, \text{ for all } m \geq m_0. \end{aligned}$$

Assume that $\xi = \sum_n \mu_n(t) \Psi^{(n)}(t)$. Then

$$\begin{aligned} (\mathfrak{N}^t \xi)_k &= \sum_n \mu_n(t) \{\mathfrak{N}^t \Psi^{(n)}(t)\}_k \\ &= \sum_n \mu_n(t) \epsilon_k^{(n)} = \mu_k(t), \text{ for all } k \in \mathbb{N}, \end{aligned}$$

which contradicts the assumption.

(2) Since $\{\Psi^{(n)}(t)\} \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ and $e \in c$, the inclusion $\{e, \Psi^{(n)}(t)\} \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ is trivial. For $\xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$, there exist unique μ satisfying equation (2.5). Thus, $\ell \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ when $\ell = \xi - \mu e$. Therefore, from part (1), the representation is unique. \square

Theorem 2.4 *The inclusion $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0} \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ strictly holds.*

Proof: Let $\chi \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$. Consequently $\mathfrak{N}^t \Delta \xi \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$. As $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0} \subset \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_c$, it follows that $\mathfrak{N}^t \Delta \xi \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_c$, thus implying that $\xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$. Thus the inclusion $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0} \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ is established.

Additionally, given that for every $\xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ we have $\mathfrak{N}^t \Delta \xi \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and the inclusion $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0} \subset \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_c$ is strict for some $\mathfrak{N}^t \Delta \xi \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_c$, it follows that $\xi \notin \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$.

On the similar lines, it can be readily established that the inclusion $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ is strict. \square

Theorem 2.5 *For $1 \leq p_n \leq p_{n+1}$, the inclusion $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_c \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$, $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0} \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ and $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty} \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ holds for all $n \in \mathbb{N}$.*

Proof: For $p = e$, inclusions are obvious (see [13]). Now we examining the case for $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty} \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$. Given $\xi \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$, we have $\Delta \xi^p \in \ell_\infty$, where $\xi^p = (\xi_k^{p_k})_{k=0}^\infty$. Choose a constant $m_0 \in \mathbb{N}$ such that $\Delta \xi_k^{p_k} < 1$ holds for all $k > m_0$. Thus for $n > m_0$, we have

$$\Delta \xi_k^{p_n} = (\Delta \xi_k^{p_k})^{\frac{p_n}{p_k}} \leq \Delta \xi_k^{p_k}, \quad m_0 \leq k \leq n. \quad (2.6)$$

Given that $p = (p_n)$ is bounded and $p_k \leq p_n$ for $k \leq n$ and $n \in \mathbb{N}$ and , it follows that for $\mathcal{K} > 0$, we get

$$\sup_n \left[\frac{1}{T_n} \sum_{i=0}^{m_0-1} \mathfrak{M}_k \left(\left\| \frac{t_{k-i} \Delta \xi_i}{\rho}, z \right\| \right) \right]^{p_n} \leq \mathcal{K} \sup_n \left(\frac{1}{T_n} \sum_{i=0}^{m_0-1} \mathfrak{M}_k \left(\left\| \frac{t_{k-i}}{\rho}, z \right\| \right) \right)^{p_n} |\Delta \xi_i|^{p_k} \quad (2.7)$$

Therefore, by applying Hölder inequality and using equation (2.6) and (2.7), we have $|\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}|^{p_n}$

$$\begin{aligned} &\leq \left[\sup_n \left(\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i}}{\rho}, z \right\| \right) \right) |\Delta \xi_i| \right]^{p_n} \\ &\leq \left[\sup_n \frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i}}{\rho}, z \right\| \right) |\Delta \xi_i|^{p_n} \right] \left[\sup_n \left(\frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i}}{\rho}, z \right\| \right) \right) \right]^{p_n-1} \\ &= \sup_n \frac{1}{T_n} \sum_{i=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i}}{\rho}, z \right\| \right) |\Delta \xi_i|^{p_n} \\ &= \sup_n \frac{1}{T_n} \left[\sum_{i=0}^{n_0-1} \mathfrak{M}_k \left(\left\| \frac{t_{k-i}}{\rho}, z \right\| \right) |\Delta \xi_i|^{p_n} + \sum_{k=n_0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i}}{\rho}, z \right\| \right) |\Delta \xi_i|^{p_n} \right] \\ &\leq \frac{\mathcal{K} + 1}{T_n} \sum_{k=0}^n \mathfrak{M}_k \left(\left\| \frac{t_{k-i}}{\rho}, z \right\| \right) |\Delta \xi_i|^{p_n} \\ &= (\mathcal{K} + 1) (\mathfrak{N}^t(\Delta \xi^p)). \end{aligned}$$

Furthermore, given that $\Delta \xi^p \in \ell_\infty$, it follows that $\mathfrak{N}^t(\Delta \xi^p) \in \ell_\infty$, [13]. Using this, the previous inequality implies that $\mathfrak{N}^t \Delta \xi \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ and consequently $\xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$. Hence the inclusion $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty} \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ holds.

Similarly, by making certain modifications, we can prove that the inclusion $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_c \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0} \subset \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ holds. \square

3. The α -, β - and γ - Duals of the Spaces $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$

Lemma 3.1 [6] *For an infinite matrix (a_{nk}) defined over the field of complex number, then the following condition holds:*

1. For $1 < p_n \leq Q < \infty$, then $A = (a_{nk}) \in (\ell(p), \ell_1)$ if and only if $R > 1 \in \mathbb{Z}$ such that

$$\sup_{N \in \mathcal{F}} \sum_k \left| \sum_{n \in \mathbb{N}} \mathfrak{M} a_{nk} R^{-1} \right|^{p'_n} < \infty, \text{ for every } n \in \mathbb{N}.$$

2. For $0 < p_n \leq 1$, then $A = (a_{nk}) \in (\ell(p), \ell_1)$ if and only if

$$\sup_{N \in \mathcal{F}} \sum_{k \in \mathbb{N}} \left| \sum_{n \in \mathbb{N}} \mathfrak{M} a_{nk} \right|^{p_n} < \infty, \text{ for every } n \in \mathbb{N}.$$

Lemma 3.2 [8] *For an infinite matrix (a_{nk}) defined over the field of complex number, then the following condition holds:*

1. For $1 < p_n \leq Q < \infty$, then $A = (a_{nk}) \in (\ell(p), \ell_\infty)$ if and only if there exists $R > 1 \in \mathbb{Z}$ such that

$$\sup_{n \in \mathbb{N}} \sum_k \mathfrak{M} |a_{nk} R^{-1}|^{p'_n} < \infty, \text{ for every } n \in \mathbb{N}. \quad (3.1)$$

2. For $0 < p_n \leq 1$, then $A = (a_{nk}) \in (\ell(p), \ell_\infty)$ if and only if

$$\sup_{n,k \in \mathbb{N}} \mathfrak{M}|a_{nk}|^{p_n} < \infty, \text{ for every } n \in \mathbb{N}. \quad (3.2)$$

Lemma 3.3 [8] For $0 < p_n \leq Q < \infty$, then $A = (a_{nk}) \in (\ell(p), c)$ if and only if inequality (3.1) and (3.2) holds for $n \in \mathbb{N}$ and $\gamma_k \in \mathbb{C}$ such that $a_{nk} \rightarrow \gamma_k$, for each $k \in \mathbb{N}$.

Theorem 3.1 For $1 \leq p_n \leq Q < \infty$, the α -dual of the spaces $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ is

$$\mathcal{D}_1 = \left\{ a \in \omega : \sup_{N \in \mathcal{F}} \sum_k \left| \sum_{n \in \mathbb{N}} \mathfrak{M}(-1)^{n-k} \mathcal{D}_{n-k} T_n \Delta a_n R^{-1} \right|^{p'_n} < \infty \right\}, n \in \mathbb{N}$$

Proof: For $a = (a_n) \in \omega$, consider the following inequality

$$a_n \xi_n = \sum_{k=0}^n \mathfrak{M}(-1)^{n-k} \mathcal{D}_{n-k} T_n \Delta a_n \chi_k = (\mathcal{C}\chi)_n, \text{ for all } n \in \mathbb{N} \quad (3.3)$$

where $\mathcal{C} = (\mathcal{C}_{nk})$ is defined as

$$\mathcal{C}_{nk} = \begin{cases} \mathfrak{M}(-1)^{n-k} \mathcal{D}_{n-k} T_n \Delta a_n, & 0 \leq k \leq n \\ 0, & k > n \end{cases},$$

for all $n, k \in \mathbb{N}$. Therefore, by combining equation (3.3) with part (a) of Lemma 3.1, we have $a\xi \in \ell_1$ when $\xi \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ or $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ if and only if $\mathcal{C}\chi \in \ell_1$ when $\chi \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_c$, $\mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$. Thus a is in α -dual of

$\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ if and only if $\mathcal{C} \in [\mathfrak{N}^t(\mathfrak{M}, \ell_1, \|\cdot, \cdot\|)_{c_0}]^\alpha = [\mathfrak{N}^t(\mathfrak{M}, \ell_1, \|\cdot, \cdot\|)_c]^\alpha = [\mathfrak{N}^t(\mathfrak{M}, \ell_1, \|\cdot, \cdot\|)_{\ell_\infty}]^\alpha$. Therefore $a \in [\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}]^\alpha = [\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c]^\alpha =$

$[\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}]^\alpha$ if and only if $\sup_{N \in \mathcal{F}} \sum_k \left| \sum_{n \in \mathbb{N}} \mathfrak{M}(\mathcal{C}_{nk} R^{-1}) \right|^{p'_n} < \infty$ which leads to the consequence

that

$$\mathcal{D}_1 = [\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}]^\alpha = [\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c]^\alpha = [\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}]^\alpha. \quad (3.4)$$

□

Theorem 3.2 Let $\mathcal{D}_2, \mathcal{D}_3, \mathcal{D}_4$ and \mathcal{D}_5 be sets defined as follow:

$$\mathcal{D}_2 = \left\{ a \in \omega : \sup_{n \in \mathbb{N}} \sum_{k \in \mathbb{N}} \left| \sum_{i=k}^n \mathfrak{M}(-1)^{i-k} \mathcal{Y} R^{-1} \right|^{p'_n} < \infty \right\}$$

$$\mathcal{D}_3 = cs$$

$$\mathcal{D}_4 = \left\{ a \in \omega : \sup_{N \in \mathcal{F}} \sum_{k \in \mathbb{N}} \left| \sum_{i=k}^n \mathfrak{M}(-1)^{n-k} \mathcal{P} \Delta a_n \right|^{p'_n} < \infty \right\}$$

$$\text{and } \mathcal{D}_5 = \left\{ a \in \omega : \sup_{n \in \mathbb{N}} \sum_{k \in \mathbb{N}} \left| \sum_{i=k}^n \mathfrak{M}(-1)^{i-k} \mathcal{P} \Delta a_i \right|^{p'_n} < \infty \right\}$$

where $\mathcal{Y} = \Delta a_i \mathcal{D}_{i-k} T_n$, $\mathcal{P} = \mathcal{D}_{i-k} T_n$, $0 < p_n \leq 1, \forall n \in \mathbb{N}$. Then,

$[\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}]^\beta = \mathcal{D}_2 \cap \mathcal{D}_3$, $[\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c]^\beta = \mathcal{D}_2 \cap \mathcal{D}_3 \cap \mathcal{D}_4$ and $[\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}]^\beta = \mathcal{D}_3 \cap \mathcal{D}_5$.

Proof: We now provide the β -dual for the space $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$. Consider the inequality

$$\sum_{k=0}^n a_k \xi_k = \sum_{k=0}^{n-1} \sum_{j=k}^n \mathfrak{M}(-1)^{j-k} \mathcal{P} \Delta a_j \chi_k + T_n \Delta a_n \chi_n = (E\chi)_n \quad (3.5)$$

where $\mathcal{P} = \mathcal{D}_{j-k} T_n$, for all $n \in \mathbb{N}$. We define the sequence $E = (e_{nk})$ as follow

$$e_{nk} = \begin{cases} \sum_{j=k}^n \mathfrak{M}(-1)^{j-k} \mathcal{D}_{j-k} T_n \Delta a_j, & 0 \leq k < n \\ T_n \Delta a_n, & k = n \\ 0, & k > n \end{cases}$$

for all $n, k \in \mathbb{N}$. Then from equation (3.4), we have $a\xi = (a_n \xi_n) \in cs$ when $\xi = (\xi_n) \in \mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$ if and only if $E\chi \in c$ when $\chi = (\chi_n) \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$. Therefore, $a = (a_n) \in [\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}]^\beta$ if and only if $E \in \mathfrak{N}^t(\mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$. Thus by part (1) of Lemma 3.2, we have

$$\sup_{n \in \mathbb{N}} \sum_k \left| \sum_{j=k}^n \mathfrak{M}(-1)^{j-k} \mathcal{D}_{j-k} T_n \Delta a_j R^{-1} \right|^{p'_n} < \infty$$

and by Lemma 3.3, $\lim_n a_{nk}$ exists for all $k \in \mathbb{N}$. Therefore $[\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}]^\beta = \mathcal{D}_2 \cap \mathcal{D}_3$. Similarly, we can obtained the β -dual for the spaces $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$. \square

Theorem 3.3 *The γ -dual of spaces $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{c_0}$, $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_c$ and $\mathfrak{N}^t(\Delta, \mathfrak{M}, p, \|\cdot, \cdot\|)_{\ell_\infty}$ is \mathcal{D}_2 .*

Proof: The proof of this theorem is directly follows from Theorem (3.2) with Lemma (3.3). \square

Acknowledgments

The authors thank to the reviewer for their valuable comments and fruitful suggestions which improve the presentation of the paper..

References

1. Bařar, F., Altay, B., Mursaleen, M., *Some generalizations of the space bv_p of p -bounded variation sequences*, Nonlinear Anal., 68(2), 273–287, (2008).
2. Et, M., olak, R., *On some generalized difference sequence spaces*, Soochow J. Math., 21(4), 377–386, (1995).
3. Freese, W. R., Cho, J., *Geometry of linear 2-normed spaces*, Nova Science Publishers, Inc., Hauppauge, NY, (2001).
4. Gähler, S., *2-metrische Räume und ihre topologische Struktur*, Math. Nachr., 26, 115–148 (1963).
5. Gunawan, H., Mashadi, *On finite-dimensional 2-normed spaces*, Soochow J. Math., 27(3), 321–329, (2001).
6. Grosse-Erdmann, K. G., *Matrix transformations between the sequence spaces of Maddox*, J. Math. Anal. Appl., 180(1), 223–238, (1993).
7. Kızmaz, H., *On certain sequence spaces*, Canad. Math. Bull., 24(2), 169–176 (1981).
8. Lascarides, G. C., Maddox, I. J., *Matrix transformations between some classes of sequences*, Proc. Cambridge Philos. Soc., 68, 99–104, (1970).
9. Lindenstrauss, J., Tzafriri, L., *On Orlicz sequence spaces*, Israel J. Math., 10, 379–390, (1971).
10. Maddox, I. J., *Elements of functional analysis*, Cambridge University Press, Cambridge, (1988).
11. Maddox, I. J., *Paranormed sequence spaces generated by infinite matrices*, Proc. Cambridge Philos. Soc., 64, 335–340, (1968).
12. Maddox, I. J., *Spaces of strongly summable sequences*, Quart. J. Math. Oxford Ser. (2), 18, 345–355 (1967).
13. Malkowsky, E., Savas, E., *Matrix transformations between sequence spaces of generalized weighted means*, Appl. Math. Comput., 147(2), 333–345, (2004).

14. Mears, M. F., *The inverse Nörlund mean*, Ann. of Math., 44, 401–410 (1943).
15. Mursaleen, M., Sharma, S. K., Kılıçman, A., *Sequence spaces defined by Musielak-Orlicz function over n -normed spaces*, Abstr. Appl. Anal., Art. ID 364743, pp. 10, (2013).
16. Mursaleen, M., *Generalized spaces of difference sequences*, J. Math. Anal. Appl., 203(3), 738–745, (1996).
17. Mursaleen, M., *On some new invariant matrix methods of summability*, Quart. J. Math. Oxford Ser., 34(133), 77–86, (1983).
18. Nakano, H., *Modulated sequence spaces*, Proc. Japan Acad., 27(2), 508–512 (1951).
19. Peyerimhoff, A., *Lectures on summability*, Lecture Notes in Mathematics, Vol. 107, (1969).
20. Raj, K., Sharma, S. K., *Some sequence spaces in 2-normed spaces defined by Musielak-Orlicz function*, Acta Univ. Sapientiae Math., 3(1), 97–109 (2011).
21. Raj, K., Sharma, S. K., Sharma, A. K. *Difference sequence spaces in n -normed spaces defined by Musielak-Orlicz function*, Armen. J. Math., 3(3), 127–141, (2010).
22. Raj, K., Sharma, A. K., Sharma, S. K., *A sequence space defined by Musielak-Orlicz function*, Int. J. Pure Appl. Math., 67(4), 475–484 (2011).
23. Simons, S., *The sequence spaces $l(p_\nu)$ and $m(p_\nu)$* , Proc. London Math. Soc., 15(3), 422–436 (1965).
24. Singh, S., Malik, T. A., *Paranormed Norlund N^t -difference sequence spaces and their α -, β - and γ -duals*, Proyecciones, 42(4), 879–892, (2023).
25. Wilansky, A., *Summability through functional analysis*, North-Holland Mathematics Studies, vol. 85, (1984).
26. Yeşilkayagil, M., Başar, F., *On the paranormed Nörlund sequence space of nonabsolute type*, Abstr. Appl. Anal., Art. ID 858704, pp. 9, (2014).

Sunil K. Sharma,
 School of Mathematics,
 Shri Mata Vaishno Devi University, Kakryal, Katra-182320, J&K,
 India.
 E-mail address: sunilksharma42@gmail.com

and

Ravi Kumar,
 Department of Mathematics,
 Central University of Jammu, Rahya Suchani (Bagla), Samba-181143, J&K,
 India.
 E-mail address: ravikumar13849@gmail.com

and

Ayhan Esi,
 Department of Basic Engi. Sci. (Math.Sect.),
 Faculty of Engineering and Natural Sciences, Malatya Turgut Ozal University-44200, Malatya,
 Türkiye.
 E-mail address: aesi23@hotmail.com; ayhan.esi@ozal.edu.tr

and

Ajay K. Sharma,
 Department of Mathematics,
 Central University of Jammu, Rahya Suchani (Bagla), Samba-181143, J&K,
 India.
 E-mail address: aksju.76@yahoo.com