Bol. Soc. Paran. Mat.
©SPM – E-ISSN-2175-1188
SPM: www.spm.uem.br/bspm

(3s.) **v. 2025 (43) 2** : 1–11.
ISSN-0037-8712
doi:10.5269/bspm.79228

A Note on Type 2 Degenerate Changhee Numbers and Polynomials

Waseem Ahmad Khan

ABSTRACT: In this paper, we introduce type 2 degenerate Changhee numbers and polynomials and investigate some properties of these numbers and polynomials. We introduce higher-order type 2 degenerate Changhee polynomials and numbers and derive their explicit expressions and some identities involving them. In addition, we give some new relations between the type 2 degenerate Changhee polynomials and degenerate Euler polynomials.

Key Words: degenerate Euler polynomials, type 2 degenerate Changhee polynomials, type 2 degenerate Euler polynomials, higher-order type 2 degenerate Changhee polynomials and numbers.

Contents

1 Introduction **1**

2 Type 2 degenerate Changhee numbers and polynomials **3**

3 Type 2 higher-order degenerate Changhee polynomials **6**

4 Conclusion **10**

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$ and $\mathbb{C}_p$ will denote the ring of p-adic integers, the filed of p-adic rational numbers and the completion of an algebraic closure of $\mathbb{Q}_p$. The p-adic norm $|\ .\ |_p$ is normalized by $|\ p\ |_p = \frac{1}{p}$. Let $C(\mathbb{Z}_p)$ be the space of continuous function on $\mathbb{Z}_p$. For $f \in C(\mathbb{Z}_p)$, the fermionic p-adic integral on $\mathbb{Z}_p$ is defined by Kim as follows

$$I(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)\mu_{-1}(x + p^N\mathbb{Z}_p)$$

$$= \lim_{N\to\infty} \sum_{x=0}^{p^N-1} f(x)(-1)^x, (\text{see [3, 21, 22]}) \tag{1.1}$$

From (1.1), we note that

$$I(f_n) + (-1)^{n-1} I(f) = 2\sum_{a=0}^{n-1} (-1)^{n-1-a} f(a), (\text{see [21, 22]}), \tag{1.2}$$

where $f_n(x) = f(x+n), (n \in \mathbb{N})$.

Let the Changhee polynomials are defined by the generating function as follows (see [5, 6])

$$\frac{2}{2+t}(1+t)^x = \sum_{n=0}^{\infty} Ch_n(x)\frac{t^n}{n!}. \tag{1.3}$$

2010 *Mathematics Subject Classification:* 11B83, 11B73, 05A19.
Submitted September 28, 2025. Published December 19, 2025

Typeset by $\mathcal{B}^{S}\mathcal{P}_{\mathcal{M}}$ style.
© Soc. Paran. de Mat.

Letting $x=0$, $Ch_n=Ch_n(0),(n\geq 0)$ are called the Changhee numbers. From (1.3), we note that

$$\int_{\mathbb{Z}_p}(1+t)^{x+y}d\mu_{-1}(y)=\frac{2}{2+t}(1+t)^x=\sum_{n=0}^{\infty}Ch_n(x)\frac{t^n}{n!}. \tag{1.4}$$

Thus, by (1.4), we have

$$\int_{\mathbb{Z}_p}(x+y)_n d\mu_{-1}(y)=Ch_n(x),(n\geq 0),(\text{see [6]}), \tag{1.5}$$

where $(x)_0=1,(x)_n=x(x-1)\cdots(x-n+1),(n\geq 1)$,

As well known, the type 2 Euler polynomials are defined by

$$\frac{2}{e^t+e^{-t}}e^{xt}=\sum_{n=0}^{\infty}E_n^*(x)\frac{t^n}{n!}. \tag{1.6}$$

In the case $x=0$, $E_n^*=E_n^*(0)$ are called the type 2 Euler numbers.

By using (1.1) and (1.6), we note that

$$\int_{\mathbb{Z}_p}e^{(2y+x+1)t}d\mu_{-1}(y)=\frac{2}{e^t+e^{-t}}e^{xt}=\sum_{n=0}^{\infty}E_n^*(x)\frac{t^n}{n!}. \tag{1.7}$$

By (1.7), we get

$$\int_{\mathbb{Z}_p}(2y+x+1)^n d\mu_{-1}(y)=E_n^*(x),(n\geq 0),(\text{see [8, 9, 10]}), \tag{1.8}$$

For $n\geq 0$, the Stirling numbers of the first kind are defined by

$$(x)_n=\sum_{l=0}^{n}S_1(n,l)x^l,(\text{see [1-10]}), \tag{1.9}$$

where $(x)_0=1$, and $(x)_n=x(x-1)\cdots(x-n+1),(n\geq 1)$. From (1.9), it is easily to see that

$$\frac{1}{k!}(\log(1+t))^k=\sum_{n=k}^{\infty}S_1(n,k)\frac{t^n}{n!},\quad (k\geq 0),(\text{see [11-20]}). \tag{1.10}$$

In the inverse expression to (1.9), the Stirling numbers of the second kind are defined by

$$x^n=\sum_{l=0}^{n}S_2(n,l)(x)_l,(\text{see [21-26]}). \tag{1.11}$$

From (1.11), it is easily to see that

$$\frac{1}{k!}(e^t-1)^k=\sum_{n=l}^{\infty}S_2(n,l)\frac{t^n}{n!},(\text{see [1-15]}). \tag{1.12}$$

From (1.3), (1.6), (1.9) and (1.10), we get

$$Ch_n(x)=\sum_{l=0}^{n}E_l^*(x)S_1(n,l), \tag{1.13}$$

and

$$E_n^*(x)=\sum_{l=0}^{n}Ch_n(x)S_2(n,l),(\text{see [6]}) \tag{1.14}$$

Recently, Kim-Kim [12] introduced the type 2 Changhee polynomials are defined by

$$\int_{\mathbb{Z}_p}(1+t)^{2y+x+1}d\mu_{-1}(y)=\frac{2}{(1+t)+(1+t)^{-1}}(1+t)^x$$
$$=\sum_{n=0}^{\infty}c_n(x)\frac{t^n}{n!}, \tag{1.15}$$

When $x=0$, $c_n=c_n(0)$ are called the type 2 Changhee numbers.

For any $\lambda\in\mathbb{R}$, degenerate version of the exponential function $e_\lambda^x(t)$ is defined as follows (see [16-26])

$$e_\lambda^x(t):=(1+\lambda t)^{\frac{x}{\lambda}}=\sum_{n=0}^{\infty}(x)_{n,\lambda}\frac{t^n}{n!}, \tag{1.16}$$

Kim introduced the degenerate Stirling numbers of the second kind (see [24]) are given by

$$\frac{1}{k!}(e_\lambda(t)-1)^k=\sum_{n=k}^{\infty}S_{2,\lambda}(n,k)\frac{t^n}{n!},\quad (k\geq 0). \tag{1.17}$$

It is clear that $\lim_{\lambda\to 0}S_{2,\lambda}(n,k)=S_2(n,k)$, where $S_2(n,k)$ are called the Stirling numbers of the second.

Recently, Kim [23] introduced the degenerate Changhee polynomials of the second kind are defined by

$$\int_{\mathbb{Z}_p}(1+\lambda\log(1+t))^{\frac{x+y}{\lambda}}d\mu_{-1}(y)=\frac{2}{1+(1+\lambda\log(1+t))}(1+\lambda\log(1+t))^{\frac{x}{\lambda}}$$
$$=\sum_{n=0}^{\infty}Ch_{n,\lambda}(x)\frac{t^n}{n!}, \tag{1.18}$$

where $\lambda\in\mathbb{C}_p$ with $\mid\lambda\mid_p\leq 1$.

When $x=0$, $Ch_{n,\lambda}=Ch_{n,\lambda}(0)$ are called the degenerate Changhee numbers of the second kind.

This paper is organized as follows. In Sect 2, we consider type 2 degenerate Changhee numbers and polynomials and investigate some properties of these numbers and polynomials. In Sect 3, we introduce higher-order type 2 degenerate Changhee polynomials and numbers which can be represented in terms of p-adic integrals on $\mathbb{Z}_p$. We derive their explicit expressions and some other polynomials. Moreover, we obtain identities involving those polynomials and some other special numbers and polynomials.

2. Type 2 degenerate Changhee numbers and polynomials

In this section, let us assume that $\lambda\in\mathbb{C}_p$ and $t\in\mathbb{C}_p$ with the condition $\mid\lambda t\mid_p< p^{-\frac{1}{p-1}}$. As is well-known, the type 2 degenerate Euler polynomials are defined by the generating function

$$\int_{\mathbb{Z}_p}e_\lambda^{2y+x+1}(t)d\mu_{-1}(y)=\frac{2}{e_\lambda(t)+e_\lambda^{-1}(t)}e_\lambda^x(t)$$
$$=\sum_{n=0}^{\infty}E_{n,\lambda}(x)\frac{t^n}{n!},\text{(see [8])}. \tag{2.1}$$

From (2.1), we have

$$\int_{\mathbb{Z}_p}(2y+x+1)_{n,\lambda}d\mu_{-1}(y)=E_{n,\lambda}(x),(n\geq 0). \tag{2.2}$$

When $x=0$, $E_{n,\lambda}=E_{n,\lambda}(0)$ are called the type 2 degenerate Euler numbers.

Inspired by (1.15) and (2.1), we introduce type 2 degenerate Changhee polynomials are defined by

$$\int_{\mathbb{Z}_p}(1+\lambda\log(1+t))^{\frac{2y+x+1}{\lambda}}d\mu_{-1}(y)=\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}}+(1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}(1+\lambda\log(1+t))^{\frac{x}{\lambda}}$$

$$=\sum_{n=0}^{\infty}\widehat{C}_{n,\lambda}(x)\frac{t^n}{n!}. \tag{2.3}$$

Note that, $\lim_{\lambda\to 0}\widehat{C}_{n,\lambda}(x)=c_n(x)$, $(n\geq 0)$, (see [12]). We note that $x=0,\widehat{C}_{n,\lambda}=\widehat{C}_{n,\lambda}(0)$ are called the type 2 degenerate Changhee.

Theorem 2.1. For $n\geq 0$, we have

$$\tilde{C}_{n,\lambda}(x)=\sum_{l=0}^{n}S_1(n,l)\int_{\mathbb{Z}_p}\binom{\frac{2y+x+1}{\lambda}}{l}l!d\mu_{-1}(y)\lambda^l$$

$$=\sum_{l=0}^{n}\int_{\mathbb{Z}_p}(2y+x+1)_{l,\lambda}d\mu_{-1}(y)\lambda^l S_1(n,l), \tag{2.4}$$

where $(x)_{0,\lambda}=1$ and $(x)_{n,\lambda}=x(x-\lambda)\cdots(x-(n-1)\lambda)$ for $n\geq 1$.

Proof. Using (2.3), we note that

$$\int_{\mathbb{Z}_p}(1+\lambda\log(1+t))^{\frac{2y+x+1}{\lambda}}d\mu_{-1}(y)=\sum_{l=0}^{\infty}\int_{\mathbb{Z}_p}\binom{\frac{2y+x+1}{\lambda}}{l}d\mu_{-1}(y)\lambda^l(\log(1+t))^l$$

$$=\sum_{n=0}^{\infty}\left(\sum_{l=0}^{n}S_1(n,l)\int_{\mathbb{Z}_p}\binom{\frac{2y+x+1}{\lambda}}{l}l!d\mu_{-1}(y)\lambda^l\right)\frac{t^n}{n!}. \tag{2.5}$$

Comparing the coefficients of on both sides of (2.3) and (2.5), we obtain the result (2.4).

Theorem 2.2. For $n\geq 0$, we have

$$E_{n,\lambda}(x)=\sum_{m=0}^{n}\widehat{C}_{m,\lambda}(x)S_2(n,m). \tag{2.6}$$

Proof By replacing t by e^t-1 in (2.3) and using (1.12), we get

$$\sum_{m=0}^{\infty}\widehat{C}_{m,\lambda}(x)\frac{(e^t-1)^m}{m!}=\frac{2}{e_\lambda(t)+e_\lambda^{-1}(t)}e_\lambda^x(t)$$

$$=\sum_{n=0}^{\infty}E_{n,\lambda}(x)\frac{t^n}{n!}. \tag{2.7}$$

On the other hand,

$$\sum_{m=0}^{\infty}\widehat{C}_{m,\lambda}(x)\frac{(e^t-1)^m}{m!}=\sum_{m=0}^{\infty}\widehat{C}_{m,\lambda}(x)\sum_{n=m}^{\infty}S_2(n,m)\frac{t^n}{n!}$$

$$=\sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}\widehat{C}_{m,\lambda}(x)S_2(n,m)\right)\frac{t^n}{n!}. \tag{2.8}$$

By (2.7) and (2.8), we get the result.

Theorem 2.3. For $n \geq 0$, we have

$$\widehat{C}_{n,\lambda}(x) = \sum_{k=0}^{n} \sum_{m=0}^{k} \binom{n}{k} (x)_{m,\lambda} S_{1,\lambda}(k,m) \widehat{C}_{n-k,\lambda}. \tag{2.9}$$

Proof. From (2.3), we note that

$$\begin{aligned}
\sum_{n=0}^{\infty} \widehat{C}_{n,\lambda}(x) \frac{t^n}{n!} &= \frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}} + (1+\lambda\log(1+t))^{-\frac{1}{\lambda}}} (1+\lambda\log(1+t))^{\frac{x}{\lambda}} \\
&= \left(\sum_{n=0}^{\infty} \widehat{C}_{n,\lambda} \frac{t^n}{n!} \right) \left(\sum_{m=0}^{\infty} \binom{\frac{x}{\lambda}}{m} (\log(1+t))^m \right) \\
&= \left(\sum_{n=0}^{\infty} \widehat{C}_{n,\lambda} \frac{t^n}{n!} \right) \left(\sum_{m=0}^{\infty} (x)_{m,\lambda} \sum_{k=m}^{\infty} S_1(k,m) \frac{t^k}{k!} \right) \\
&= \left(\sum_{n=0}^{\infty} \widehat{C}_{n,\lambda} \frac{t^n}{n!} \right) \left(\sum_{k=0}^{\infty} \left(\sum_{m=0}^{k} (x)_{m,\lambda} S_1(k,m) \right) \frac{t^k}{k!} \right) \\
&= \sum_{n=0}^{\infty} \left(\sum_{k=0}^{n} \sum_{m=0}^{k} \binom{n}{k} (x)_{m,\lambda} S_{1,\lambda}(k,m) \widehat{C}_{n-k,\lambda} \right) \frac{t^n}{n!}.
\end{aligned} \tag{2.10}$$

Therefore, by (2.4) and (2.10), we obtain at the required result.

Theorem 2.4. For $n \geq 0$, we have

$$\widehat{C}_{n,\lambda}(1) + \widehat{C}_{n,\lambda} = \begin{cases} 2, & \text{if } n = 0, \\ 0, & \text{if } n \geq 1, \end{cases} \tag{2.11}$$

Proof. By (1.2), we easily get

$$\int_{\mathbb{Z}_p} f(x+1) d\mu_{-1}(x) + \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = 2f(0). \tag{2.12}$$

Now, equation (2.12) can be written as

$$\int_{\mathbb{Z}_p} (1+\lambda\log(1+t))^{\frac{2y+2}{\lambda}} d\mu_{-1}(y) + \int_{\mathbb{Z}_p} (1+\lambda\log(1+t))^{\frac{2y+1}{\lambda}} d\mu_{-1}(y) = 2. \tag{2.13}$$

From (2.3) and (2.13), we have

$$\begin{aligned}
&\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}} + (1+\lambda\log(1+t))^{-\frac{1}{\lambda}}} (1+\lambda\log(1+t))^{\frac{1}{\lambda}} \\
&\quad + \frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}} + (1+\lambda\log(1+t))^{-\frac{1}{\lambda}}} = 2.
\end{aligned} \tag{2.14}$$

From (2.11) and (2.14), we have

$$\sum_{n=0}^{\infty} \left(\widehat{C}_{n,\lambda}(1) + \widehat{C}_{n,\lambda} \right) \frac{t^n}{n!} = 2. \tag{2.15}$$

In view of (2.15), we complete the proof.

Theorem 2.6. For $n \geq 0$, we have

$$\widehat{C}_{n,\lambda}(x+1) + \widehat{C}_{n,\lambda}(x) = \sum_{k=0}^{n} \binom{n}{k} \sum_{m=0}^{k} \widehat{C}_{n-k,\lambda}(x) (1)_{m,\lambda} S_1(k,m). \tag{2.16}$$

Proof. Suppose that

$$\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}}+(1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}(1+\lambda\log(1+t))^{\frac{x+1}{\lambda}}+\frac{2(1+\lambda\log(1+t))^{\frac{x}{\lambda}}}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}}+(1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}$$

$$=\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}}+(1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}(1+\lambda\log(1+t))^{\frac{x}{\lambda}}(1+\lambda\log(1+t))^{\frac{1}{\lambda}}. \tag{2.17}$$

Thus, by (2.3) and (2.17), we get

$$\sum_{n=0}^{\infty}\left(\widehat{C}_{n,\lambda}(x+1)+\widehat{C}_{n,\lambda}(x)\right)\frac{t^n}{n!}$$

$$=\left(\sum_{n=0}^{\infty}\widehat{C}_{n,\lambda}(x)\frac{t^n}{n!}\right)\left(\sum_{m=0}^{\infty}(1)_{m,\lambda}\frac{1}{m!}(\log(1+t))^m\right)$$

$$=\left(\sum_{n=0}^{\infty}\widehat{C}_{n,\lambda}(x)\frac{t^n}{n!}\right)\left(\sum_{k=0}^{\infty}\sum_{m=0}^{k}(1)_{m,\lambda}S_1(k,m)\frac{t^k}{k!}\right)$$

$$=\sum_{n=0}^{\infty}\left(\sum_{k=0}^{n}\binom{n}{k}\sum_{m=0}^{k}\widehat{C}_{n-k,\lambda}(x)(1)_{m,\lambda}S_1(k,m)\right)\frac{t^n}{n!}. \tag{2.18}$$

By comparing the coefficients of t, we get (2.16).

3. Type 2 higher-order degenerate Changhee polynomials

In this section, we introduce type 2 degenerate Changhee polynomials of order r which are derived from the multivariate fermionic p-adic integral on $\mathbb{Z}_p$.

For $r\in\mathbb{N}$, we define the type 2 degenerate Changhee polynomials of order r which are given multivariate fermionic p-adic integral on $\mathbb{Z}_p$ as follows:

$$\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(1+\lambda\log(1+t))^{\frac{2(x_1+\cdots+x_r)+x+1}{\lambda}}d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)$$

$$=\left(\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}}+(1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}\right)^r(1+\lambda\log(1+t))^{\frac{x}{\lambda}}$$

$$=\sum_{n=0}^{\infty}\widehat{C}_{n,\lambda}^{(r)}(x)\frac{t^n}{n!}. \tag{3.1}$$

When $x=0,\widehat{C}_{n,\lambda}^{(r)}=\widehat{C}_{n,\lambda}^{(r)}(0)$ are called the type 2 degenerate Changhee numbers of order α.

Theorem 3.1. For $n\geq 0$ and $r\in\mathbb{N}$, we have

$$\widehat{C}_{n,\lambda}^{(r)}(x)=\sum_{m=0}^{n}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(2(x_1+\cdots+x_r)+x+1)_{\lambda,m}d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)S_1(n,m)$$

Proof. From (3.1), we note that

$$\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(1+\lambda\log_\lambda(1+t))^{\frac{2(x_1+\cdots+x_r)+x+1}{\lambda}}d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)$$

$$=\sum_{m=0}^{\infty}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}\binom{\frac{2(x_1+\cdots+x_r)+x+1}{\lambda}}{m}d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)\lambda^m(\log(1+t)))^m$$

$$= \sum_{m=0}^{\infty} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (2(x_1+\cdots+x_r)+x+1)_{\lambda,m} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) \frac{1}{m!}(\log(1+t)))^m$$

$$= \sum_{n=0}^{\infty} \left(\sum_{m=0}^{n} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (2(x_1+\cdots+x_r)+x+1)_{\lambda,m} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r) S_1(n,m) \right) \frac{t^n}{n!}. \tag{3.2}$$

For $r \in \mathbb{N}$, we have

$$\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (1+\lambda t)^{\frac{2(x_1+\cdots+x_r)+x+1}{\lambda}} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)$$
$$= \left(\frac{2}{(1+\lambda t)^{\frac{1}{\lambda}} + (1+\lambda t)^{-\frac{1}{\lambda}}} \right)^r (1+\lambda t)^{\frac{x}{\lambda}}. \tag{3.3}$$

Now, we define the type 2 degenerate Euler polynomials of order r which are given by

$$\left(\frac{2}{(1+\lambda t)^{\frac{1}{\lambda}} + (1+\lambda t)^{-\frac{1}{\lambda}}} \right)^r (1+\lambda t)^{\frac{x}{\lambda}} = \sum_{n=0}^{\infty} E_{n,\lambda}^{(r)}(x) \frac{t^n}{n!}. \tag{3.4}$$

Thus, by (3.3) and (3.4) we get

$$\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (2(x_1+\cdots+x_r)+x+1)_{\lambda,m} d\mu_{-1}(x_1)\cdots d\mu_{-1} = E_{n,\lambda}^{(r)}(x), (m \geq 0). \tag{3.5}$$

Theorem 3.2. For $n \geq 0$, we have

$$\widehat{C}_{n,\lambda}^{(r)}(x) = \sum_{m=0}^{n} E_{m,\lambda}^{(r)} S_1(n,m).$$

Proof. By using (3.2), (3.4) and (3.5), we obtain the result.

Theorem 3.3. For $n \geq 0$, we have

$$E_{n,\lambda}^{(r)}(x) = \sum_{m=0}^{n} \widehat{C}_{m,\lambda}^{(r)}(x) S_2(n,m).$$

Proof. By changing t by $e^t - 1$ in (3.1), we have

$$\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} (1+\lambda t)^{\frac{2(x_1+\cdots+x_r)+x+1}{\lambda}} d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)$$
$$= \sum_{m=0}^{\infty} \widehat{C}_{m,\lambda}^{(r)}(x) \frac{(e^t-1)^m}{m!}$$
$$= \sum_{n=0}^{\infty} \left(\sum_{m=0}^{n} \widehat{C}_{m,\lambda}^{(r)}(x) S_2(n,m) \right) \frac{t^n}{n!}. \tag{3.6}$$

Therefore, by (3.3) and (3.6), we get the result.

Theorem 3.4. For $n \geq 0$, we have

$$\widehat{C}_{n,\lambda}^{(r)}(x) = \sum_{l=0}^{n} \binom{n}{l} \widehat{C}_{n-l,\lambda}^{(r)} \widehat{C}_{l,\lambda}^{(r-k)}(x).$$

Proof. From (3.1), we have

$$\sum_{n=0}^{\infty} \widehat{C}_{n,\lambda}^{(r)}(x) \frac{t^n}{n!} = \left(\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}} + (1+\lambda\log(1+t))^{-\frac{1}{\lambda}}} \right)^r (1+\lambda\log(1+t))^{\frac{x}{\lambda}}$$

$$= \left(\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}} + (1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}\right)^{r}$$

$$\times \left(\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}} + (1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}\right)^{r-k} (1+\lambda\log(1+t))^{\frac{x}{\lambda}}$$

$$= \left(\sum_{n=0}^{\infty} \widehat{C}_{n,\lambda}^{(r)} \frac{t^n}{n!}\right) \left(\sum_{l=0}^{\infty} \widehat{C}_{l,\lambda}^{(r-k)}(x) \frac{t^l}{l!}\right)$$

$$= \sum_{n=0}^{\infty} \left(\sum_{l=0}^{n} \binom{n}{l} \widehat{C}_{n-l,\lambda}^{(r)} \widehat{C}_{l,\lambda}^{(r-k)}(x)\right) \frac{t^n}{n!}. \tag{3.7}$$

In view of (3.7), we complete the proof.

Theorem 3.5. For $n \geq 0$, we have

$$\widehat{C}_{n,\lambda}^{(r)}(x) = \sum_{k=0}^{n} \sum_{m=0}^{k} \binom{n}{k} (x)_{m,\lambda} S_{1,\lambda}(k,m) \widehat{C}_{n-k,\lambda}^{(r)}. \tag{3.8}$$

Proof. From (3.1), we note that

$$\sum_{n=0}^{\infty} \widehat{C}_{n,\lambda}^{(r)}(x) \frac{t^n}{n!} = \left(\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}} + (1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}\right)^{r} (1+\lambda\log(1+t))^{\frac{x}{\lambda}}$$

$$= \left(\sum_{n=0}^{\infty} \widehat{C}_{n,\lambda}^{(r)} \frac{t^n}{n!}\right) \left(\sum_{m=0}^{\infty} \binom{\frac{x}{\lambda}}{m} (\log(1+t))^m\right)$$

$$= \left(\sum_{n=0}^{\infty} \widehat{C}_{n,\lambda}^{(r)} \frac{t^n}{n!}\right) \left(\sum_{m=0}^{\infty} (x)_{m,\lambda} \sum_{k=m}^{\infty} S_1(k,m) \frac{t^k}{k!}\right)$$

$$= \left(\sum_{n=0}^{\infty} \widehat{C}_{n,\lambda}^{(r)} \frac{t^n}{n!}\right) \left(\sum_{k=0}^{\infty} \left(\sum_{m=0}^{k} (x)_{m,\lambda} S_1(k,m)\right) \frac{t^k}{k!}\right)$$

$$= \sum_{n=0}^{\infty} \left(\sum_{k=0}^{n} \sum_{m=0}^{k} \binom{n}{k} (x)_{m,\lambda} S_{1,\lambda}(k,m) \widehat{C}_{n-k,\lambda}^{(r)}\right) \frac{t^n}{n!}. \tag{3.9}$$

Therefore, by (3.1) and (3.9), we obtain at the required result.

Theorem 3.6. For $n \geq 0$, we have

$$\sum_{m=0}^{n} \widehat{C}_{m,\lambda}^{(r)}(x) S_2(n,m) = \sum_{m=0}^{n} S_{2,\lambda}(n,m) C_m^{(r)}(x).$$

Proof. Now, we observe that

$$(1+\lambda t)^{\frac{2(x_1+\cdots+x_r)+x+1}{\lambda}} = \left((1+\lambda t)^{\frac{1}{\lambda}} - 1 + 1\right)^{2(x_1+\cdots+x_r)+x+1}$$

$$= \sum_{m=0}^{\infty} \binom{2(x_1+\cdots+x_r)+x+1}{m} ((1+\lambda t)^{\frac{1}{\lambda}} - 1)^m$$

$$= \sum_{m=0}^{\infty} (2(x_1+\cdots+x_r)+x+1)_m \sum_{n=m}^{\infty} S_{2,\lambda}(n,m) \frac{t^n}{n!}$$

$$= \sum_{n=0}^{\infty} \left(\sum_{m=0}^{n} S_{2,\lambda}(n,m)(2(x_1+\cdots+x_r)+x+1)_m\right) \frac{t^n}{n!}. \tag{3.10}$$

Thus, by (3.5) and (3.10), we get

$$\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{2(x_1+\cdots+x_r++x+1}{\lambda}}d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)$$

$$=\sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}S_{2,\lambda}(n,m)\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(2(x_1+\cdots+x_r)+x+1)_m d\mu_{-1}(x_1)\cdots d\mu_{-1}(x_r)\right)\frac{t^n}{n!}$$

$$=\sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}S_{2,\lambda}(n,m)C_m^{(r)}(x)\right)\frac{t^n}{n!}. \tag{3.11}$$

Therefore, by (3.5) and (3.11), we obtain the result.

Theorem 3.7. For $n\geq 0$, we have

$$\widehat{C}_{n,\lambda}^{(r)}(x)=\sum_{m=0}^{n}E_{m,\lambda}^{(r)}(x)S_1(n,m).$$

Proof. By replacing t by $\log(1+t)$ in (3.4), we have

$$\left(\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}}+(1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}\right)^r(1+\lambda\log(1+t))^{\frac{x}{\lambda}}=\sum_{m=0}^{\infty}E_{m,\lambda}^{(r)}(x)\frac{(\log(1+t))^m}{m!}$$

$$=\sum_{m=0}^{\infty}E_{m,\lambda}^{(r)}(x)\sum_{n=m}^{\infty}S_1(n,m)\frac{t^n}{n!}$$

$$=\sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}E_{m,\lambda}^{(r)}(x)S_1(n,m)\right)\frac{t^n}{n!}. \tag{3.12}$$

On the other hand, we have

$$\left(\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}}+(1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}\right)^r(1+\lambda\log(1+t))^{\frac{x}{\lambda}}=\sum_{n=0}^{\infty}\widehat{C}_{n,\lambda}^{(r)}(x)\frac{t^n}{n!}. \tag{3.13}$$

In view of (3.12) and (3.13), we obtain the result.

Theorem 3.8. For $n\geq 0$, we have

$$\widehat{C}_{n,\lambda}^{(r)}(x+y)=\sum_{k=0}^{n}\sum_{m=0}^{k}\binom{n}{k}(y)_{m,\lambda}S_1(k,m)\widehat{C}_{n-k,\lambda}^{(r)}(x).$$

Proof. From (3.1), we have

$$\sum_{n=0}^{\infty}\widehat{C}_{n,\lambda}^{(r)}(x+y)\frac{t^n}{n!}=\left(\frac{2}{(1+\lambda\log(1+t))^{\frac{1}{\lambda}}+(1+\lambda\log(1+t))^{-\frac{1}{\lambda}}}\right)^r(1+\lambda\log(1+t))^{\frac{x+y}{\lambda}}$$

$$=\left(\sum_{n=0}^{\infty}\widehat{C}_{n,\lambda}^{(r)}(x)\frac{t^n}{n!}\right)\left(\sum_{m=0}^{\infty}\binom{\frac{y}{\lambda}}{m}(\log(1+t))^m\right)$$

$$=\left(\sum_{n=0}^{\infty}\widehat{C}_{n,\lambda}^{(r)}\frac{t^n}{n!}(x)\right)\left(\sum_{m=0}^{\infty}(y)_{m,\lambda}\sum_{k=m}^{\infty}S_1(k,m)\frac{t^k}{k!}\right)$$

$$=\left(\sum_{n=0}^{\infty}\widehat{C}_{n,\lambda}^{(r)}(x)\frac{t^n}{n!}\right)\left(\sum_{k=0}^{\infty}\left(\sum_{m=0}^{k}(y)_{m,\lambda}S_1(k,m)\right)\frac{t^k}{k!}\right)$$

$$=\sum_{n=0}^{\infty}\left(\sum_{k=0}^{n}\sum_{m=0}^{k}\binom{n}{k}(y)_{m,\lambda}S_1(k,m)\widehat{C}_{n-k,\lambda}^{(r)}(x)\right)\frac{t^n}{n!}. \tag{3.14}$$

Thus, by (3.14), we obtain at the required result.

4. Conclusion

In this article, we introduced type 2 degenerate Changhee numbers and polynomials and investigated some properties of these numbers and polynomials. We introduced higher-order type 2 degenerate Changhee polynomials and numbers and derived their explicit expressions and some identities involving them. In addition, we given some new relations between the type 2 degenerate Changhee polynomials and degenerate Euler polynomials.

References

[1] M. S. Alatawi, W. A. Khan, *New type of degenerate Changhee-Genocchi polynomials*, Axioms, **2022**, 11, 355. https://doi.org/10.3390/axioms11080355.

[2] D. V. Dolgy, W. A. Khan, *A note on type two degenerate poly-Changhee polynomials of the second kind*, Symmetry. **13(579)**(2021), 1-12.

[3] L. C. Jang, D. S. Kim, T. Kim, H. Lee, *p-Adic integral on $\mathbb{Z}_p$ associated with degenerate Bernoulli polynomials of the second kind*, Adv. Diff. Equ. **2020**, 278, 1-20.

[4] H.-I. Kwon, T. Kim, J. J. Seo, *A note on degenerate Changhee numbers and polynomials*, Proc. Jangjeon Math. Soc., **18(3)**(2015), 295-305.

[5] D. S. Kim, T. Kim, J. J. Seo, *A note on Changhee polynomials and numbers*, Adv. Studies Theo. Phys., **7(20)**(2013), 993-1003.

[6] D. S. Kim, J. J. Seo, S.-H. Lee, *Higher-order Changhee numbers and polynomials*, Adv. Studies Theo. Phys., **8(8)**(2014), 365-373.

[7] D. S. Kim, H. Y. Kim, D. Kim, T. Kim, *Identities of symmetry for type 2 Bernoulli and Euler polynomials*, Symmetry **613(11)**(2019).

[8] D. S. Kim, T. Kim, C. S. Ryoo, *Generalized type 2 degenerate Euler numbers*, Adv. Stud. Contemp. Math., **30(2)**(2020), 165-169.

[9] D. S. Kim, T. Kim, *A note on new type of degenerate Bernoulli numbers*, Russ. J. Math. Phys. **27(2)**(2020), 227-235.

[10] G. W. Jang, T. Kim, *A note on type 2 degenerate Euler and Bernoulli polynomials*, Adv. Stud. Contemp. Math., **29(1)**(2019), 147-159.

[11] T. Kim, D. S. Kim, *Degenerate central factorial numbers of the second kind*, Rev. R. Acad. Cienc. Exactas. Fs. Nat. Ser. A Mat. RACSAM, **113(4)** (2019), 3359-3367.

[12] T. Kim, D. S. Kim, *A note on type 2 Changhee and Daehee polynomials*, Rev. R. Acad. Cienc. Exactas. Fs. Nat. Ser. A Mat. RACSAM, **113(3)** (2019), 2783-2791.

[13] W. A. Khan, M. S. Alatawi, *A note on modified degenerate Changhee-Genocchi polynomials of the second kind*, Symmetry, **15:136** (2023) , 1-12.

[14] W. A. Khan, V. Yadav, *A study on q-analogue of degenerate Changhee numbers and polynomials*, Southeast Asian Journal of Mathematics and Mathematical Sciences, **19(1)** (2023), 29-42.

[15] W. A. Khan, *A note on q-analogues of degenerate Catalan-Daehee numbers and polynomials*, Journal of Mathematics, **2022**, Volume 2022, Article ID 9486880, 9 pages.

[16] W. A. Khan, *A note on q-analogue of degenerate Catalan numbers associated p-adic integral on $\mathbb{Z}_p$*, Symmetry, **14(119)**(2022), 1-10.

[17] W. A. Khan, *A study on q-analogue of degenerate $\frac{1}{2}$-Changhee numbers and polynomials*, Southeast Asian Journal of Mathematics and Mathematical Sciences, **18(2)** (2022), 1-12.

[18] W. A. Khan, H. Haroon, *Higher-order degenerate Hermite-Bernoulli arising from p-adic integral on $\mathbb{Z}_p$*, Iranian Journal of Mathematical Sciences and Informatics, **17(2)** (2022), 171-189.

[19] W. A. Khan, J. Younis, U. Duran, A. Iqbal, *The higher-order type Daehee polynomials associated with p-adic integrals on $\mathbb{Z}_p$*, Applied Mathematics in Science and Engineering, **30(1)** (2022) , 573-582.

[20] W. A. Khan, M. Acikgoz, U. Duran, *Note on the type 2 degenerate multi-poly-Euler polynomials*, Symmetry. **12**(2020), 1-10.

[21] T. Kim, *Some identities on the q-Euler polynomials of higher order and q-Stirling numbers by the fermionic p-adic integral on $\mathbb{Z}_p$*, Russ. J. Math. Phys., **16(4)**, (2009), 484-491.

[22] T. Kim, *Symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on $\mathbb{Z}_p$*, Russ. J. Math. Phys., **16(1)** (2009), 93-96.

[23] T. Kim, *Degenerate Changhee numbers and polynomials of the second kind*, arXiv:1707.09721v1 [math.NT] 31 Jul 2017.

[24] T. Kim, *A note on degenerate Stirling numbers of the second kind*, Proc. Jangjeon Math. Soc.. **20(3)**(2017), 319-331.

[25] S. K. Sharma, W. A. Khan, S. Araci, S. S. Ahmed, *New type of degenerate Daehee polynomials of the second kind*, Adv. Differ. Equ. **428**(2020), 1-14.

[26] S. K. Sharma, W. A. Khan, S. Araci, S. S. Ahmed, *New construction of type 2 of degenerate central Fubini polynomials with their certain properties*, Adv. Differ. Equ. **587**(2020), 1-11.

Waseem Ahmad Khan,
Department of Electrical Engineering,
Prince Mohammad Bin Fahd University,
P.O Box 1664, Al Khobar 31952
Saudi Arabia.
E-mail address: wkhan1@pmu.edu.sa