

Minimum Vertex Degree Index for Graphs

Shashirekha S.¹, Smitha M.^{*1}, Sujatha R.¹, Ravi K. M.² and Veena C. R.³

ABSTRACT: In this research work, we introduce the notion of new topological index called Minimum Vertex Degree Index (*MVDI*), for a graph G and obtain *MVDI* for few graphs. Also, we obtain *MVDI* for few chemical structures and develop a report comprising data analysis for properties of lower alkanes.

Key Words: Minimum vertex degree index, elliptic delta and modified elliptic delta.

Contents

1	Introduction	1
2	Definition and Computation of Minimum Vertex Degree Index of Some Special and Standard Graphs	2
3	Computation of Minimum Vertex Degree Index for Chemical Structure of Certain Networks	3
3.1	Minimum Vertex Degree Index for Silicate Network:	3
3.2	Minimum Vertex Degree Index for Honeycomb Network	4
3.3	Minimum Vertex Degree Index for Oxide Network:	4
3.4	Minimum Vertex Degree Index for Hexagonal Network:	5
4	A QSPR Analysis for MVDI	6

1. Introduction

This entire manuscript studies graphs, that are connected, simple, undirected and finite, which are given in [7] by F.Harary. Here vertex set is represented by $V(G)$ and edge set by $E(G)$ for a graph G with n vertices. We represent degree of vertex v_i as $d(v_i)$. The graph energy and the topological index for graphs are very attractive due their importance in mathematical chemistry. In the research field of drug development, the task of topological indices is well defined. In the process of drug invention technology, a sequence of definitions for topological indices are discussed. There exists many distance based and degree based topological indices defined for graphs.

A graphical structure attains numerical value through a topological index. A huge variety of graph indices are introduced and their applications in various disciplines are studied in [1,2,3,4,5,6,8,9,10,11,12, 13,14,15,16,17,20,21]. These indices are graph invariant and are used for QSPR (Quantitative Structure Property Relationship) and QSAR (Quantitative Structure Activity Relationship).

We consider the following definitions from papers of V. R. Kulli [18,19], in the process of developing the paper.

The δ vertex degree at vertex u is defined as

$$\delta_u = d_u - \delta(G) + 1$$

Here $\delta(G)$ represents minimum degree among the vertices of G .

Definition 1.1 The elliptic delta index of the graph G is defined as,

$$E\delta(G) = \sum_{uv \in E(G)} (\delta_u + \delta_v) \sqrt{\delta_u^2 + \delta_v^2}$$

* Corresponding author.

2010 Mathematics Subject Classification: 05C35, 05C50.

Submitted October 10, 2025. Published December 19, 2025

Definition 1.2 The modified elliptic delta index of the graph G is defined as,

$${}^m E\delta(G) = \sum_{uv \in E(G)} \frac{1}{(\delta_u + \delta_v)\sqrt{\delta_u^2 + \delta_v^2}}$$

Motivated by the above works, we introduce a new topological index called Minimum Vertex Degree Index ($MVDI$), in section 2 and compute ($MVDI$) for few special and standard graphs. In section 3 we compute $MVDI$ for chemical structure of certain networks. In section 4 we give QSPR analysis for physical properties of lower alkanes and also give regression models for the analysis.


2. Definition and Computation of Minimum Vertex Degree Index of Some Special and Standard Graphs

Let G be a simple graph with $\{v_1, v_2, \dots, v_n\}$ as vertex set and $d(u_i)$ and $d(v_i)$ denote the degree of u_i and v_i respectively for, $i = 1, 2, 3, \dots, n$. Then, we define $MVDI$ as follows,

$$MVDI(G) = \sum_{u_i v_i \in E} \min[d(u_i), d(v_i)] \quad (2.1)$$

Using the above definition, we have the following computations:

Example 1:



The $MVDI$ for above Peterson graph is 30.

Now we give $MVDI$ for few standard graphs, 1. For a path P_n with n vertices we have,

$$MVDI(P_n) = 2n - 4.$$

2. For a cycle C_n with n vertices we have,

$$MVDI(C_n) = 2|E| = 2n.$$

3. For a complete graph K_n with n vertices we have,

$$MVDI(K_n) = \frac{n(n-1)^2}{2}.$$

4. For a fan graph F_n with n vertices we have,

$$MVDI(F_n) = 6(n - 1) - 1.$$

5. For a Star graph S_n with n vertices we have,

$$MVDI(S_n) = (n - 1)^2.$$

6. For a Double Star graph $S_n \times S_n$ with n vertices we have,

$$MVDI(S_n \times S_n) = (n - 2) + 4.$$

3. Computation of Minimum Vertex Degree Index for Chemical Structure of Certain Networks

3.1. Minimum Vertex Degree Index for Silicate Network:

Among the minerals found in nature silicates are accepted as complicated, largest and interesting compound. They are tetrahedron denoted by SiO_4 . When we represent it in two-dimensional plane, we see it as a ring formed by tetrahedrons, which are joined by oxygen nodes and we notice it as a sheetlike structure.

When we represent silicate network (SL_n) as a graph G , the number of silicon nodes equals the product of number of nodes of SL_n , also the number of edges in G is $6 \times 6n^2 = 36n^2$. Based on the degree at the vertex, it has the following three types of edges,

$$E_{3,3} = \{u_i v_i \in E(G) \mid d(u_i) = d(v_i) = 3\}, |E_{3,3}| = 6n.$$

$$E_{3,6} = \{u_i v_i \in E(G) \mid d(u_i) = 3, d(v_i) = 6\}, |E_{3,6}| = 18n^2 + 6n.$$

$$E_{6,6} = \{u_i v_i \in E(G) \mid d(u_i) = 6, d(v_i) = 6\}, |E_{6,6}| = 18n^2 - 12n.$$

Theorem 3.1 *Let G be the graphical representation of a Silicate Network SL_n . Then,*

$$MVDI(G) = 162n^2 - 36n.$$

Proof: *By definition we have,*

$$\begin{aligned} MVDI(G) &= \sum_{uv \in E} \min[d(u_i), d(v_i)] \\ &= 162n^2 - 36n. \end{aligned} \tag{3.1}$$

□

Also we have the following bounds for G which represents SL_n ,

(i) $MVDI(G) < E\delta(G)$

i.e, $162n^2 - 36n < (90\sqrt{17} + 576\sqrt{2})n^2 + (30\sqrt{17} - 372\sqrt{2})n$.

(ii) $MVDI(G) >^m E\delta(G)$

i.e, $162n^2 - 36n > \frac{6n}{2\sqrt{2}} + \frac{18n^2+6n}{5\sqrt{17}} + \frac{18n^2-12n}{32\sqrt{2}}$.

3.2. Minimum Vertex Degree Index for Honeycomb Network

A honeycomb network is a regular network (HC_n) with a hexagonal topology that can be constructed in any dimension. Honeycomb networks are often used to study complex systems and their properties. The applications of Honeycomb networks include Material science, Physics, Computer science, and Communication network design. In [19] V.R. Kulli obtained that the graphical structure of HC_n has $6n^2$ vertices and $9n^2 - 3n$ edges. Based on the degree at the vertex, it has the following three types of edges,

$$\begin{aligned} E_{2,2} &= \{u_i v_i \in E(G) \mid d(u_i) = d(v_i) = 2\}, |E_{2,2}| = 6. \\ E_{2,3} &= \{u_i v_i \in E(G) \mid d(u_i) = 2, d(v_i) = 3\}, |E_{2,3}| = 12n - 12. \\ E_{3,3} &= \{u_i v_i \in E(G) \mid d(u_i) = d(v_i) = 3\}, |E_{3,3}| = 9n^2 - 15n + 6. \end{aligned}$$

Theorem 3.2 *Let G be the graphical representation of a honeycomb network HC_n . Then,*

$$MVDI(G) = 27n^2 - 21n + 6.$$

Proof: By definition we have,

$$\begin{aligned} MVDI(G) &= \sum_{u_i v_i \in E} \min[d(u_i), d(v_i)] \\ &= 27n^2 - 21n + 6. \end{aligned} \tag{3.2}$$

□

Also we have the following bounds for G which represents HC_n ,

(i) $MVDI(G) < E\delta(G)$

i.e, $27n^2 - 21n + 6 < 72\sqrt{2}n^2 + (36\sqrt{5} - 120\sqrt{2})n + 60\sqrt{2} - 36\sqrt{5}$.

(ii) $MVDI(G) >^m E\delta(G)$

i.e, $27n^2 - 21n + 6 > \frac{6}{2\sqrt{2}} + \frac{12n-12}{3\sqrt{5}} + \frac{9n^2-15n+6}{8\sqrt{2}}$.

3.3. Minimum Vertex Degree Index for Oxide Network:

An oxide network (OX_n) is a network that is created by removing all the silicon nodes in a silicate network, hence the diameter of both SL_n and OX_n are same. An n -dimensional oxide network is represented by OX_n . Oxide networks are important for studying the metric and structural properties of silicate networks.

Let G be the graph of an oxide network OX_n . The graph G has $9n^2 + 3n$ vertices and $18n^2$ edges. Based on the degree at the vertex, it has the following two types of edges,

$$E_{2,4} = \{u_i v_i \in E(G) \mid d_G(u_i) = 2, d_G(v_i) = 4\}, |E_{2,4}| = 12n.$$

$$E_{4,4} = \{u_i v_i \in E(G) \mid d_G(u_i) = d_G(v_i) = 4\}, |E_{4,4}| = 18n^2 - 12n.$$

Theorem 3.3 Let G be the graphical representation of an oxide network OX_n . Then,

$$MVDI(G) = 72n^2 - 24n.$$

Proof: By definition we have,

$$\begin{aligned} MVDI(G) &= \sum_{u_i v_i \in E} \min[d(u_i), d(v_i)] \\ &= 72n^2 - 24n. \end{aligned} \tag{3.3}$$

□

Also we have the following bounds for G which represents OX_n ,

(i) $MVDI(G) < E\delta(G)$

$$\text{i.e., } 72n^2 - 24n < 162\sqrt{2}n^2 + (24\sqrt{10} - 54\sqrt{2})n.$$

(ii) $MVDI(G) >^m E\delta(G)$

$$\text{i.e., } 72n^2 - 24n > \frac{6n}{4\sqrt{10}} + \frac{9n^2 - 3n}{18\sqrt{2}}.$$

3.4. Minimum Vertex Degree Index for Hexagonal Network:

In graph theory, a hexagonal network is a network that is modeled by a planar graph and is based on a triangular plane tessellation. This means that the network is made up of equilateral triangles, with each node having up to six neighbours. In a hexagonal network the total count of number of edges and vertices are $(3n(n - 1) + 1)$ and $(9n - 15)n + 6$ respectively. Based on the degree at the vertex, it has the following five types of edges,

$$E_{3,4} = \{u_i v_i \in E(G) \mid d(u_i) = 3, d(v_i) = 4\}, |E_{3,4}| = 12.$$

$$E_{3,6} = \{u_i v_i \in E(G) \mid d(u_i) = 3, d(v_i) = 6\}, |E_{3,6}| = 6.$$

$$E_{4,4} = \{u_i v_i \in E(G) \mid d(u_i) = d(v_i) = 4\}, |E_{4,4}| = 16n - 18.$$

$$E_{4,6} = \{u_i v_i \in E(G) \mid d(u_i) = 4, d(v_i) = 6\}, |E_{4,6}| = 12n - 24.$$

$$E_{6,6} = \{u_i v_i \in E(G) \mid d(u_i) = d(v_i) = 6\}, |E_{6,6}| = 9n^2 - 33n + 30.$$

Theorem 3.4 Let G be the graphical representation of a Hexagonal Network HX_n . Then,

$$MVDI(G) = 54n^2 - 126n + 66.$$

Proof: By definition we have,

$$\begin{aligned} MVDI(G) &= \sum_{u_i v_i \in E} \min[d(u_i), d(v_i)] \\ &= 54n^2 - 126n + 66. \end{aligned} \tag{3.4}$$

□

Also we have the following bounds for G which represents HX_n ,

(i) $MVDI(G) < E\delta(G)$

$$\text{i.e., } 54n^2 - 126n + 66 < 288\sqrt{2}n^2 + (144\sqrt{5} - 1008\sqrt{2})n + 30\sqrt{17} - 252\sqrt{5} + 816\sqrt{2}.$$

(ii) $MVDI(G) >^m E\delta(G)$

$$\text{i.e., } 54n^2 - 126n + 66 > \frac{12}{3\sqrt{5}} + \frac{6}{5\sqrt{17}} + \frac{6n-18}{8\sqrt{2}} + \frac{12n-24}{12\sqrt{5}} + \frac{9n^2-33n+30}{32\sqrt{2}}.$$

4. A QSPR Analysis for MVDI

In this section, we carry out QSPR analysis of molecular structures of graphs using MVDI for properties of lower alkanes. The MVDI for molecular structures of graphs of lower alkanes are obtained further they are tabulated in the Table 1. For the analysis, we used Practical values of the properties - surface tensions (st) dyne cm^{-1} , critical temperatures (ct) C , molar refractions (mr) cm^3 , critical pressures (cp) atm, heats of vaporization (hv) kJ molar volumes (mv) cm^3 , and boiling points (bp) C for lower alkanes, presented in [17].

Following Table 1 shows the corelation between the properties of lower alkanes and MVDI. The corelation coefficient r is a measure of how strongly a pair of variables are related. It usually varies among -1 and +1. If r is nearer to +1 then we say that variables are set to be strongly, positively and linearly proportional. If r is nearer to -1 then we say that variables are set to be strongly, negatively and linearly inversely proportional. If r close to zero or r is equal to zero, then we conclude that there is no relationship between the variables. (we observe from Table 1 that properties of lower alkanes are positively proportional and is closer to the value+1)

Table 1: Minimum Vertex Degree Index of low alkanes

Alkane	MVDI	Alkane	MVDI
2,3,4-Trimethylpentane	11	Ethane	1
Nonane	14	Propane	4
2-Methyloctane	13	Butane	4
3-Methyloctane	13	2-Methylpropane	3
4-Methyloctane	13	Pentane	6
3-Ethylheptane	13	2-Methylbutane	5
4-Ethylheptane	13	2,2-Dimethylpropane	4
2,2-Dimethylheptane	12	Hexane	8
2,3-Dimethylheptane	13	2-Methylpentane	7
2,3,4-Trimethylpentane	11	Ethane	1
Nonane	14	Propane	4
2-Methyloctane	13	Butane	4
3-Methyloctane	13	2-Methylpropane	3
4-Methyloctane	13	Pentane	6
3-Ethylheptane	13	2-Methylbutane	5
4-Ethylheptane	13	2,2-Dimethylpropane	4
2,2-Dimethylheptane	12	Hexane	8
2,3-Dimethylheptane	13	2-Methylpentane	7
2,4-Dimethylheptane	12	3-Methylpentane	7

Alkane	MVDI	Alkane	MVDI
2,5-Dimethylheptane	12	2,2-Dimethylbutane	6
2,6-Dimethylheptane	12	2,3-Dimethylbutane	7
3,3-Dimethylheptane	12	Heptane	9
3,4-Dimethylheptane	13	2-Methylhexane	9
3,5-Dimethylheptane	12	3-Methylhexane	9
4,4-Dimethylheptane	13	3-Ethylhexane	11
3-Ethyl-2-methylhexane	13	2,2-Dimethylpentane	8
4-Ethyl-2-methylhexane	12	2,3-Dimethylpentane	9
3-Ethyl-3-methylhexane	12	2,4-Dimethylpentane	8
3-Ethyl-4-methylhexane	13	3,3-Dimethylpentane	8
2,2,3-Trimethylhexane	12	2,3,3-Trimethylbutane	8
2,2,4-Trimethylhexane	11	Octane	12
2,2,5-Trimethylhexane	11	2-Methylheptane	11
2,3,3-Trimethylhexane	12	3-Methylheptane	11
2,3,4-Trimethylhexane	13	4-Methylheptane	11
2,3,5-Trimethylpentane	11	4-Ethylhexane	11
2,4,4-Trimethylhexane	11	2,2-Dimethylhexane	10
3,3,4-Trimethylhexane	12	2,3-Dimethylhexane	11
3,3,4-Trimethylhexane	12	2,4-Dimethylhexane	10
3,3-Diethylpentane	12	2,5-Dimethylhexane	10
2,2-Dimethyl-3-ethylpentane	12	3,3-Dimethylhexane	10
2,3-Dimethyl-3-ethylpentane	12	3-Ethyl-2-methylpentane	11
2,4-Dimethyl-3-ethylpentane	13	3,4-Dimethylhexane	11
2,2,3,3-Tetramethylpentane	12	3-Ethyl-3-methylpentane	10
2,2,3,4-Tetramethylpentane	12	2,2,3-Trimethylpentane	10
2,2,4,4-Tetramethylpentane	10	2,2,4-Trimethylpentane	9
2,3,3,4-Tetramethylpentane	12	2,3,3-Trimethylpentane	10

Table 2: r , r^2 and p -value between MVDI and physical properties

Y	r	r^2	value	Std error
B_p	0.9525	0.9073	$2.5361e - 36$	0.5355
M_v	0.8908	0.7935	$1.2124e - 24$	0.4413
M_r	0.9131	0.8337	$8.3913e - 28$	0.1205
H_v	0.9572	0.9163	$8.2717e - 38$	0.0872
C_t	0.9271	0.8595	$2.9028e - 30$	0.8012
C_p	-0.8167	0.6671	$1.1765e - 17$	0.0850
St	0.9023	0.8141	$8.3752e - 26$	0.0504

Regression models The linear model for the properties of the lower alkanes with $MVDI$ is provided by the following equations.

$$y_{B_p} = 13.7131MVDI - 30.8657 \quad (4.1)$$

$$y_{M_v} = 7.0810MVDI + 87.5205 \quad (4.2)$$

$$y_{M_r} = 2.2085MVDI + 16.0566 \quad (4.3)$$




$$y_{H_v} = 2.3626MVDI + 13.7306 \quad (4.4)$$




$$y_{C_t} = 16.2226MVDI + 119.1618 \quad (4.5)$$


$$y_{C_p} = -0.9847MVDI + 36.7448 \quad (4.6)$$

$$y_{St} = 0.8571MVDI + 11.7596 \quad (4.7)$$

The variables are shown to coincide with one another in the following graphs. The coefficient of determination is high if the model's traces are closer to the real values. Therefore, there is no need to improve the current state of the model.

Figure 1: Regression model for *bp*Figure 2: Regression model for *mv*Figure 3: Regression model for *mr*

Figure 4: Regression model for $h\nu$ Figure 5: Regression model for ct Figure 6: Regression model for cp

Figure 7: Regression model for *st*

Conclusion:

Topological indices serve as an important predictive tool for the chemical and physical properties of compounds. The various extended versions of topological indices have become the most interesting aspect of the research due to their extensive applications in the field of chemical sciences. Motivated by the applications, we introduced *MVDI* and obtained a numerical value of *MVDI* for some standard and special graphs in terms of their vertices and constants. Further, we obtained *MVDI* for hexagonal, oxide, silicate and honeycomb networks. We also compared the values of *MVDI* with the modified elliptic delta index and the elliptic delta index. A QSPR analysis of molecular structures of graphs using *MVDI* for properties of lower alkanes is done, which shows that, *MVDI* is strongly correlated with physical properties of lower alkanes.

Conflicts of interest : The authors declare no conflict of interest.

References

1. A. Ashraf and S. Akhter, Revan indices and Revan polynomials of silicon carbide graphs, International Journal of Engineering Research & Technology, 8(9) (2019), 353-361.
2. A. Q. Baig, M. Naeem and W. Gao, Revan and Hyper-Revan indices of octahedral and icosahedral networks, Applied Mathematics and Nonlinear Sciences, 3(1) (2018), 33-40.
3. A. Mahboob, G. Muhiuddin, I. Siddique and S. M. Alam, A view of Banhatti and Revan indices in chemical graphs, Journal of Mathematics, 2022(8) (2022).
4. B. Basavanagoud and S. Policepatil, Chemical applicability of Gourava and hyper Gourava indices, Nano Systems: Physics, Chemistry, Mathematics, 12(2) (2021), 142-150.
5. D. Narasimhan, R. Vignesh and K. Desikan, Results on Revan and hyper Re van indices of some HEX derived networks, Fuzzy Mathematical Analysis and Advances in Computational Mathematics, (2022), 209-220.
6. F. Dayan, M. Javaid and M. A. Rehman, On leap Gourava indices of some wheel related graphs, Scientific Inquiry and Review, 2(4) (2018), 14-24.
7. F. Harary, Graph Theory, Addison Wesley, Reading, 1972.
8. G. N. Adithya, N. D. Soner and M. Kirankumar, Gourava indices for Jahangir graph and phase transfer catalyst, Journal of Emerging Technologies and Innovative Research, 10(6) (2023), 394-399.
9. I. Gutman and N. Trinajstic, Graph Theory and Molecular Orbitals. Total phi electron energy of alternant hydrocarbons, Chemical Physics Letters, 17(4) (1972), 535-538.
10. I. Gutman, V. R. Kulli, B. Chaluvaraju and H. S. Boregowda, On Banhatti and Zagreb indices, Journal of the International Mathematical Virtual Institute, 7(2017), 53-67.
11. K. G. Mirajkar and B. Pooja, On Gourav indices of some chemical graphs, International Journal of Applied Engineering Research, 14(3) (2019), 743-749.

12. M. Aruvi, J. M. Joseph and E. Ramganesh, The second Gourava index of some graph products, *Advances in Mathematics: Scientific Journal*, 9(12) (2020), 10241-10249.
13. M. Kamran, N. Salamat, R. H. Khan, M. A. Ullah, M. S. Hameed and M. K. Pandit, Computation of Revan topological indices for phenol-formaldehyde resin, *Journal of Mathematics*, (2022), 10 pages.
14. M.V. Diudea (ed.) *QSPR/QSAR Studies by Molecular Descriptors*, NOVA New York, (2001).
15. R. Aguilar-Sanchez, I. F. Herrera-Gonzalez, J. A. Mendez-Bermudez and J. M. Sigaareta, Revan degree indices on random graphs, arXiv: 2210.04749v1 [math. CO], (2022).
16. R. Rajendra, P. Siva Kota Reddy and M. Prabhavathi, Computation of Wiener index, Reciprocal Wiener index and Peripheral Wiener index using adjacency matrix, *Southeast Asian J. of Mathematics and Mathematical Sciences*, Vol. 18, No 3(2022), pp. 275-282.
17. S. Wagner and H. Wang, *Introduction to Chemical Graph Theory*, Boca Raton, CRC Press, (2018).
18. V. R. Kulli, δ -Sombor index and its exponential of certain nanotubes, *Annals of Pure and Applied Mathematics*, 23(1) (2021), 37-42.
19. V. R. Kulli, Elliptic Delta and Modified Elliptic Delta Indices of Certain Networks, *International Journal of Mathematics and its Applications*, ISSN: 2347-1557, 12(2) (2024), 105-116.
20. V. R. Kulli, Graph indices, in *Handbook of Research on Advanced Applications of Application Graph Theory in Modern Society*, M. Pal, S. Samanta & A. Pal (eds), IGI Global, USA, (2019), 66-91.
21. V. R. Kulli, B. Chaluvaraju and H. S. Boregowda, K Banhatti and K hyper Banhatti indices of windmill graphs, *South East Asian J. of Math. Math. Sci.*, 13(1) (2017), 11-18.

¹ Department of Mathematics, JSS Science and Technology University, Mysuru-57006, Karnataka, INDIA.

E-mail address: shashirekha.s157@gmail.com, smitham@jssstuniv.in, rsujatha@sjce.ac.in

and

²Department of Mathematics, R V College of Engineering, Bengaluru-560059, Karnataka, INDIA.

E-mail address: drravikm@rvce.edu.in

and

³Department of Mathematics, JSS College of Arts, Commerce and Science, Ooty Road, Mysuru-570 025, Karnataka, INDIA.

E-mail address: veenacr.maths@gmail.com