

Explicit properties of q -Cosine and q -Sine Fubini-type polynomials and numbers

Waseem Ahmad Khan and Manoj Sharma

ABSTRACT: In recent years, q -special polynomials, such as q -Bernoulli, q -Genocchi, q -Euler and q -Frobenius-Genocchi have been studied and investigated by many mathematicians, as well physicists. It is important that any polynomials have explicit formulas, symmetric identities, summation formulas, and relations with other polynomials. In this work, the q -Cosine and q -Sine Fubini type polynomials are introduced and multifarious above mentioned properties for these polynomials are derived by utilizing some series manipulation methods. Moreover, several correlations related to both the q -Bernoulli, q -Euler, and q -Genocchi polynomials and the q -Stirling numbers of the second kind.

Key Words: Quantum calculus, Fubini type polynomials, Parametric q -Fubini type polynomials, generating functions, Stirling numbers of the second kind.

Contents

1	Introduction	1
2	Bivariate kinds of q-Cosine and q-Sine Fubini type polynomials	4
3	Relationship between q-Bernoulli, q-Euler and q-Genocchi polynomials and q-Stirling numbers of the second kind	9
4	Conclusions	11

1. Introduction

Recently, many authors [1,2,3,5,6,35] have introduced and constructed generating functions for new families of special polynomials including two parametric kinds of polynomials as Bernoulli, Euler, Genocchi, etc. They have given fundamental properties of these polynomials. Also, they have established more identities, and relations among trigonometric functions, two parametric kinds of special polynomials by using generating functions. Special polynomials have important role in several subjects of mathematics, approximation theory, engineering and theoretical physics. Applying the partial derivative operator to these generating functions, some derivative formulae, and finite combinatorial sums involving the aforementioned polynomials and numbers are obtained. In addition, these special polynomials allow the derivation of different useful identities in a fairly straightforward way and help in introducing new families of special polynomials. The Fubini-kind polynomials seem in combinatorial mathematics and play an crucial role in the principle and applications of arithmetic, hence many wide variety idea and combinatorics experts have extensively studied their residences and received a series of exciting results (see [7,8,13,14,15,16,17]). By inspiring and motivating the above polynomials, in this study, we are purpose to define a parametric kinds of q -Fubini type polynomials by introducing the two specific q -exponential generating functions. Also, we show many formulation and family members for those polynomials, such as a few implicit summation formulas, differentiation policies, and correlations with the sooner polynomials with the aid of utilizing some collection manipulation approach.

The concern of q -calculus started performing in the 19th century due to its packages in various fields of mathematics, physics and engineering. The definitions and notations of q -calculus reviewed right here are taken from (see [4,18,19,20,21,22]):

2010 *Mathematics Subject Classification:* 11B68, 11B73, Secondary 05A15, 05A19.

Submitted October 15, 2025. Published December 19, 2025

The q -analogue of the shifted factorial $(\alpha)_\omega$ is given by

$$(\alpha; q)_0 = 1, (\alpha; q)_\omega = \prod_{\gamma=0}^{\omega-1} (1 - q^\gamma \alpha) \quad \omega \in \mathbb{N}.$$

The q -analogue of a complex number α and of the factorial function is given by

$$[\alpha]_q = \frac{1 - q^\alpha}{1 - q} \quad q \in \mathbb{C} - \{1\}; \alpha \in \mathbb{C},$$

$$[\omega]_q! = \prod_{\gamma=1}^{\omega} [\gamma]_q = [1]_q [2]_q \cdots [\omega]_q = \frac{(q; q)_\omega}{(1 - q)^\omega} \quad q \neq 1; \omega \in \mathbb{N},$$

$$[0]_q! = 1, q \in \mathbb{C}; 0 < q < 1.$$

The Gauss q -binomial coefficient $\binom{\omega}{k}_q$ is given by

$$\binom{\omega}{\gamma}_q = \frac{[\omega]_q!}{[\gamma]_q! [\omega - \gamma]_q!} = \frac{(q; q)_\omega}{(q; q)_\gamma (q; q)_{\omega-\gamma}} \quad \gamma = 0, 1, \dots, \omega.$$

The q -analogue of the function $(x + y)_q^\omega$ is given by

$$(x + y)_q^\omega = \sum_{\gamma=0}^{\omega} \binom{\omega}{\gamma}_q q^{\gamma(\gamma-1)/2} x^{\omega-\gamma} y^\gamma \quad \omega \in \mathbb{N}_0. \quad (1.1)$$

The q -analogue of exponential functions are given by [23,24,25,26,27]

$$e_q(x) = \sum_{\omega=0}^{\infty} \frac{x^\omega}{[\omega]_q!} = \frac{1}{((1-q)x; q)_\infty} \quad 0 < |q| < 1; |x| < |1-q|^{-1}, \quad (1.2)$$

and

$$E_q(x) = \sum_{\omega=0}^{\infty} \frac{q^{\binom{\omega}{2}}}{[\omega]_q!} x^\omega = (-(1-q)x; q)_\infty \quad 0 < |q| < 1; x \in \mathbb{C}. \quad (1.3)$$

These two functions are related by the equation (see [9,10])

$$e_q(x) E_q(-x) = 1.$$

The q -derivative operator is defined by

$$D_q f(z) = \frac{f(qz) - f(z)}{qz - z}, \quad 0 < |q| < 1,$$

and $D_q f(0) = f'(0)$ provided that f is differentiable at $x = 0$.

The q -derivative fulfills the following product and quotient rules

$$D_{q,z}(f(z)g(z)) = f(z)D_{q,z}g(z) + g(qz)D_{q,z}f(z), \quad (1.4)$$

and

$$D_{q,z} \left(\frac{f(z)}{g(z)} \right) = \frac{g(qz)D_{q,z}f(z) - f(qz)D_{q,z}g(z)}{g(z)g(qz)}. \quad (1.5)$$

Definition 1.1 The q -trigonometric functions are (see [11]):

$$\sin_q(x) = \frac{e_q(ix) - e_q(-ix)}{2i}, \quad \text{SIN}_q(x) = \frac{E_q(ix) - E_q(-ix)}{2i},$$

and

$$\cos_q(x) = \frac{e_q(ix) + e_q(-ix)}{2}, \quad \text{COS}_q(x) = \frac{E_q(ix) + E_q(-ix)}{2},$$

where $\text{SIN}_q(x) = \sin_{q^{-1}}(x)$, $\text{COS}_q(x) = \cos_{q^{-1}}(x)$.

The Apostol-type q -Bernoulli polynomials $\mathbb{B}_{\omega,q}^{(\alpha)}(x; \lambda)$ of order α , the Apostol-type q -Euler polynomials $\mathbb{E}_{\omega,q}^{(\alpha)}(x; \lambda)$ of order α and the Apostol-type q -Genocchi polynomials $\mathbb{G}_{\omega,q}^{(\alpha)}(x; \lambda)$ of order α are defined by (see [28, 29, 30, 33]):

$$\left(\frac{t}{\lambda e_q(t) - 1} \right)^\alpha e_q(xt) = \sum_{\omega=0}^{\infty} \mathbb{B}_{\omega,q}^{(\alpha)}(x; \lambda) \frac{t^\omega}{[\omega]_q!} \quad (|t + \log \lambda| < 2\pi), \quad (1.6)$$

$$\left(\frac{2}{\lambda e_q(t) + 1} \right)^\alpha e_q(xt) = \sum_{\omega=0}^{\infty} \mathbb{E}_{\omega,q}^{(\alpha)}(x; \lambda) \frac{t^\omega}{[\omega]_q!} \quad (|t + \log \lambda| < \pi), \quad (1.7)$$

and

$$\left(\frac{2t}{\lambda e_q(t) + 1} \right)^\alpha e_q(xt) = \sum_{\omega=0}^{\infty} \mathbb{G}_{\omega,q}^{(\alpha)}(x; \lambda) \frac{t^\omega}{[\omega]_q!} \quad (|t + \log \lambda| < \pi), \quad (1.8)$$

respectively.

Clearly, we have

$$\mathbb{B}_{\omega,q}^{(\alpha)}(\lambda) = \mathbb{B}_{\omega,q}^{(\alpha)}(0; \lambda), \quad \mathbb{E}_{\omega,q}^{(\alpha)}(\lambda) = \mathbb{E}_{\omega,q}^{(\alpha)}(0; \lambda), \quad \mathbb{G}_{\omega,q}^{(\alpha)}(\lambda) = \mathbb{G}_{\omega,q}^{(\alpha)}(0; \lambda).$$

Kang and Ryoo [12] introduced and defined the q -Bernoulli and q -Euler polynomials of complex variable as follows:

$$\frac{t}{e_q(t) - 1} e_q(xt) \text{COS}_q(yt) = \sum_{j=0}^{\infty} \frac{\mathbb{B}_{j,q}((x \oplus iy)_q) + \mathbb{B}_j((x \ominus iy)_q)}{2} \frac{t^j}{[j]_q!} = \sum_{j=0}^{\infty} \mathbb{B}_{j,q}^{(C)}(x, y) \frac{t^j}{[j]_q!}, \quad (1.9)$$

$$\frac{t}{e_q(t) - 1} e_q(xt) \text{SIN}_q(yt) = \sum_{j=0}^{\infty} \frac{\mathbb{B}_{j,q}((x \oplus iy)_q) - \mathbb{B}_{j,q}((x \ominus iy)_q)}{2i} \frac{t^j}{[j]_q!} = \sum_{j=0}^{\infty} \mathbb{B}_{j,q}^{(S)}(x, y) \frac{t^j}{[j]_q!}, \quad (1.10)$$

and

$$\frac{2}{e_q(t) + 1} e_q(xt) \text{COS}_q(yt) = \sum_{j=0}^{\infty} \frac{\mathbb{E}_{j,q}((x \oplus iy)_q) + \mathbb{E}_{j,q}((x \ominus iy)_q)}{2} \frac{t^j}{[j]_q!} = \sum_{j=0}^{\infty} \mathbb{E}_{j,q}^{(C)}(x, y) \frac{t^j}{[j]_q!}, \quad (1.11)$$

$$\frac{2}{e_q(t) + 1} e_q(xt) \text{SIN}_q(yt) = \sum_{j=0}^{\infty} \frac{\mathbb{E}_j((x \oplus iy)_q) - \mathbb{E}_j((x \ominus iy)_q)}{2i} \frac{t^j}{[j]_q!} = \sum_{j=0}^{\infty} \mathbb{E}_{j,q}^{(S)}(x, y) \frac{t^j}{[j]_q!}, \quad (1.12)$$

respectively.

Also they proved that (see [34]):

$$e_q(xt) \text{COS}_q(yt) = \sum_{r=0}^{\infty} C_{r,q}(x, y) \frac{t^r}{[r]_q!}, \quad (1.13)$$

and

$$e_q(xt) \text{SIN}_q(yt) = \sum_{r=0}^{\infty} S_{r,q}(x, y) \frac{t^r}{[r]_q!}, \quad (1.14)$$

where

$$C_{r,q}(x, y) = \sum_{j=0}^{\lfloor \frac{r}{2} \rfloor} (-1)^j \binom{r}{2j}{}_q (-1)^j q^{2j-1} x^{r-2j} y^{2j}, \quad (1.15)$$

and

$$S_{r,q}(x, y) = \sum_{j=0}^{\lfloor \frac{r-1}{2} \rfloor} \binom{r}{2j+1}{}_q (-1)^j q^{(2j+1)j} x^{r-2j-1} y^{2j+1}. \quad (1.16)$$

For $\lambda \in \mathbb{C}$, the generalized q -Stirling numbers of the second kind $S_{m,q}^\omega(\lambda)$ are given by (see [26]):

$$\frac{(\lambda e_q(t) - 1)^m}{m!} = \sum_{\omega=0}^{\infty} S_{m,q}^\omega(\lambda) \frac{t^\omega}{[\omega]_q!} \quad m \in \mathbb{N}_0 = \{0, 1, 2, \dots\}. \quad (1.17)$$

Taking $\lambda = 1$, equation (1.17) reduces to the q -Stirling numbers of the second kind as follows

$$\frac{(e_q(t) - 1)^m}{m!} = \sum_{\omega=m}^{\infty} S_{2,q}(\omega, m) \frac{t^\omega}{[\omega]_q!}. \quad (1.18)$$

The q -type Geometric polynomials of two variables are defined by (see [28]):

$$\frac{1}{1 - y(e_q(t) - 1)} e_q(xt) = \sum_{n=0}^{\infty} \mathbb{F}_{n,q}(x; y) \frac{t^n}{[n]_q!}. \quad (1.19)$$

When $x = 0$, $\mathbb{F}_{n,q}(y) = \mathbb{F}_{n,q}(0; y)$ are called the q -type Geometric polynomials and $\mathbb{F}_{n,q}(0; 1) = \mathbb{F}_{n,q}$ are called the q -type Geometric numbers.

The Fubini type polynomials of order α are defined by the following generating function [36]

$$\frac{2^\alpha}{(2 - e^t)^{2\alpha}} e^{xt} = \sum_{n=0}^{\infty} a_n^{(\alpha)}(x) \frac{t^n}{n!}, \quad \alpha \in \mathbb{N}_0; \quad |t| < \log(2). \quad (1.20)$$

When $x = 0$, $a_n^{(\alpha)} = a_n^{(\alpha)}(0)$ are called the Fubini type numbers of order α .

2. Bivariate kinds of q -Cosine and q -Sine Fubini type polynomials

In this section, we consider the q -Cosine and q -Sine Fubini type polynomials of complex variable and deduce some identities of these polynomials. First, we present the following definition as.

$$\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) E_q(ity) = \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(\alpha)}((x + iy)_q) \frac{t^n}{[n]_q!}. \quad (2.1)$$

On the other hand, we suppose that

$$e_q(xt) E_q(ity) = e_q(xt) (\text{COS}_q(yt) + i \text{SIN}_q(yt)). \quad (2.2)$$

Thus, by (2.1) and (2.2), we have

$$\sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(\alpha)}((x + iy)_q) \frac{t^n}{[n]_q!} = \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) E_q(ity) = \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) (\text{COS}_q(yt) + i \text{SIN}_q(yt)), \quad (2.3)$$

and

$$\sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(\alpha)}((x - iy)_q) \frac{t^n}{[n]_q!} = \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) E_q(-ity) = \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) (\text{COS}_q(yt) - i \text{SIN}_q(yt)). \quad (2.4)$$

From (2.3) and (2.4), we get

$$\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) COS_q(yt) = \sum_{n=0}^{\infty} \left(\frac{\mathcal{F}_{n,q}^{(\alpha)}((x \oplus iy)_q) + \mathcal{F}_{n,q}^{(\alpha)}((x \ominus iy)_q)}{2} \right) \frac{t^n}{[n]_q!}, \quad (2.5)$$

and

$$\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) SIN_q(yt) = \sum_{n=0}^{\infty} \left(\frac{\mathcal{F}_{n,q}^{(\alpha)}((x \oplus iy)_q) - \mathcal{F}_{n,q}^{(\alpha)}((x \ominus iy)_q)}{2} \right) \frac{t^n}{[n]_q!}. \quad (2.6)$$

Definition 2.1 Let $n \geq 0$. We define two parametric kinds of q -Cosine Fubini type polynomials $\mathcal{F}_{n,q}^{(C,\alpha)}(x, y)$ and q -Sine Fubini type polynomials $\mathcal{F}_{n,q}^{(S,\alpha)}(x, y)$, for non negative integer n are defined by

$$\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) COS_q(yt) = \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!}, \quad (2.7)$$

and

$$\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) SIN_q(yt) = \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(S,\alpha)}(x, y) \frac{t^n}{[n]_q!}, \quad (2.8)$$

respectively.

Note that

$$\mathcal{F}_{n,q}^{(C,\alpha)}(0, 0) = \mathcal{F}_{n,q}^{(\alpha)}, \quad \mathcal{F}_{n,q}^{(S,\alpha)}(0, 0) = 0. \quad (n \geq 0).$$

From (2.5)-(2.8), we have

$$\mathcal{F}_{n,q}^{(C,\alpha)}(x, y) = \frac{\mathcal{F}_{n,q}^{(\alpha)}((x \oplus iy)_q) + \mathcal{F}_{n,q}^{(\alpha)}((x \ominus iy)_q)}{2}, \quad (2.9)$$

and

$$\mathcal{F}_{n,q}^{(S)}(x, y) = \frac{\mathcal{F}_{n,q}^{(\alpha)}((x \oplus iy)_q) - \mathcal{F}_{n,q}^{(\alpha)}((x \ominus iy)_q)}{2i}. \quad (2.10)$$

Remark 2.1 For $x = 0$ in (2.7) and (2.8), we get new type of q -Cosine Fubini type polynomials $\mathcal{F}_{n,q}^{(C,\alpha)}(y)$ and q -Sine Fubini type polynomials $\mathcal{F}_{n,q}^{(S,\alpha)}(y)$ as

$$\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} COS_q(yt) = \sum_{j=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(y) \frac{t^n}{[n]_q!}, \quad (2.11)$$

and

$$\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} SIN_q(yt) = \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(S,\alpha)}(y) \frac{t^n}{[n]_q!}, \quad (2.12)$$

respectively.

It is clear that

$$\mathcal{F}_{n,q}^{(C,\alpha)}(0) = \mathcal{F}_{n,q}^{\alpha}, \quad \mathcal{F}_{n,q}^{(S,\alpha)}(0) = 0, \quad n \geq 0.$$

Now, we start some basic properties of these polynomials.

Theorem 2.1 Let n integer. Then

$$\mathcal{F}_{n,q}^{(C,\alpha)}(y) = \sum_{v=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n+v}{2v}_q (-1)^v q^{(2v-1)v} y^{2v} \mathcal{F}_{n-2v,q}^{(\alpha)}, \quad (2.13)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha)}(y) = \sum_{v=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n+v}{2v+1}_q (-1)^v q^{(2v+1)v} y^{2v+1} \mathcal{F}_{n-2v-1,q}^{(\alpha)}. \quad (2.14)$$

Proof: By (2.11) and (2.12), we can derive the following equations

$$\begin{aligned} \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(y) \frac{t^n}{[n]_q!} &= \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} CO S_q(yt) = \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(\alpha)} \frac{t^n}{[n]_q!} \sum_{v=0}^{\infty} (-1)^v q^{(2v-1)v} \eta^{2v} \frac{t^v}{[2v]_q!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{v=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n+v}{2v}_q (-1)^v q^{(2v-1)v} \eta^{2v} \mathcal{F}_{n-2v,q}^{(\alpha)} \right) \frac{t^n}{[n]_q!}, \end{aligned} \quad (2.15)$$

and

$$\begin{aligned} \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(S,\alpha)}(y) \frac{t^n}{[n]_q!} &= \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} SIN_q(yt) \\ &= \sum_{n=0}^{\infty} \left(\sum_{v=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{2v+1}_q (-1)^v q^{(2v+1)v} y^{2v+1} \mathcal{F}_{n-2v-1,q}^{(\alpha)} \right) \frac{t^n}{[n]_q!}. \end{aligned} \quad (2.16)$$

Therefore, by (2.15) and (2.16), we get (2.13) and (2.14). \square

Theorem 2.2 *Let n integer. Then*

$$\mathcal{F}_{n,q}^{(\alpha)}((x \oplus iy)_q) = \sum_{k=0}^n \binom{n}{k}_q (x \oplus iy)_q^k \mathcal{F}_{n-k,q} = \sum_{k=0}^n \binom{n}{k}_q (iy)_q^k \mathcal{F}_{n-k,q}^{(\alpha)}(x), \quad (2.17)$$

and

$$\mathcal{F}_{n,q}^{(\alpha)}((x \ominus iy)_q; z) = \sum_{k=0}^n \binom{n}{k}_q (x \ominus iy)_q^k \mathcal{F}_{n-k,q} = \sum_{k=0}^n \binom{n}{k}_q (-1)^k (iy)_q^k \mathcal{F}_{n,q}^{(\alpha)}(x). \quad (2.18)$$

Proof: By using (2.3) and (2.4), we obtain (2.17) and (2.18). So we omit the proof. \square

Theorem 2.3 *Let n integer. Then*

$$\mathcal{F}_{n,q}^{(C,\alpha)}(x, y) = \sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(\alpha)} \mathcal{C}_{n-k,q}(x, y), \quad (2.19)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha)}(x, y) = \sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(\alpha)} \mathcal{S}_{n-k,q}(x, y). \quad (2.20)$$

Proof: Consider

$$\left(\sum_{n=0}^{\infty} a_n \frac{t^n}{n!} \right) \left(\sum_{k=0}^{\infty} b_k \frac{t^k}{k!} \right) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^j a_{n-k} b_k \right) \frac{t^n}{n!}.$$

Now

$$\begin{aligned} \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!} &= \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) CO S_q(yt) \\ &= \left(\sum_{k=0}^{\infty} \mathbb{F}_{k,q}^{(\alpha)} \frac{t^k}{[k]_q!} \right) \left(\sum_{n=0}^{\infty} \mathcal{C}_{n,q}(x, y) \frac{t^n}{[n]_q!} \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(\alpha)} \mathcal{C}_{n-k,q}(x, y) \right) \frac{t^n}{[n]_q!}, \end{aligned}$$

which proves (2.19). The proof of (2.20) is similar. \square

Theorem 2.4 Let $n \geq 0$. Then

$$\mathcal{F}_{n,q}^{(C,\alpha)}(x+r, y) = \sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(C,\alpha)}(x, y) r^{n-k}, \quad (2.21)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha)}(x+r, y) = \sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(S,\alpha)}(x, y) r^{n-k}. \quad (2.22)$$

Proof: By changing x with $x+r$ in (2.7), we have

$$\begin{aligned} \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x+r, y) \frac{t^n}{[n]_q!} &= \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q(xt) \text{COS}_q(yt) e^{rt} \\ &= \left(\sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!} \right) \left(\sum_{k=0}^{\infty} r^k \frac{t^k}{[k]_q!} \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(C,\alpha)}(x, y) r^{n-k} \right) \frac{t^n}{[n]_q!}, \end{aligned}$$

which complete the proof (2.21). The result (2.22) can be similarly proved. \square

Theorem 2.5 Let n integer. Then

$$\frac{\partial}{\partial x} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) = [n]_q \mathcal{F}_{n-1,q}^{(C,\alpha)}(x, y), \quad (2.23)$$

$$\frac{\partial}{\partial y} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) = -[n]_q \mathcal{F}_{n-1,q}^{(S,\alpha)}(x, qy), \quad (2.24)$$

and

$$\frac{\partial}{\partial x} \mathcal{F}_{n,q}^{(S,\alpha)}(x, y) = [n]_q \mathcal{F}_{n-1,q}^{(S,\alpha)}(x, y), \quad (2.25)$$

$$\frac{\partial}{\partial y} \mathcal{F}_{n,q}^{(S,\alpha)}(x, y) = [n]_q \mathcal{F}_{n-1,q}^{(C,\alpha)}(x, qy). \quad (2.26)$$

Proof: Equation (2.7) yields

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\partial}{\partial x} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!} &= \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} \frac{\partial}{\partial x} e_q(xt) \text{COS}_q(yt) = \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) \frac{t^{n+1}}{[n]_q!} \\ &= \sum_{n=1}^{\infty} \mathcal{F}_{n-1,q}^{(C,\alpha)}(x, y) \frac{t^n}{[(n-1)]_q!} = \sum_{n=1}^{\infty} [n]_q \mathcal{F}_{n-1,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!}, \end{aligned}$$

proving (2.23). Other (2.24), (2.25) and (2.26) can be similarly derived. \square

Theorem 2.6 Let $N \in \mathbb{N}^*$, the following formula holds true. Then

$$\mathcal{F}_{n,q}^{(C,\alpha)}(2x, y) = \sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(C,\alpha)}(x, y) x^{n-k}, \quad (2.27)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha)}(2x, y) = \sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(S,\alpha)}(x, y) x^{n-k}. \quad (2.28)$$

Proof: By using Definition 2.1, we can easily proof of equations (2.27) and (2.28). We omit the proof. \square

Theorem 2.7 For $n \geq 0$, we have

$$\mathcal{F}_{n,q}^{(C,\alpha)}((1 \oplus x)_q, y) = \sum_{r=0}^n \binom{n}{r}_q \mathcal{F}_{n-r,q}^{(C,\alpha)}(x, y), \quad (2.29)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha)}((1 \oplus x)_q, y) = \sum_{r=0}^n \binom{n}{r}_q \mathcal{F}_{n-r,q}^{(S,\alpha)}(x, y). \quad (2.30)$$

Proof: Using the generating function (2.7), we have

$$\begin{aligned} \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}((1 \oplus x)_q, y) \frac{t^n}{[n]_q!} - \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!} &= \left(\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} \right) (e_q(t) - 1) e_q(xt) \text{COS}_q(yt) \\ &= \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!} \left(\sum_{r=0}^{\infty} \frac{t^r}{[r]_q!} - 1 \right) \\ &= \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!} \sum_{r=0}^{\infty} \frac{t^r}{[r]_q!} - \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!} \\ &= \sum_{n=0}^{\infty} \sum_{r=0}^n \binom{n}{r}_q \mathcal{F}_{n-r,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!} - \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x, y) \frac{t^n}{[n]_q!}. \end{aligned}$$

Finally, equating the coefficients of the like powers of t on both sides, we get (2.29). The proof of (2.30) is similar. \square

Theorem 2.8 Let n be integer. Then

$$\mathcal{F}_{n,q}^{(C,\alpha+\beta)}(x, y) = \sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(\alpha)} \mathcal{F}_{n-k,q}^{(C,\beta)}(x, y) \quad (2.31)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha+\beta)}(x, y) = \sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(\alpha)} \mathcal{F}_{n-k,q}^{(S,\beta)}(x, y) \quad (2.32)$$

Proof: By (2.7), we get

$$\begin{aligned} \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha+\beta)}(x, y) \frac{t^n}{[n]_q!} &= \frac{2^{\alpha+\beta}}{(2 - e_q(t))^{2(\alpha+\beta)}} e_q(xt) \text{COS}_q(yt) \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \mathcal{F}_{k,q}^{(\alpha)} \mathcal{F}_{n-k,q}^{(C,\beta)}(x, y) \frac{t^k}{[k]_q!} \frac{t^n}{[n]_q!} = \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k}_q \mathcal{F}_{k,q}^{(\alpha)} \mathcal{F}_{n-k,q}^{(C,\beta)}(x, y) \frac{t^n}{[n]_q!}, \end{aligned}$$

which proves the result (2.31). The assertion (2.32) can be proved similarly. \square

3. Relationship between q -Bernoulli, q -Euler and q -Genocchi polynomials and q -Stirling numbers of the second kind

In this section, we prove some relationships for two bivariate kind of q -Cosine Fubini type polynomials and q -Sine Fubini type polynomials related to q -Bernoulli polynomials, q -Euler polynomials and q -Genocchi polynomials and q -Stirling numbers of the second kind. We start a following theorem.

Theorem 3.1 *Each of the following relationships holds true:*

$$\mathcal{F}_{n,q}^{(C,\alpha)}(x,y) = \sum_{s=0}^{n+1} \binom{n+1}{s}_q \left[\sum_{k=0}^s \binom{s}{k}_q \mathcal{B}_{s-k,q}(x) - \mathcal{B}_{s,q}(x) \right] \frac{\mathcal{F}_{n+1-s,q}^{(C,\alpha)}(y)}{[n+1]_q}, \quad (3.1)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha)}(x,y) = \sum_{s=0}^{n+1} \binom{n+1}{s}_q \left[\sum_{k=0}^s \binom{s}{k}_q \mathcal{B}_{s-k,q}(x) - \mathcal{B}_{s,q}(x) \right] \frac{\mathcal{F}_{n+1-s,q}^{(S,\alpha)}(y)}{[n+1]_q}. \quad (3.2)$$

Proof: By using (1.6) and (2.7), we have

$$\begin{aligned} \left(\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} \right) e_q(xt) \text{COS}_q(yt) &= \left(\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} \right) \frac{t}{e_q(t) - 1} \frac{e_q(t) - 1}{t} e_q(xt) \text{COS}_q(yt) \\ &= \frac{1}{t} \sum_{n=0}^{\infty} \left(\sum_{k=0}^s \binom{s}{k}_q \mathbb{B}_{s-k,q}(x) p^{\binom{k}{2}} \right) \frac{t^s}{[s]_q!} \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(y) \frac{t^n}{[n]_q!} \\ &\quad - \frac{1}{t} \sum_{s=0}^{\infty} \mathcal{B}_{s,q}(x) \frac{t^s}{[s]_q!} \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(y) \frac{t^n}{[n]_q!} \\ &= \frac{1}{t} \sum_{n=0}^{\infty} \left[\sum_{s=0}^n \binom{n}{s}_q \sum_{k=0}^s \binom{s}{k}_q \mathcal{B}_{s-k,q}(x) \right] \mathcal{F}_{n-s,q}^{(C,\alpha)}(y) \frac{t^n}{[n]_q!} \\ &\quad - \frac{1}{t} \sum_{n=0}^{\infty} \left[\sum_{s=0}^n \binom{n}{s}_q \mathcal{B}_{s,q}(x) \right] \mathcal{F}_{n-s,q}^{(C,\alpha)}(y) \frac{t^n}{[n]_q!}. \end{aligned}$$

By using Cauchy product and comparing the coefficients of $\frac{t^n}{[n]_q!}$, we arrive at the required result (3.1). The proof of (3.2) is similar. \square

Theorem 3.2 *Each of the following relationships holds true:*

$$\mathcal{F}_{n,q}^{(C,\alpha)}(x,y) = \sum_{s=0}^n \binom{n}{s}_q \left[\sum_{k=0}^s \binom{s}{k}_q \mathcal{E}_{s-k,q}(x) + \mathcal{E}_{s,q}(x) \right] \frac{\mathcal{F}_{n-s,q}^{(C,\alpha)}(y)}{[2]_q}, \quad (3.3)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha)}(x,y) = \sum_{s=0}^n \binom{n}{s}_q \left[\sum_{k=0}^s \binom{s}{k}_q \mathcal{E}_{s-k,q}(x) + \mathcal{E}_{s,q}(x) \right] \frac{\mathcal{F}_{n-s,q}^{(S,\alpha)}(y)}{[2]_q}. \quad (3.4)$$

Proof: By using definitions (1.7) and (2.7), we have

$$\begin{aligned} \left(\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} \right) e_q(xt) \text{COS}_q(yt) &= \left(\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} \right) \frac{[2]_q}{e_q(t) + 1} \frac{e_q(t) + 1}{[2]_q} e_q(xt) \text{COS}_q(yt) \\ &= \frac{1}{[2]_q} \left[\sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_q \mathcal{E}_{n-k,q}(x) \right) \frac{t^n}{[n]_q!} + \sum_{n=0}^{\infty} \mathcal{E}_{n,q}(x) \frac{t^n}{[n]_q!} \right] \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(y) \frac{t^n}{[n]_q!} \\ &= \frac{1}{[2]_q} \sum_{n=0}^{\infty} \left[\sum_{s=0}^n \binom{n}{s}_q \sum_{k=0}^s \binom{s}{k}_q \mathcal{E}_{s-k,q}(x) + \sum_{s=0}^n \binom{n}{s}_q \mathcal{E}_{s,q}(x) \right] \mathcal{F}_{n-s,q}^{(C,\alpha)}(y) \frac{t^n}{[n]_q!}. \end{aligned}$$

Comparing the coefficients of $\frac{t^n}{[n]_q!}$, we arrive at the desired result (3.3). The proof of (3.4) is similar. \square

Theorem 3.3 *Let n be integer. Then*

$$\mathcal{F}_{n,q}^{(C,\alpha)}(x, y) = \sum_{s=0}^n \binom{n+1}{s}_q \left[\sum_{k=0}^s \binom{s}{k}_q \mathcal{G}_{s-k,q}(x) + \mathcal{G}_{s,q}(x) \right] \frac{\mathcal{F}_{n+1-s,q}^{(C,\alpha)}(y)}{[2]_q[n+1]_q}, \quad (3.5)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha)}(x, y) = \sum_{s=0}^n \binom{n+1}{s}_q \left[\sum_{k=0}^s \binom{s}{k}_q \mathcal{G}_{s-k,q}(x) + \mathcal{G}_{s,q}(x) \right] \frac{\mathcal{F}_{n+1-s,q}^{(S,\alpha)}(y)}{[2]_q[n+1]_q}. \quad (3.6)$$

Proof: By (1.8) and (2.7), we have

$$\begin{aligned} \left(\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} \right) e_q(xt) \text{COS}_q(yt) &= \left(\frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} \right) e_q(xt) \text{COS}_q(yt) \frac{[2]_q t}{e_q(t) + 1} \frac{e_q(t) + 1}{[2]_q t} e_q(xt) \text{COS}_q(yt) \\ &= \frac{1}{[2]_q t} \left[\sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k}_q \mathcal{G}_{n-k,q}(x) p^{\binom{k}{2}} \right) \frac{t^n}{[n]_q!} + \sum_{n=0}^{\infty} \mathcal{G}_{n,q}(x) \frac{t^n}{[n]_q!} \right] \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(y) \frac{t^n}{[n]_q!} \\ &= \frac{1}{[2]_q} \sum_{n=0}^{\infty} \left[\sum_{s=0}^n \binom{n}{s}_q \sum_{k=0}^s \binom{s}{k}_q \mathcal{G}_{s-k,q}(x) + \sum_{s=0}^n \binom{n}{s}_q \mathcal{G}_{s,q}(x) \right] \mathcal{F}_{n+1-s,q}^{(C,\alpha)}(y) \frac{t^n}{[n+1]_q!}. \end{aligned}$$

Comparing the coefficients of $\frac{t^n}{[n]_q!}$, then we have the asserted result (3.5). The proof of (3.6) is similar. \square

Theorem 3.4 *Let n be integer. Then*

$$\mathcal{F}_{n,q}^{(C,\alpha)}(x+r, y) = \sum_{k=0}^n \binom{n}{k}_q \sum_{m=0}^k \mathcal{F}_{n-k,q}^{(C,\alpha)}(y) x^m S_{2,q}(k+r, m+r), \quad (3.7)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha)}(x+r, y) = \sum_{k=0}^n \sum_{m=0}^k \binom{n}{k}_q \mathcal{F}_{n-k,q}^{(S,\alpha)}(y) x^m S_{2,q}(k+r, m+r) \quad (3.8)$$

Proof: Using (1.18) and (2.7), we have

$$\begin{aligned} \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(x+r, y) \frac{t^n}{[n]_q!} &= \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q((r)t) (e_q(t) - 1 + 1)^x \text{COS}_q(yt) \\ &= \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q((r)t) \text{COS}_q(yt) \sum_{m=0}^{\infty} x^k (e_q(t) - 1)^m \frac{t^m}{[m]_q!} \\ &= \frac{2^\alpha}{(2 - e_q(t))^{2\alpha}} e_q((r)t) \text{COS}_q(yt) \sum_{k=0}^{\infty} \sum_{m=0}^k x^k S_{2,q}(k, m) \frac{t^k}{[k]_q!} \\ &= \sum_{n=0}^{\infty} \mathcal{F}_{n,q}^{(C,\alpha)}(y) \frac{t^n}{[n]_q!} \sum_{k=0}^{\infty} \sum_{m=0}^k x^k S_{2,q}(k, m) \frac{t^k}{[k]_q!} \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k}_q \sum_{m=0}^k \mathcal{F}_{n-k,q}^{(C,\alpha)}(y) x^m S_{2,q}(k+r, m+r) \frac{t^n}{[n]_q!}. \end{aligned}$$

Comparing the coefficients of $\frac{t^n}{[n]_q!}$, then we have the asserted result (3.7). The proof of (3.8) is similar. \square

Remark 3.1 Let n be integer. Then

$$\mathcal{F}_{n,q}^{(C,\alpha)}(x,y) = \sum_{k=0}^n \binom{n}{k} {}_q \sum_{m=0}^k \mathcal{F}_{n-k,q}^{(C,\alpha)}(y) x^m S_{2,q}(k,m), \quad (3.9)$$

and

$$\mathcal{F}_{n,q}^{(S,\alpha)}(x,y) = \sum_{k=0}^n \sum_{m=0}^k \binom{n}{k} {}_q \mathcal{F}_{n-k,q}^{(S,\alpha)}(y) x^m S_{2,q}(k,m) \quad (3.10)$$

4. Conclusions

Utilizing q -numbers and q -concepts, Mahmudov [31,32] considered q -Genocchi polynomials and numbers, q -Bernoulli polynomials and numbers and q -Euler polynomials and numbers and provided many properties and formulas for these polynomials. Inspired and motivated by this consideration, many authors have introduced q -special numbers and polynomials and have described their several identities and properties. In this paper, using the q -Cosine polynomials and q -Sine polynomials, we have introduced novel kinds of q -extensions of Fubini polynomials and have acquired multifarious properties and identities by making use of some series manipulation methods. Furthermore, we have computed the q -derivative operator rules for these polynomials. Moreover, we have determined the approximate root movements of the new mentioned polynomials in a complex plane, utilizing the Newton method and illustrating them in figures. The structure of the approximate roots will come out in various ways, depending on the condition of the variables, and new methods and theorems related to this topic need to be created and proven.

Not only can the ideas presented in this paper be utilized for similar polynomials, but these polynomials may also have possible applications in other scientific areas besides the applications described at the end of the paper. We would like to continue to study this line of research in the future.

References

1. N. Alam, W. A. Khan. C. Kizilates and C S Ryoo, Two variable q -General-Appell polynomials within the context of the monomiality principle. *Mathematics*, 13,765, 2025.
2. N. Ahmad and W. A. Khan. *Insights into new generalization of q -Legendre-based Appell polynomials: Properties and quasi monomiality*. *Mathematics*, 13,955, 2025.
3. N. Ahmad and W. A. Khan, A new generalization of of q -Laguerre-based Appell polynomials and Quasi monomiality. *Symmetry*, 17,439, 2025.
4. L. C. Andrews, *Special Functions for Engineers and Applied Mathematicians*, Macmillan Publishing Company, New York, 1985.
5. W. A. Khan, D. Srivastava and K. S. Nisar, A new class of generalized polynomials associated with Milne-Thomson-based poly-Bernoulli polynomials. *Miskolc Math. Notes*. 25(2):793-803, 2024.
6. W. A. Khan, J. Younis and M. Nadeem, Construction of partially degenerate Laguerre-Bernoulli polynomials of the first kind. *Applied Mathe. in Science and Engineering*. 30(1):362-375, 2022.
7. M. Zayed, W. A. Khan, C. S. Ryoo and U. Duran, An exploratory study on bivariate extended q -Laguerre-based Appell polynomials with some applications. *AIMS Mathematics*. 10(6):12841-12867, 2025.
8. M. S. Alatawi, W. A. Khan, C. S. Ryoo, Explicit properties of q -Cosine and q -Sine Array-type polynomials containing symmetric structures. *Symmetry*, **2022**, 14, 1675.
9. H. F. Jackson, q -Difference equations. *Am. J. Math.* **1910**, 32, 305-314.
10. H. F. Jackson, On q -functions and a certain difference operator. *Trans. R. Soc. Edinb.* **2013**, 46, 253-281.
11. J. Y. Kang, W. A. Khan, A new class of q -Hermite based Apostol type Frobenius Genocchi polynomials. *Communication of the Korean Mathematical Society*. **2020**, 35(3), 759-771.
12. J. Y. Kang, C. S. Ryoo, Various structures of the roots and explicit properties of q -cosine Bernoulli polynomials and q -sine Bernoulli polynomials. *Mathematics*, **2020**, 8, 463.
13. W. A. Khan, Some results on q -analogue type of Fubini numbers and polynomials. *Journal of Mathematical Control Science and Applications*, **2021**, 7(2), 141-154.

14. W. A. Khan, K. S. Mohamed, F.A. Costabile, C. Kizilates, C. S. Ryoo, A new generalization of q -truncated polynomials associated with q -General polynomials within the context of the monomiality principle. *Mathematics*, 2025, 13, 1964, 1-19.
15. W. A. Khan, K. S. Mohamed, F.A. Costabile, C. Kizilates, C. S. Ryoo, Finding the q -Appell convolution of certain polynomials within the context of quantum calculus. *Mathematics*, 2025, 13, 2073, 1-24.
16. H. Qawqneh, W. A. Khan, H. Aydi, U. Duran, C. S. Ryoo, A comprehensive study of generalized bivariate q -Laguerre polynomials: Structural properties and applications. *European Journal of Pure and Applied Mathematics*, 18 (3) (2025), 1-23.
17. S. A. Wani, M. Riyasat, R. William, W. A. Khan, Multivariate q -Hermite-based Appell polynomials: structural properties and applications. *Afrika Matematika* , (2025), 36:97.
18. A. Aledamat, W. A. Khan, U. Duran, S. A. K. Kirmani, C. S. Ryoo, A study on differential equations associated with (q, h) -Frobenius-Euler polynomials. *Journal of Mathematics and Computer Science*, 36(3) (2025), 386-398.
19. A. Alshejari, W. A. Khan, U. Duran, C. S. Ryoo, A study on differential equations associated with (q, h) -Frobenius-Genocchi polynomials. *Journal of Mathematics and Computer Science*, 37(3) (2025), 406-422.
20. W. A. Khan, M. Alhazmi, T. Nahid, A new family of q -Mittag-Leffler-based Bessel and Tricomi functions via umbral approach. *Symmetry*, (2024), 16:1580, 1-13.
21. W. A. Khan, Construction of degenerate q -Changhee polynomials with weight alpha and its applications. *Southeast Asian Journal of Mathematics and Mathematical Sciences*, 20(3) (2024), 65-80.
22. G. Fang, S. Koparal, N. Omur, O. Duran, W. A. Khan, On three dimensional q -Riordan arrays. *Demonstratio Mathematica*, 57(1) (2024), 1-12.
23. M. Fadel, M. S. Alatawi, W. A. Khan, Two variable q -Hermite based Appell polynomials and their applications. *Mathematics*, (2024), 12:1358, 1-17.
24. A. Aledamat, W. A. Khan, U. Duran, C. S. Ryoo, Construction for q -hypergeometric Bernoulli polynomials of a complex variable with applications computer modeling. *Advanced Mathematical Models and Applications*, 9(1) (2024), 68-92.
25. H. Guan, N. Omur, S. Koparal, W. A. Khan, q -analogue of $H_{n,m}$ and their applications. *Mathematics*, (2023), 15:943, 1-10.
26. Y. Rao, W. A. Khan, S. Araci, C. S. Ryoo, Explicit properties of Apostol-type Frobenius-Euler polynomials involving q -trigonometric function with applications in computer modeling. *Mathematics*, (2023), 11:2386, 1-21.
27. Z. Chen, N. Omur, S. Koparal, W. A. Khan, Some identities with multi generalized q -hyper harmonic numbers of order r. *Symmetry*, (2023), 15:917, 1-10.
28. W. A. Khan, M. S Alatawi, C. S. Ryoo, U. Duran, Novel properties of q -Sine based and q -Cosine-based q -Fubini polynomials. *Symmetry*, (2023), 15:356, 1-18.
29. W. A. Khan, A note on q -analogue of degenerate Catalan numbers associated p-adic integral on Z_p . *Symmetry Journal*. 14(119) (2022), 1-10. doi.org/10.3390/sym14061119.
30. W. A. Khan, A note on q -analogues of degenerate Catalan-Daehee numbers and polynomials. *Journal of Mathematics*, (2022), Article ID 9486880, 9 pages <https://doi.org/10.1155/2022/9486880>.
31. N. I. Mahmudov, q -analogues of the Bernoulli and Genocchi polynomials and the Srivastava-Pinter addition theorems. *Discrete and Dynamics in Nature and Soc.* Article number 169348, 2012. doi:10.1155/2012/169348.
32. N. I. Mahmudov, On a class of q -Bernoulli and q -Euler polynomials. *Adv. in Diff. Equa.* **2013**. doi:10.1186/1687-1847-2013-1.
33. K. S. Nisar, W. A. Khan, Notes on q -Hermite based unified Apostol type polynomials. *Journal of Interdisciplinary Mathematics*. **2019**, 22(7), 1185-1203.
34. C. S. Ryoo, J. Y. Kang, Explicit properties of q -Cosine and q -Sine Euler polynomials containing symmetric structures. *Symmetry*, **2020**, 12, 1247.
35. S. K. Sharma, W. A. Khan, C. S. Ryoo, A parametric kind of Fubini polynomials of a complex variable. *Mathematics*, **2020**, 8, 643; DOI: 10.3390/math8040643.
36. N. Kilar, Y. Simsek, A new family of Fubini type numbers and polynomials associated with Apostol type Bernoulli numbers and polynomials. *J. Korean Math. Soc.* 54(2017), 1605-1621.

*Waseem Ahmad Khan,
Department of Electrical Engineering,
Prince Mohammad Bin Fahd University,
P.O Box 1664, Al Khobar 31952
Saudi Arabia.*

E-mail address: `wkhan1@pmu.edu.sa`

and

Manoj Sharma

*Department of Mathematics,
Rustumji Institute of Technology,
BSF, Academy, Tekanpur, Gwalior, India.
E-mail address:* `manoj240674@yahoo.co.in`