



Even-Paired Domination with Modified Forgotten, Hyper Zagreb Indices of Graph

Pushpa N., Dhananjayamurthy B. V.

ABSTRACT: In this study, we define a new parameter for each vertex, referred to as the Even-paired domination degree, and denote it by $d_{coe}(v)$. Building on Even-paired domination degrees, we propose several Even-paired domination indices. We also determine the precise values of the Even-paired domination Forgotten, Refined Forgotten index/ Modified Forgotten invariant, Hyper-Zagreb graph invariant for certain families of well-known graphs. Additionally, we identify bounds, both lower and upper, for these indices across various graph structures.

Keywords: Topological Indices, Even-paired dominating sets, degree, Modified Forgotten, hyper Zagreb indices.

Contents

1 Introduction	1
2 Graph's Even-paired Domination Indices	2
3 Some bounds for Modified Forgotten, hyper Zagreb Indices of Graph and Even-paired Domination Forgotten	11

1. Introduction

In mathematics, theory of graph can be viewed as a new tool to deal, with sciences of all types and is applied in the disciplines of topological graphs [2,3], fuzzy graphs [4,5,6] etc. Mathematical chemistry is the most common application of graph theory in the last decade. Mathematical analysis of chemical phenomena can be done using graph theory tools. Topological indices help describe a graph's structure through numerical values that remain fixed for the entire graph. They're widely applied in building structure-activity relationships (QSAR), where they link chemical structures to biological activity and other traits-often via degree-based indices [7,8] or distance-based ones [9,10,11,13,14]. Note that the Zagreb index stands apart as a distance-based measure, defined as [15,17].

$$M_1(G) = \sum_{v \in V(G)} d(v)^2$$

$$M_2(G) = \sum_{e_1=uv \in E(G)} d(u)d(v).$$

The index has been overlooked in the references. For detailed insights into the Zagreb index and related topological indices, readers may consult the existing survey literature. In a simple graph, let Δ denote the maximum degree of a vertex and let δ denote the minimum degree. A Pendant-augmented n-wheel graph is obtained from a n-wheel graph with n vertices on its cycle by attaching a pendant edge to each vertex of the n-cycle [18]. Consider two path graphs, say P_m and P_n . Their Cartesian product, particularly when one of the paths is P_2 , results in a ladder graph with $(2n)$ vertices and $(3n - 2)$ edges. A dominating set D in a graph G is a set of vertices such that every vertex $v \in V(G)$ either belongs to D or is adjacent to some vertex $u \in D$. The domination number of G , denoted by $\gamma(G)$, is the minimum cardinality of such a set. A. M. Hanan Ahmed and her co-authors determined the domination number of the path graph P_n . For a general graph G , the domination number is denoted by $\gamma(G)$. However, the concept of co-equal domination is not applicable in this context. In this paper, we introduce a new parameter $\gamma_{coe}(G)$, called the *co-pair domination number* or *even-paired domination number* of a graph G . This

2020 *Mathematics Subject Classification:* 05C69, 05C90, 05C35.

Submitted January 21, 2026. Published February 26, 2026

parameter serves as a novel metric associated with the vertices of the graph. Along with this concept, we define several new co-pair domination indices based on classical competitive graph structures. We derive exact expressions for the co-pair Zagreb indices of important families of graphs. Furthermore, lower and upper bounds for the Zagreb index are established using control chart techniques.

2. Graph's Even-paired Domination Indices

The definition of Equivalent Dominance Forgotten, Modified Forgotten, and hyper Zagreb Indices was presented in this part, along with a formula for calculating the accuracy of these indices for specific lines in the Known class of graphs. Graph's even-paired Domination Indices

Definition 2.1 [1] *The Even-paired domination degree of a vertex $v \in V$ is*

$$d_{coe}(v) = |\{D \subseteq V : D \text{ is a even-paired dominating vertices, } v \in D\}|.$$

In other words,

$$d_{coe}(v) = \{D : D \text{ is a even-paired domination set, } v \in D \subseteq V\}$$

' $\delta_{coe}(G)$ ' and ' $\Delta_{coe}(G)$ ' denote the minimal and maximum even-paired domination degrees of G , respectively. where; $\delta_{coe}(G) = \min\{d_{coe}(v) : v \in V\}$, $\Delta_{coe}(G) = \max\{d_{coe}(v) : v \in V\}$

Definition 2.2 *Suppose G be a graph, and graph's even-paired domination set be D then the graph's Even-paired Domination Forgotten, Modified Forgotten, and Hyper Zagreb Indices are defined as :*

$$D_{coe}F(G) = \sum_{v \in V} d_{coe}^3(v). \quad (2.1)$$

$$D_{coe}F^*(G) = \sum_{e_1=uv \in E} \left[d_{coe}^2(u) + d_{coe}^2(v) \right] \quad (2.2)$$

$$D_{coe}HM_1(G) = \sum_{e_1=uv \in E} \left[d_{coe}(u) + d_{coe}(v) \right]^2. \quad (2.3)$$

Proposition 2.1 *If S_n is a star graph, then*

1. *If n is even*

$$D_{coe}F(S_n) = \sum_{v \in V} (1^2) = n.$$

$$D_{coe}F^*(S_n) = \sum_{e_1=uv \in E} (1^2 + 1^2) = 2n - 2.$$

$$D_{coe}HM(S_n) = \sum_{e_1=uv \in E} (1 + 1)^2 = 4n - 4.$$

2. *If n is odd;*

$$D_{coe}F(S_n) = 8n - 7.$$

$$D_{coe}F^*(S_n) = 5(n - 1).$$

$$D_{coe}HM(S_n) = 9(n - 1).$$

Proof: Let a star graph be isomorphic to a graph G . Assume the collection of n vertices of G is $\{v_1, v_2, \dots, v_{n-1}, w\}$, with w being the center vertex.

Case 1: There is only single even-paired dominating set if n is even, say $v_1, v_2, \dots, v_{n-1}, w$. Hence $d_{coe}(v) = 1, T_{coe}(v) = 1$, so we have only $\forall v \in V$, then by the definition 2.2

$$\begin{aligned} D_{coe}F(S_n) &= \sum_{v \in V} (1^2) = n. \\ D_{coe}F^*(S_n) &= \sum_{e_1=uv \in E} (1^2 + 1^2) = 2(n-1). \\ D_{coe}HM_1(S_n) &= \sum_{e_1=uv \in E} (1+1)^2 = 4(n-1). \end{aligned}$$

Case 2: If n is odd, we have

$D_{coe1} = \{v_1, v_2, \dots, v_{n-1}, w\}$, $D_{coe2} = \{v_1, v_2, \dots, v_{n-1}\}$, $T_{coe}(S_n) = 2$. Hence ,

$$d_{coe}(v) = \begin{cases} 1, & \text{if } v = w; \\ 2, & \text{if otherwise.} \end{cases}$$

then by definition 2.2, we get

$$\begin{aligned} D_{coe}F(S_n) &= 8n - 7. \\ D_{coe}F^*(S_n) &= 5(n-1). \\ D_{coe}HM_1(S_n) &= 9(n-1). \end{aligned}$$

□

Proposition 2.2 Consider a wheeled graph (W_n) , then

1. Suppose n is even, then

$$\begin{aligned} D_{coe}F(W_n) &= n. \\ D_{coe}F^*(W_n) &= 4n - 4. \\ D_{coe}HM_1(W_n) &= 8n - 8. \end{aligned}$$

2. Suppose n is odd, then

$$\begin{aligned} D_{coe}F(W_n) &= 8n - 7. \\ D_{coe}F^*(W_n) &= 5n - 5. \\ D_{coe}HM_1(W_n) &= 9n - 9. \end{aligned}$$

Proof: We can prove this as similar to Proposition 2.1

□

Proposition 2.3 Suppose G is a K_n graph (Fully connected graph) with $n \geq 3$, then

1. when n is even, then

$$\begin{aligned} D_{coe}F(G) &= n. \\ D_{coe}F^*(G) &= 2n^2 - 2n. \\ D_{coe}HM_1(G) &= 4n^2 - 4n. \end{aligned}$$

2. when n is odd, then

$$\begin{aligned} D_{coe}F(G) &= n2^{3n-3}. \\ D_{coe}F^*(G) &= (n^2 - n)2^{2n-2}. \\ D_{coe}HM_1(G) &= (n^2 - n)2^{(n^2-n)}. \end{aligned}$$

Proof: If G is a fully connected graph (K_n) , with $n \geq 3$, then

Case 1: Let n be even, the only even-paired dominating set of G is the vertex set of K_n . i.e, $D_{coe} = \{v_1, v_2, \dots, v_n\}$, Hence $d_{coe}(v) = 1$ for all $v \in V(K_n)$, and

$$\begin{aligned} D_{coe}F(K_n) &= n. \\ D_{coe}F^*(K_n) &= 2n(n-1). \\ D_{coe}HM_1(K_n) &= 4n(n-1). \end{aligned}$$

Case 2: Let n be odd, then the first row of all vertices is even, so there is $\binom{n}{1}$. A co-double dominant set has only one vertex with the first number $\binom{n}{2}$ co-dominant set with two vertices has the same number of degrees and has $\binom{n}{3}$. Co-pair dominated set includes all vertices but not all of the above-mentioned even-pair dominated sets include all vertices at a single level. The total set of public pair keys is: $\binom{n}{1} + \binom{n}{k} + \dots + \binom{n}{n} = 2^n - 1$. If we use the definition of co-pair dominance, we get $d_{coe}(v) = 2^{n-1}$. Using the definition of 2.2, we obtain the dominant index of graph which is even-paired.

$$\begin{aligned} D_{coe}F(K_n) &= \sum_{v \in V} \left(2^{n-1}\right)^3 = n2^{3n-3}. \\ D_{coe}F^*(K_n) &= \sum_{e_1=uv \in E} \left[(2^{n-1})^2 + (2^{n-1})^2 \right] = n(n-1)2^{2(n-1)}. \\ D_{coe}HM_1(K_n) &= n(n-1)2^{2n-1} \end{aligned}$$

□

Proposition 2.4 Let $K_{m,n}$ represent a complete bipartite graph with vertices of m, n .

1. Suppose n is even and m is odd, then

$$\begin{aligned} D_{coe}F(K_{m,n}) &= (m+8n)2^{3m-3}. \\ D_{coe}F^*(K_{m,n}) &= 5mn2^{2m-2}. \\ D_{coe}HM_1(K_{m,n}) &= 9mn2^{2m-2}. \end{aligned}$$

2. Suppose n is odd and m is even, then

$$\begin{aligned} D_{coe}F(K_{m,n}) &= (8m+n)2^{3n-3}. \\ D_{coe}F^*(K_{m,n}) &= 3mn2^{2n-1}. \\ D_{coe}HM_1(K_{m,n}) &= 9mn2^{2n-2}. \end{aligned}$$

3. suppose m and n both are odd, then

$$\begin{aligned} D_{coe}F(K_{m,n}) &= (m+n). \\ D_{coe}F^*(K_{m,n}) &= 2mn. \\ D_{coe}HM_1(K_{m,n}) &= 4mn. \end{aligned}$$

Proof: Assume that the first partition's vertex set is A , and the second partition's vertex set is B . It follows that $A = m, B = n$

Case 1: When m is odd and n is even, the degree of each vertex in A becomes even, while every vertex in B attains an odd degree in this case there exists $\binom{m}{1}$ jointly double-dominated sets containing a single one vertex from A , $\binom{m}{2}$ jointly such sets involving two vertices from A , and so on up to $\binom{m}{m}$. Consequently, the co-pair dominating set consists of all vertices belonging to A . The co-pair dominant set contains all vertices which belongs A . (This does not mean that all vertices of B are in Co-equal

dominant declines). Therefore, all co-double dominant clusters are set to B and $\binom{m}{1} + \binom{m}{2} + \dots + \binom{m}{m} = 2^m$. Using the concept of co-dominance, one obtains the total number as 2^m .

$$d_{coe}(v) = \begin{cases} 2^m, & \text{if } v \in B; \\ 2^{m-1}, & \text{if } v \in A. \end{cases}$$

By Definition 2.2, we get

$$\begin{aligned} D_{coe}F(K_{m,n}) &= \sum_{v \in A} (2^{m-1})^3 + \sum_{v \in B} (2^m)^3 = (m + 8n)2^{3m-3}. \\ D_{coe}F^*(K_{m,n}) &= \sum_{e_1=uv \in E(k_{m,n})} \left(2^{2m} + 2^{2m-2} \right) \\ &= 2^{2m} \left(1 + \frac{1}{4} \right) mn = 5mn2^{2m-2}. \\ D_{coe}HM_1(K_{m,n}) &= \sum_{e_1=uv \in E(k_{m,n})} \left(2^m + 2^{m-1} \right)^2 \\ &= \left(2^{2m} + 2^{2m} + 2^{2m-2} \right) = 9mn2^{2m-2} \end{aligned}$$

Case 2: suppose m even and n odd. By considering the same method as in Case 1, we get,

$$d_{coe}(v) = \begin{cases} 2^{n-1}, & \text{if } v \in B; \\ 2^n, & \text{if } v \in A. \end{cases}$$

From definition 2.2, we get

$$\begin{aligned} D_{coe}F(K_{m,n}) &= \sum_{v \in A} (2^n)^3 + \sum_{v \in B} (2^{n-1})^3. \\ &= m2^{3n} + n2^{3n-3} = (8m + n)2^{3n-3} \\ D_{coe}F^*(K_{m,n}) &= \sum_{e_1=uv \in E(k_{m,n})} \left[(2^n)^n + (2^{n-1})^2 \right] = 2^{2n-1}(mn). \\ D_{coe}HM_1(K_{m,n}) &= \sum_{e_1=uv \in E(k_{m,n})} \left(2^n + 2^{n-1} \right)^2 = 9mn2^{2n-2} \end{aligned}$$

Case 3: When m and n both are odd. In this case there is only single even-paired dominating set, which is the set of all vertices of $K_{m,n}$, hence $d_{coe}(v) = 1$ and by using the Definition 2.2 we get the results. \square

$$D_{coe}F(K_{m,n}) = m + n.$$

$$D_{coe}F^*(K_{m,n}) = 2mn.$$

$$D_{coe}HM_1(K_{m,n}) = 4n.$$

Definition 2.3 For any $v \in V$, the graph G is referred to be the k - even-paired dominance regular graph iff $d_{coe}(v) = k$.

examples:

1. Single even-paired dominating regular graph is S_n , where n is even.
2. Single even-paired dominating regular graph $K_n, n \geq 3$, with even number of vertices n

3. 2^{n-1} Even-paired dominating regular graphs of K_n with n odd.

Remark 2.1 (Observation) [1] Let G be graph with D_1, D_2, \dots, D_t as even-paired domination sets and $(\gamma_{coe}(G))$ of $G = (n, m)$ be the even-paired domination number, then

$$t\gamma_{coe}(G) \leq \sum_{V \in \mathcal{V}} d_{coe}(V) \leq t|V|$$

Remark 2.2 (Observation) [1] Suppose G is the Pendant-augmented n -wheel graph, then

$$1 \leq d_{coe}(v) \leq T_{coe}(G).$$




Figure 1: Pendant-augmented n -wheel graph H_4 , and H_5 .

Theorem 2.1 Let H_n be Pendant-augmented n -wheel graph with $n \geq 4$. Then

1. If n is even

$$\begin{aligned} D_{coe}F(H_n) &= (9n - 1)2^{3n-3}. \\ D_{coe}F^*(H_n) &= (n - 1) \left(10 * 2^{2n-4} + 2^{2n-3} \right). \\ D_{coe}HM_1(H_n) &= (n - 1) \left(6 * 2^{2n-2} + 2^{2n-2} \right). \end{aligned}$$

2. If n is odd

$$\begin{aligned} D_{coe}F(H_n) &= (n - 1) \left(2^{3n} + 2^{6n-6} + 3 * 2^{5n-5} + 3 * 2^{4n-4} + 2^{3n-3} \right). \\ D_{coe}F^*(H_n) &= (n - 1) \left(2^{2n-1} + 2^{n-2} + 1 \right). \\ D_{coe}HM_1(H_n) &= (n - 1) \left(7 + (9 * 2^{2n-2}) + 2^{2n} + 2^{2n-2} \right). \end{aligned}$$

Proof: Assume a Pendant-augmented n -wheel graph (H_n) with $n \geq 4$.

Case 1: There are $n - 1$ Vertices of even first degree $d(v)$ in this case if n is even. Therefore, $\binom{n-1}{1} + \binom{n-1}{2} + \dots + \binom{n-1}{n-1} = 2^{n-1} - 1$ is the number of Even-paired dominant sets generated by all the various substitutions for $(n - 1)$ vertices. The $2^{n-1} - 1$ even-paired dominant sets all include vertices of odd initial degree. Furthermore, another even-paired dominant set that only has the vertices of $d(v)$ odd is present. It follows that $T_{coe}(G) = 2^{n-1} - 1 + 1 = 2^{n-1}$. Thus, applying the even-paired domination degree concept, we obtain

$$d_{coe}(v) = \begin{cases} 2^{n-2}, & \text{for even } d(v); \\ 2^{n-1}, & \text{for odd } d(v). \end{cases}$$

Therefore by definition 2.2, we get

$$\begin{aligned} D_{coe}F(H_n) &= \sum_{d(v) \text{ is even}} (2^{n-2})^3 + \sum_{d(v) \text{ is even}} (2^{n-1})^3 \\ &= (n-1)2^{3n-6} + n2^{3n-3} = (9n-1)2^{3n-3}. \end{aligned}$$

Let E_1 represent the set of edges joining vertices with odd first degrees to Vertices with even first degrees. Let E_2 represent the set of edges joining vertices with even first degrees to other vertices with even first degrees, therefore

$$\begin{aligned} D_{coe}F^*(H_n) &= \sum_{d(v)=\text{even}} \left((2^{n-2})^2 + (2^{n-1})^2 \right) + \sum_{d(v)=\text{even}} \left((2^{n-2})^2 + (2^{n-2})^2 \right) \\ &= 5 * 2^{2n-4}(2n-4) + (n-1)2^{2n-3} \\ &= (n-1) \left(10 * 2^{2n-4} + 2^{2n-3} \right) \\ D_{coe}HM_1(H_n) &= \sum_{d(v)=\text{even}} (2^{n-2} + 2^{n-1})^2 + \sum_{d(v)=\text{even}} (2^{n-2} + 2^{n-2})^2 \\ &= \sum_{d(v)=\text{even}} (2^{2n-2} + 2^{2n-2} + 2^{2n-2}) + \sum_{d(v)=\text{even}} (2^{2n-2}) \\ &= (n-1) \left(6 * 2^{2n-2} + 2^{2n-2} \right) \end{aligned}$$

Case 2: Here, there exist n vertices of even first degree when n is odd. Therefore, the number of distinct ways to substitute n is equal to the number of even-paired dominant sets. even-paired dominant sets: $\binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n} = 2^n - 1$. It should be noted that every $2^n - 1$ even-paired dominating set contains every pendent vertex.

Another even-paired dominating set with the center vertex and all pendent vertices is also present. Hence $T_{coe}(H_n) = 2^n$.

By the definition of Even-paired dominating degree, we get

$$d_{coe}(v) = \begin{cases} 2^n, & \text{for off } d(v); \\ 2^{n-1}, & \text{excluding the central vertex in } d(v); \\ 2^{n-1} + 1, & \text{for } v \text{ even and } v \text{ the center vertices.} \end{cases}$$

Let B be the set of all vertices on the cycle, and A be the set of all pendent vertices.

By considering the Definition 2.2, we get

$$\begin{aligned} D_{coe}F(H_n) &= \sum_{v \in A} (2^{3n}) + \sum_{v \in B} \left((2^{3n-3})(2^{n-1} + 1)^3 \right) \\ &= (n-1)2^{3n} + (n-1) \left(2^{6n-6} + 3 * 2^{5n-5} + 3 * 2^{4n-4} + 2^{3n-3} \right) t \end{aligned}$$

Let $E_1 = \{e_1 = uv : d_{coe}(u) = 2^n, d_{coe}(v) = 2^{n-1}\}$,
 $E_2 = \{e_1 = uv : d_{coe}(u) = 2^{n-1}, d_{coe}(v) = 2^{n-1}\}$,
 $E_3 = \{e_1 = uv : d_{coe}(u) = 2^{n-1}, d_{coe}(v) = 2^{n-1} + 1\}$, where; $|E_1| = (n-1)$, $|E_2| = (n-1)$, $|E_3| = (n-1)$

Applying the Definition 2.2, we get

$$\begin{aligned}
D_{coe}F^*(H_n) &= \sum_{E_1} \left(2^{2n} + 2^{2n-2}\right) + \sum_{E_2} \left(2^{2n-2} + 2^{2n-2}\right) + \sum_{E_3} \left(2^{2n-2} + 2^{2n-2} + 2^{n-2} + 1\right). \\
&= (n-1) * \left(2^{2n} + 2^{2n-2}\right) + (n-1) * \left(2^{2n-2} + 2^{2n-2}\right) + (n-1) * \left(2^{2n-2} + 2^{2n-2} + 2^{n-2} + 1\right) \\
&= (n-1) * \left(2^{3n} + 2^{6n-6} + 3 * 2^{5n-5} + 3 * 2^{4n-4} + 2^{3n-3}\right). \\
D_{coe}HM_1(H_n) &= \sum_{E_1} \left(2^n + 2^{n-1}\right)^2 + \sum_{E_2} \left(2^{n-1} + 2^{n-1}\right)^2 + \sum_{E_3} \left(2^{n-1} + 2^{n-1} + 1\right)^2 \\
&= \sum_{E_1} \left(2^{2n} + 2^{2n} + 2^{2n-2}\right) + \sum_{E_2} \left(2^{2n}\right) \\
&= \sum_{E_3} \left(2^{2n-2} + 2^{2n-1} + 2^n + 1\right) \\
&= (n-1)2^{2n} \left(2 + \frac{1}{4}\right) + (n-1)2^{2n} + (n-1) \left(2^{2n-2} + 2^{2n-1} + 2^n + 1\right) \\
&= 9(n-1)2^{2n-2} + (n-1)2^{2n} + 7(n-1) + (n-1)2^{2n-2} \\
&= (n-1) * \left(7 + (9 * 2^{2n-2}) + 2^{2n} + 2^{2n-2}\right).
\end{aligned}$$

□




Figure 2: Ladder graph L_n .

Proposition 2.5 Let $P_2 \times P_n$ be Ladder graph.

Then

$$\begin{aligned}
D_{coe}F(P_2 \times P_n) &= 8192n - 14336. \\
D_{coe}F^*(P_2 \times P_n) &= 1535n - 3440. \\
D_{coe}HM_1(P_2 \times P_n) &= 1536n - 1280.
\end{aligned}$$

Proof: Let $P_2 \times P_n$ be Ladder graph. Then from the Figure 2. the first degree of the vertices sets $\{w_1, w_2, \dots, w_n, w'_1, w'_2, \dots, w'_n\}$ are three, and the first degree of the vertices sets $\{v_1, v_2, v'_1, v'_2\}$ are two. Then the number of even-paired dominating sets are :

$\binom{4}{1} + \binom{4}{2} + \binom{4}{3} + \binom{4}{4}$ + the set of all vertices of first degree 3 ($d(v)=3$). Hence $T_{coe}((P_2 \times P_n)) = 15 + 1 = 16$ even-paired dominating sets.

By considering, definition of even-paired dominating degree, we get

$$d_{coe}(v) = \begin{cases} 8, & \text{if } d(v) \text{ is even;} \\ 16, & \text{if } d(v) \text{ is odd.} \end{cases}$$

Therefore by the definition

$$\begin{aligned} D_{coe}F(P_2 \times P_n) &= \sum_{d(v)=\text{even}} 8^3 + \sum_{d(v)=\text{odd}} 16^3 \\ &= 2048 + 4096(2n - 4) \\ &= 2048 + 8192n - 16384 \\ &= 8192n - 14336. \end{aligned}$$

Let $|E_1 = \{e_1 = uv : d(u) \text{ is even and } d(v) \text{ is even}\}, |E_1| = 2$.

$E_2 = \{e_1 = uv : d(u) \text{ is odd and } d(v) \text{ is even}, |E_2| = 4$.

$E_3 = \{e_1 = uv : d(u) \text{ is odd and } d(v) \text{ is odd}, |E_3| = 3n - 8$. Then

$$\begin{aligned} D_{coe}F^*(P_2 \times P_n) &= \sum_{e_1=uv \in E} (P_2 \times P_n) \\ &= \sum_{e_1=uv \in E_1(G)} (P_2 \times P_n) + \sum_{e_1=uv \in E_2(G)} (P_2 \times P_n) + \sum_{e_1=uv \in E_3(G)} (P_2 \times P_n) \\ &= \sum_{e_1=uv \in E_1(G)} (64 + 64) + \sum_{e_1=uv \in E_2(G)} (64 + 256) + \sum_{e_1=uv \in E_3} 512(3n - 8) \\ &= 1535n - 3440. \end{aligned}$$

$$\begin{aligned} D_{coe}HM_1(P_2 \times P_n) &= \sum_{e_1=uv \in E} (P_2 \times P_n) \\ &= \sum_{e_1=uv \in E_1(G)} (P_2 \times P_n) + \sum_{e_1=uv \in E_2(G)} (P_2 \times P_n) + \sum_{e_1=uv \in E_3(G)} (P_2 \times P_n) \\ &= 1536n - 1280. \end{aligned}$$

□



Figure 3: double Star graph.

Theorem 2.2 Let $S_{m,n}$ be double Star graph with w_1, w_2 as center vertices of first star and second star respectively.

1. If m and n are even,

$$\begin{aligned} D_{coe}F(S_{m,n}) &= n + m + 2. \\ D_{coe}F^*(S_{m,n}) &= 2(n + m + 1). \\ D_{coe}HM_1(S_{m,n}) &= 2D_{coe}F^*(S_{m,n}). \end{aligned}$$

2. If m and n are odd,

$$\begin{aligned} D_{coe}F(S_{m,n}) &= 16 + 27(m+n). \\ D_{coe}F^*(S_{m,n}) &= 8 + 13(m+n). \\ D_{coe}HM_1(S_{m,n}) &= 16 + 25(m+n). \end{aligned}$$

3. If m is even and n is odd,

$$\begin{aligned} D_{coe}F(S_{m,n}) &= 8(m+n+1) + 1. \\ D_{coe}F^*(S_{m,n}) &= 3 + 8m + 5n. \\ D_{coe}HM_1(S_{m,n}) &= 3 + 16m + 9n. \end{aligned}$$

4. If m odd and n is odd,

$$\begin{aligned} D_{coe}F(S_{m,n}) &= 10 + 9(m+n). \\ D_{coe}F^*(S_{m,n}) &= 3 + 5m + 8n. \\ D_{coe}HM_1(S_{m,n}) &= 3 + 9m + 16n. \end{aligned}$$

Proof: Let $S_{m,n}$ be star graph with $n, m \geq 3$. Then

1. If m and n are even. In this case there is only single even-paired dominating set, $D = V(S_{m,n})$. Hence $d_{coe}(v) = 1$ for all $v \in V(S_{m,n})$. Applying the Definition 2.2, we get

$$\begin{aligned} D_{coe}F(S_{m,n}) &= \sum_{v \in V} 1^3 = n + m + 2. \\ D_{coe}F^*(S_{m,n}) &= \sum_{e_1 = uv \in E} (1 + 1) = 2(n + m + 1). \\ D_{coe}HM_1(S_{m,n}) &= 2(m + n + 1) = \sum_{e_1 = uv \in E} (1 + 1)^2 = 2D_{coe}M_2(S_{m,n}). \end{aligned}$$

2. When m and n are both odd, In this case the Even-paired dominating sets are

$$\begin{aligned} D_1 &= \{\text{all pendant vertices } w_1\}, D_2 = \{\text{all pendant vertices } w_2\}, \\ D_3 &= \{\text{all pendant vertices } w_1, w_2\}, D_4 = \{\text{all pendant vertices}\} \text{Hence} \\ d_{coe}(v) &= \begin{cases} 3, & \text{if } d(v) \text{ is pendant vertex or } d(v) = 1; \\ 2, & \text{if } d(v) = w_1, \text{ or } w_2. \end{cases} \end{aligned}$$

$$D_{coe}F(S_{m,n}) = \sum_{v \in A} 3^3 + \sum_{v \in \{w_1, w_2\}} 2^3 = 16 + 27(m+n).$$

Let $E_1 = \{e_1 = uv : d(u) \text{ is even and } d(v) = 1\}$. Then

$$\begin{aligned} D_{coe}F^*(S_{m,n}) &= 8 + \sum_{e_1 = uv \in E_1(S_{m,n})} (3^2 + 2^2) = 8 + 13(m+n), \\ D_{coe}HM_1(S_{m,n}) &= 16 + 25(m+n). \end{aligned}$$

3. If m is even and n is odd. In this case, we have $D_1 = v(S_{m,n})$, $D_2 = \{v(S_{m,n}) - w_2\}$, Hence

$$d_{coe}(v) = \begin{cases} 2, & \text{if } v \neq w_2; \\ 1, & \text{if } v = w_2. \end{cases}$$

$$D_{coe}F(S_{m,n}) = \sum_{V-1} 2^3 + 1 = 8(m+n+1).$$

let $E_1 = \{v_i w_1 : i = 1, 2, \dots, m\}$, and
 $E_2 = \{u_i w_2 : i = 1, 2, \dots, n\}$

$$D_{coe}F^*(S_{m,n}) = 3 + \sum_{e_1=uv \in E_1} (2^2 + 2^2) + \sum_{e_1=uv \in E_2} (1 + 2^2) = 3 + 8m + 5n.$$

$$D_{coe}HM_1(S_{m,n}) = 3 + \sum_{e_1=uv \in E_1} 16 + \sum_{e_1=uv \in E_2} 9 = 3 + 16m + 9n.$$

4. If m odd and n is odd.

In this Case we have $D_1 = v(S_{m,n})$, $D_2 = \{v_{m,n} - w_1\}$, Hence

$$d_{coe}(V) = \begin{cases} 2, & \text{if } v \neq w_1; \\ 1, & \text{if } v = w_1. \end{cases}$$

$$D_{coe}F(S_{m,n}) = \sum_{V(S_{m,n-1})} 2^3 + 1 = 10 + 9(m+n).$$

Let $E_1 = \{V_i w_1 : i = 1, 2, \dots, m\}$, then we get

$$D_{coe}F^*(S_{m,n}) = 3 + \sum_{e_1=uv \in E_1} 15 + \sum_{e_1=uv \in E_2} 8 = 3 + 5m + 8n,$$

$$D_{coe}HM_1(S_{m,n}) = 3 + \sum_{e_1=uv \in E_1} 9 + \sum_{e_1=uv \in E_2} 16 = 3 + 9m + 16n.$$

□

3. Some bounds for Modified Forgotten, hyper Zagreb Indices of Graph and Even-paired Domination Forgotten

In the present section, we find some lower and upper bounds on even-paired domination, Zagreb indices of graph, and some graph operations.

Proposition 3.1 *Let $G = (V, E)$. Then*

$$D_{coe}F(G) \leq nT_{coe}^2(G)$$

$$D_{coe}F(G) \leq T_{coe}^2(G)|E|$$

$$D_{coe}M_3(G) \leq 2T_{coe}(G)|E|.$$

The equality holds iff G is one even-paired dominating regular graph.

Proof: Case:1 Let $G = (V, E)$. Then

$$d_{coe}(v) \leq T_{coe}(G)$$

$$d_{coe}^3(v) \leq T_{coe}^3(G)$$

$$\sum_{v \in V} d_{coe}^3(v) \leq \sum_{v \in V} T_{coe}^3(G)$$

$$D_{coe}F(G) \leq T_{coe}^3(G)|V|.$$

Case 2: Let $G = (V, E)$. Then

$$\begin{aligned}
 d_{coe}(v) &\leq T_{coe}(G) \\
 d_{coe}^2(v) &\leq T_{coe}^2(G) \\
 d_{coe}^2(v) + d_{coe}^2(u) &\leq T_{coe}^2(G) + T_{coe}^2(G) \\
 \sum_{e_1=uv \in E} d_{coe}^2(v) + d_{coe}^2(u) &\leq \sum_{e_1=uv \in E} T_{coe}^2(G) + T_{coe}^2(G) D_{coe} F^*(G) \leq 2T_{coe}^2(G)|E|.
 \end{aligned}$$

Case 3: Let G be a graph. Then

$$\begin{aligned}
 d_{coe}(v) &\leq T_{coe}(G) \\
 d_{coe}(v) + d_{coe}(u) &\leq T_{coe}(G) + T_{coe}(G) \\
 \left(d_{coe}(v) + d_{coe}(u) \right)^2 &\leq \left(T_{coe}(G) + T_{coe}(G) \right)^2 \\
 \sum_{e_1=uv \in E} \left(d_{coe}(v) + d_{coe}(u) \right)^2 &\leq \sum_{e_1=uv \in E} \left(T_{coe}(G) + T_{coe}(G) \right)^2 \\
 \sum_{e_1=uv \in E} \left(d_{coe}(v) + d_{coe}(u) \right)^2 &\leq \sum_{e_1=uv \in E} \left(4T_{coe}^2(G) \right) \\
 D_{coe} HM_1(G) &\leq 4T_{coe}^2(G)|E|.
 \end{aligned}$$

To prove the equality if G has only one Even-paired domination set, then the only Even-paired domination set is the set of all Vertices of G , so $d_{coe}(v) = 1$, and (1), (2), (3) holds. □

References

1. Ahmed A., Omran, Ahmed H., Alsinai A., Vishnu Narayan Mishra, *Coeven Domination Zagreb Indices Of Graph*, Italin Journal of pure mathematics, (2021).
2. AlDzhabri K. S., Omran A. A., and Al-Harere M. N., *DG-domination topology in Digraph*, Journal of Prime Research in Mathematics, 17(2), 93-100, (2021).
3. Azari M., Iranmanesh A., *Harary Index of Some Nano-structures*, MATCH Commun. MAath. Comput. Chem., 71, 373-382, (2014).
4. Al-Fozan T., Manuel P., Rajasingh I., Sundara Rajan R., *Computing Szeged Index of Certain Nanosheets Using Partition Technique*, MATCH Commun. Math. Comput. Chem., 72, 339-353, (2014).
5. Alsini A., Alwardi A., and Soner N. D., *TOPOLOGICAL PROPERTIES OF GRAPHENE USING ψ_k polynomial*, Proceedings of the Jangjeon Mathematical Society, 24(3), 375-388, (2021).
6. Alsinai A., Alwardi A., Soner N. D., *On reciprocals leap function of graph*, Journal of Discrete Mathematical Sciences and Cryptography, 24(2), 307-324, (2021).
7. Alsinai A., Alwardi A., Ahmed H., Soner N. D., *Leap Zagreb Indices For The Central Graph Of Graph*, Journal of Prime Research in Mathematics, 12(2), 100-110, (2021).
8. Hanan Ahmed A. M., Alwardi A., and Salestina M. R., *On domination topological indices of graphs*, International Journal of Analysis and Applications, 19(1), 47-64, (2020).
9. Ahmed A. H., Alwardi A., Salestina M. R., and Soner N. D., *Forgotten domination, hyper domination and modified forgotten domination indices of graphs*, Journal of Discrete Mathematical Sciences and Cryptography, 24(2), 353-368, (2021).
10. Gutman I., Trinajstic N., *Graph theory and molecular orbitals. Total f-electron energy of alternant hydrocarbons*, Chem. Phys. Lett., 17, 535-538, (1972).
11. Gutman I., Rušćic B., Trinajstic N., Wilcox C. F., *Graph theory and molecular orbitals. XII. Acyclic polyenes*, J. Chem. Phys., 62, 3399-3405, (1975).
12. Gutman I., MiloVanoVic E., and MiloVanoVic I., *Beyond the Zagreb indices*, AKCE, Int. J. Graph Combin., (2018).

13. Kahat S. S., Omran A. A., and Al-Harere M. N., *Fuzzy equality co-neighborhood domination of graphs*, The International Journal of Nonlinear Analysis and Applications (IJNAA), 12(2),2008-6822, (2021).
14. Jabor A. A., and Omran A. A., *Topological domination in graph theory*, In AIP Conference Proceedings, 2334, 1, (2021).
15. Omran A. A., and Ibrahim T. A., *Fuzzy Even-paired domination of strong fuzzy graphs*, Int. J.
16. Shalaan M. M., Omran A. A., *Even-paired Domination in Graph*, International Journal of Control and Automation, 13(3), 330-334, (2020).
17. Yousif H.J. and Omran A. A., *Inverse 2- Anti Fuzzy Domination in Anti fuzzy graphs*, IOP Publishing Journal of Physics: Conference Series, 1818, 012072, (2021).
18. Raju S., Puttaswamy, Nayaka S..R. , *ABC, GA and AG Even-paired Domination Indices of Graph*,International Journal of Mathematics And its Applications,1(1), 75-84,(2023).

Pushpa N.

Department of Mathematics,

REVA UniVersity,

Bangalore, Karnataka-560064, India

E-mail address: pushparao3@gmail.com

and

Dhananjayamurthy B. V.,

Department of Mathematics,

Nitte Meenakshi Institute of Technology,

Yelahanka, Bengaluru Karnataka-560064, India

E-mail address: dmgyashams@yahoo.co.in