



## Existence of Solutions for Some Quasilinear Elliptic System with Weight and Measure-Valued Right Hand Side

El Houcine RAMI, Abdelkrim BARBARA and Elhoussine AZROUI

ABSTRACT: We prove the existence of a solution  $u$  for the nonlinear elliptic system:

$$\begin{aligned} -\operatorname{div} \sigma(x, u, Du) &= \mu + g(x, u, Du) \text{ in } \Omega \\ u &= 0 \text{ on } \partial\Omega \end{aligned}$$

where  $\mu$  is Radon measure on  $\Omega$  with finite mass and  $g$  satisfies some standards continuity and growth conditions. In particular, we show that if the coercivity rate of  $\sigma$  lies in the range  $]\frac{s+1}{s}, (\frac{s+1}{s})(2 - \frac{1}{n})]$  then  $u$  is approximately differentiable and the equation holds with  $Du$  replaced by  $\operatorname{ap} Du$ . The proof relies on an approximation of  $\mu$  by smooth functions  $f_k$  and a compactness result for the corresponding solutions  $u_k$ . This follows from a detailed analysis of the Young measure  $\{\delta_{u_k}(x) \otimes \vartheta(x)\}$  generated by the sequence  $(u_k, Du_k)$ , and the div-curl type inequality  $\langle \vartheta(x), \sigma(x, u, \cdot) \rangle \leq \bar{\sigma}(x) \langle \vartheta(x), \cdot \rangle$  for the weak limit  $\bar{\sigma}$  of the sequence.

Keywords: Nonlinear elliptic system, measure-valued, young measure, the div-curl type inequality.

### Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Hypothesis</b>	<b>2</b>
<b>3</b>	<b>Some Preliminary Lemmas</b>	<b>4</b>
<b>4</b>	<b>Approximate Solutions and a Priori Bounds</b>	<b>6</b>
<b>5</b>	<b>A Div-Curl Inequality</b>	<b>9</b>
<b>6</b>	<b>Passage to the Limit</b>	<b>9</b>

### 1. Introduction

We consider the existence and compactness questions for elliptic systems of the form

$$-\operatorname{div} \sigma(x, u(x), Du(x)) = \mu + g(x, u, Du) \text{ in } \Omega \tag{1.1}$$

$$u = 0 \text{ on } \partial\Omega \tag{1.2}$$

with measure-valued right hand side  $\mu \in M(\Omega, \mathbb{R}^m)$  on an open, bounded domain  $\Omega$  in  $\mathbb{R}^n$ . And the weight functions  $\omega^*$  defined by  $\omega = \{\omega_{ij}, 0 \leq i \leq n, 1 \leq j \leq m\}$  with  $\omega_{ij}^* = \omega_{ij}^{1-p'}$  on this paper we are interested in the solution  $u$  in the Sobolev space  $W_0^{1,p}(\Omega, \omega, \mathbb{R}^m)$ , and estimation of the weak Lebesgues spaces. We assume that  $\sigma$  satisfies the following hypotheses  $(H_0) - (H_3)$  and  $g$  satisfies  $G_0 - G_1$ . Here  $M^{m \times n}$  denotes the space of real  $m \times n$  matrices equipped with inner product  $M : N = M_{ij} N_{ij}$  ( we use the usual summation convention) and the tensor product  $a \otimes b$  of two vectors  $a, b \in \mathbb{R}^n$  is defined to be the matrix  $(a_i b_j)_{i,j=1,\dots,m}$ . Let  $\omega = \{\omega_{ij} | 0 \leq i \leq n; 1 \leq j \leq m\}$  the system for weight functions on  $\Omega$  satisfying:

$$\omega_{ij} \in L^1(\Omega), \quad \omega_{ij}^{-\frac{1}{p-1}} \in L^1_{loc}(\Omega) \tag{1.3}$$

$$\omega_{ij}^{-s} \in L^1(\Omega) \tag{1.4}$$

---

2020 *Mathematics Subject Classification*: 35J46, 35J62.

Submitted February 13, 2026. Published April 17, 2026

for same  $s$ . The Sobolev space  $W^{1,p}(\Omega, \omega, \mathbb{R}^m)$  with

$$W^{1,p}(\Omega, \omega, \mathbb{R}^m) = \{u \in L^p(\Omega, \bar{\omega}_0, \mathbb{R}^m) \mid \bar{\omega}_0 = \omega_{0j}; 1 \leq j \leq m; \frac{\partial u_j}{\partial x_i} \in L^p(\Omega, \omega_{ij}); 1 \leq j \leq m; 1 \leq i \leq n\}$$

The Sobolev space  $W^{1,p}(\Omega, \omega, \mathbb{R}^m)$  equipped with the norm

$$\|u\|_{1,p,\omega}^p = \sum_{j=1}^m \int_{\Omega} \omega_{0j} |u_j|^p dx + \sum_{i,j} \int_{\Omega} \omega_{ij} |D_{ij}u|^p dx.$$

The condition (1.3) we give  $(W^{1,p}(\Omega, \omega, \mathbb{R}^m), \|\cdot\|_{1,p,\omega})$  Banach space and  $C_0^\infty(\Omega, \mathbb{R}^m)$  subspace of  $(W^{1,p}(\Omega, \omega, \mathbb{R}^m))$ . The space  $(W_0^{1,p}(\Omega, \omega, \mathbb{R}^m))$  is the fermenter of  $C_0^\infty(\Omega, \mathbb{R}^m)$  in  $(W^{1,p}(\Omega, \omega, \mathbb{R}^m))$  for the norm  $\|\cdot\|_{1,p,\omega}^p$ . The condition (1.4), we give

$$(W^{1,p}(\Omega, \omega, \mathbb{R}^m)) \hookrightarrow (W^{1,p_s}(\Omega, \mathbb{R}^m)) \hookrightarrow L^r(\Omega, \mathbb{R}^m), \quad (1.5)$$

for all  $1 \leq r < p_s^*$  if  $p.s < n(s+1)$ , and

$$\forall r \geq 1 \text{ if } p.s \succ n(s+1) \text{ with } p_s = \frac{p.s}{s+1} \text{ and } p_s^* = \frac{n.p.s}{n(s+1)-p.s}$$

## 2. Hypothesis

( $H_0$ ) (Continuity)  $\sigma : \Omega \times \mathbb{R}^m \times \mathbb{M}^{m \times n} \rightarrow \mathbb{M}^{m \times n}$  is a Carathéodory function, i.e:

$x \mapsto \sigma(x, u, p)$  is measurable for every  $(u, p)$  and  $(u, p) \mapsto \sigma(x, u, p)$  is continuous for almost every  $x \in \Omega$ .

( $H_1$ ) (Coercivity and growth): There exist constants  $c_1, c_2, \beta > 0$  and  $\lambda_1 \in L^{p'}(\Omega)$ ,  $\lambda_2 \in L^1(\Omega)$ ,  $\lambda_3 \in L^{(\frac{p}{\theta})'}(\Omega)$ ,  $0 < \theta < p$ ,  $1 < q < \frac{p^2}{\theta}$ , such that, for all  $1 \leq r \leq n$ , and  $1 \leq s \leq m$ :

$$|\sigma_{rs}(x, u, F)| \leq \beta \omega_{rs}^{\frac{1}{p}} [\lambda_1 + c_1 \sum_{j=1}^m \gamma_j^{\frac{1}{p'}} |u_j|^{\frac{q}{p'}} + c_1 \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |F_{ij}|^{p-1}]$$

$$\sigma(x, u, F) : F \geq -\lambda_2 - \sum_{j=1}^m \lambda_3 \gamma_j^{\frac{\theta}{p}} |u_j|^{\frac{q\theta}{p}} + c_2 \sum_{i,j} \omega_{ij} |F_{ij}|^p.$$

( $H_2$ ) (Monotonicity)  $\sigma$  satisfies one following conditions:

a)- For all  $x \in \Omega$ ,  $u \in \mathbb{R}^m$  the function  $F \mapsto \sigma(x, u, F)$  is a  $C^1$  and monotone function, which means  $(\sigma(x, u, F) - \sigma(x, u, G)) : (F - G) \geq 0$ , for all  $x \in \Omega$ ,  $u \in \mathbb{R}^m$ , and  $F, G \in \mathbb{M}^{m \times n}$ .

b)- There exist a function  $W : \Omega \times \mathbb{R}^m \times \mathbb{M}^{m \times n} \rightarrow \mathbb{R}$  such that:

$\sigma(x, u, F) = \frac{\partial W}{\partial F}(x, u, F)$ , and the function  $F \mapsto W(x, u, F)$  is a  $C^1$  and convex function.

c)-  $\sigma$  is strictly monotone, i-e;  $\sigma$  is monotone and  $(\sigma(x, u, F) - \sigma(x, u, G)) : (F - G) = 0$ . implies  $F = G$ .

d) A function  $F \mapsto \sigma(x, u, F)$  is strictly  $p$ -quasi-monotone, i-e:

$\int_{\mathbb{M}^{m \times n}} (\sigma(x, u, \lambda) - \sigma(x, u, \bar{\lambda})) : (\lambda - \bar{\lambda}) d\nu(\lambda) > 0$ , for all homogeneous  $W^{1,p}$ -gradient Young measures  $\nu$  with center of mass  $\bar{\lambda} = \langle \nu; Id \rangle = \int_{\mathbb{M}^{m \times n}} \lambda d\nu(\lambda)$  which are not a single Dirac mass.

( $H_3$ ) (structure conditions)

i)- (Angle condition) for all  $x \in \Omega$ ,  $u \in \mathbb{R}^m$  and  $F \in \mathbb{M}^{m \times n}$  there holds

$\sigma(x, u, F) : MF \geq 0$ , for all matrix  $M \in \mathbb{M}^{m \times m}$  of the form  $M = Id - a \otimes a$  with  $|a| \leq 1$ .

ii)- (The sign condition) for all  $x \in \Omega$ ,  $u \in \mathbb{R}^m$  and  $F \in \mathbb{M}^{m \times n}$ ,  $\sigma_j(x, u, F) : F_j \geq 0$ , for all  $1 \leq j \leq m$  with  $F_j$  and  $\sigma_j$  are the cologne  $j$  of matrix  $F$  and  $\sigma$ .

( $H_4$ ) (The Hardy-Type Inequality) There exist  $c > 0$ , one weighted function  $\gamma = (\gamma_j)_{1 \leq j \leq m}$ , and the parameter  $1 < q < \frac{p^2}{\theta}$  ( $H_1$ ), such that:

$$\left( \sum_{j=1}^m \int_{\Omega} \gamma_j |u_j|^q dx \right)^{\frac{1}{q}} \leq c \left( \sum_{i,j} \int_{\Omega} \omega_{ij} |D_{ij}u|^p dx \right)^{\frac{1}{p}}$$

and the expression  $\left( \sum_{i,j} \int_{\Omega} \omega_{ij} |D_{ij}u|^p dx \right)^{\frac{1}{p}}$  is a norm equivalent to the norm  $\|\cdot\|_{1;p;\omega}$  in  $W_0^{1,p}(\Omega, \omega, \mathbb{R}^m)$ .

( $G_0$ ) (Continuity)  $g : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^m$  is Caratheodory function.

( $G_1$ ) (Growth) There exist:  $b_1 \in L^{p'}(\Omega)$ ,  $C_1, C'_1 > 0$  such that  $|g_j(x, u, F)| \leq [b_1(x) + C_1 \gamma_j^{\frac{1}{p'}} |u_j|^{\frac{q}{p'}} + C'_1 \sum_{r,s} \omega_{rs}^{\frac{1}{p'}} |F_{rs}|^{p-1}] \omega_{0j}^{\frac{1}{p}}$

### Remark 2.1

1. Assumption ( $H_0$ ) ensures that  $\sigma(x, u(x), U(x))$  is measurable on  $\Omega$  for measurable function  $u : \Omega \rightarrow \mathbb{R}^m$  and  $U : \Omega \rightarrow \mathbb{M}^{m \times n}$ . A typical example for a function  $\sigma$  satisfying ( $H_0$ ) is  $\sigma(x, u, p) = \zeta(x, u, p)p$  with a real valued non-negative function  $\zeta$ .
2. A serious technical obstacle is that for  $p_s \in (1, 2 - 1/n]$  solutions of the system (1.1) in general do not belongs to the Sobolev space  $W^{1,1}(\Omega, \omega, \mathbb{R}^m)$ .

This fact has led to the use of normalized solutions in [2] and generalized entropy solutions in [5] for elliptic equations of the above type. We will use a notion of solution where the weak derivative  $Du$  is replaced by the approximate derivative  $apDu$ . Recall that a measurable function  $u$  is said to be approximately differentiable at  $x \in \Omega$  if there exists a matrix  $F_x \in \mathbb{M}^{m \times n}$  such that:

for all  $\epsilon > 0$ ,  $\lim_{r \rightarrow 0} \frac{1}{r^n} \text{meas} \{y \in B(x, r) : |u(y) - u(x) - F_x(y - x)| > \epsilon r\} = 0$ . We write:  
 $apDu(x) = F_x$ .

**Definition 2.1** A measurable function  $u : \Omega \rightarrow \mathbb{R}^m$  is called a solution of the system (1.1) if:

- (i)  $u$  is almost everywhere approximately differentiable.
- (ii)  $\eta \circ u \in W^{1,1}(\Omega, \omega, \mathbb{R}^m)$ , for all,  $\eta \in C_0^1(\mathbb{R}^m, \mathbb{R}^m)$ .
- (iii)  $\sigma(x, u, apDu) \in L^1(\Omega, \mathbb{M}^{m \times n})$ ;
- (iv) The equation:  $-\text{div} \sigma(x, u(x), Du(x)) = \mu + g(x, u, Du)$  holds in the sense of distributions. Moreover we say that  $u$  satisfies the boundary condition (1.2) if  $\eta \circ u \in W_0^{1,1}(\Omega, \omega, \mathbb{R}^m)$ , for all,  $\eta \in C^1(\mathbb{R}^m, \mathbb{R}^m) \cap L^\infty(\mathbb{R}^m, \mathbb{R}^m)$  with  $\eta = \text{id}$  on  $B(0, r)$ , for some  $r > 0$ , and  $|D\eta(y)| \leq c \cdot (1 + |y|)^{-1}$ , with  $c < \infty$ .

### Remark 2.2

1. The conditions in Definition (2.1) (except (ii)) are the weakest possible in order to define the system (1.1) in the sense of distributions. Note that if  $u$  is approximately differentiable, then  $apDu$  is measurable, so  $\sigma(\cdot, u, apDu)$  is measurable.
2. The assumption  $\eta \circ u \in W^{1,1}(\Omega, \mathbb{R}^m)$  ensures minimal regularity of  $u$ . For example, if  $\mu = 0$ , and  $\sigma(x, u, p) = \sigma(p)$  with  $\sigma(0) = 0$ , then piecewise constant functions  $u$  satisfy  $apDu = 0$  a.e, but are not admissible solutions. The following theorem is the main result in this paper .

**Theorem 2.1** *Let  $\Omega$  be a bounded, open set. We suppose that the hypotheses  $(H_0) - (H_2) - (H_3)$  and the coercivity condition in  $(H_1)$  are satisfied and  $g$  satisfies  $(G_0) - (G_1)$ . Let  $\mu$  denote a  $\mathbb{R}^m$ -valued Radon measure on  $\Omega$  with finite mass. Then the system (1.1)-(1.2) has a solution  $u$  in the sense of definition 1, which satisfies the weak Lebesgue space estimate:*

$$\|u\|_{L^{t_{p_s}^*, \infty}(\Omega, \mathbb{R}^m)}^* + \|apDu\|_{L^{t_{p_s}, \infty}(\Omega, M^{m \times n})}^* \leq C, \quad (2.1)$$

with the constant  $C$  depends of  $|\Omega|, c, c_2$ , and  $\|\lambda_3\|_{L^{(\frac{p}{\beta})}(\Omega)}$ , with  $t_{p_s} = \frac{n(p_s-1)}{n-1}$  and  $t_{p_s}^* = \frac{n(p_s-1)}{n-p_s}$  is the Sobolev exposed of  $t_{p_s}$ . If  $c_2 = 0$  the right hand side of (1.3) reduces to  $C(c_1) \left\| \mu^{\frac{1}{p-1}} \right\|_M$ .

### Remark 2.3

1. If  $p_s > 2 - \frac{1}{n}$ , then  $t_{p_s} > 1$  and  $Du \in L^1(\Omega, M^{m \times n})$ .
2. If  $p > n$  one can replace the  $L^{s, \infty}$ -norm of  $u$  in (1.3) by the  $C^0, \beta$ -norm with  $\beta = 1 - \frac{n}{p}$ . For  $p = q = n$  it is an open question whether  $Du \in L^{n, \infty}$ . See Section 7 [4] for the (weaker) inclusion  $u \in BMO_{loc}$ .
3. The exponent in (1.3) are optimal as can be seen from the nonlinear Green's function  $G_p(x) = c|x|^{-\frac{n}{s^*}}$  for the  $p$ -Laplace equation:  $-div(|Du|^{p-2} Du) = \delta_0$  in  $\mathbb{R}^m, n \geq 3$ . In particular,  $L^{s, \infty}$  cannot be replaced by  $L^s$ . with  $(L^{s, \infty})$ , is a Laurent space.
4. The pointiest monotonicity condition can be replaced by a weaker integrated version, called quasi-monotonicity,

The Key point in the proof of the theorem, is the div-curl inequality for the Young measure  $\{\{\vartheta_x\}_{x \in \Omega}\}$  generated by a sequence  $Du_k$  of gradients of approximate solutions. Together with the identity. (1.5):

$apDu(x) = \langle \vartheta_x, Id \rangle$ . The div-curl inequality implies easily that  $\sigma(\cdot, u_k, Du_k)$  converges weakly in  $L^1$  to  $\sigma(\cdot, u, apDu)$ . (1.5) is a consequence of general properties of young measures if  $p_s > 2 - \frac{1}{n}$  since in this case  $Du_k$  is bounded in  $L^s$  for some  $s > 1$ . If  $1 < p_s \leq 2 - \frac{1}{n}$  one only has the weaker bounds.

### 3. Some Preliminary Lemmas

In this section, we will also use the Young measures, and Inequality div-curl for assume the convergence of subsequence  $u_k \rightarrow u$  in measure and for almost every subsequence, with  $u$  is approximately differentiable, and  $apDu = \langle \nu_x, id \rangle$ ,  $\nu_x$  is the Young measures generated by a sequence  $Du_k$ .

**Lemma 3.1** *Let  $u_k: \Omega \rightarrow \mathbb{R}^m$  a sequence of measurable functions such that:*

$$\sup_{k \in \mathbb{N}} \int_{\Omega} |u_k|^s dx < +\infty \text{ for some } s > 0. \quad (3.1)$$

*We suppose that for each  $\alpha > 0$  the sequence of truncated functions  $\{T_{\alpha}(u_k)\}_{k \in \mathbb{N}}$  is precompact in  $L^1(\Omega, \mathbb{R}^m)$ . Then there exists a measurable function  $u$  on  $\Omega$  such that for a subsequence  $u_k \rightarrow u$  in measure.*

#### Proof

Choose a subsequence of  $\{u_k\}$  (not relabeled) which generates a Young measure  $\{\vartheta_x\}_{x \in \Omega}$ . By 3.1 and Theorem (Young, Tartar, Ball) the measure  $\nu_x$  are probability measure for almost every a  $x \in \Omega$  and  $T_{\alpha}(u_k) \rightarrow v_{\alpha} = \langle \nu_x; T_{\alpha} \rangle$ , weakly in  $L^1(\Omega, \mathbb{R}^m)$  and in fact strongly since  $T_{\alpha}(u_k)$  is precompact in  $L^1$ . Consequently there exists a subsequence such that:  $T_{\alpha}(u_{k_l}) \rightarrow v_{\alpha}$  almost uniformly, i-e:

$$T_{\alpha}(u_{k_l}) \rightarrow v_{\alpha} \text{ uniformly up to a set of arbitrary small measure.} \quad (3.2)$$

Let  $M_{\alpha} = \{x \in \Omega : |v_{\alpha}(x)| < \alpha\}$ . Then for each  $\epsilon > 0$  and  $\delta > 0$  there exists a set  $E_{\epsilon}$  of measure  $\text{meas}(E_{\epsilon}) < \epsilon$  and an index  $l_0(\epsilon; \delta)$  such that:  $|T_{\alpha}(u_{k_l})| < |v_{\alpha}(x)| + \delta$  for all  $x \in M_{\alpha} \setminus E_{\epsilon}$  and all  $l > l_0$ .

It follows that  $u_{k_l}(x) \rightarrow v_\alpha(x)$  for almost every  $x \in M_\alpha \setminus E_\epsilon$  consider first  $x \in M_\beta$ ;  $\beta < \alpha$  and then the union over  $\beta < \alpha$ ). Since  $\epsilon > 0$  was arbitrary it follows that  $v_x = \delta_{v_\alpha}(x)$  for almost every  $x \in M_\alpha$ . In view of the Ball's theorem it suffices to show that  $\cup M_\alpha$  has full measure. Now clearly  $M_\alpha \subset M_\beta$  for  $\alpha < \beta$  since  $T_\beta(u_{k_l}) \rightarrow T_\beta(v_\alpha) = v_\alpha$  almost everywhere in  $M_\alpha$  and therefore  $v_\alpha = v_\beta$  on  $M_\alpha$ . By (3.2) there exists for each  $\epsilon > 0$  a set  $E_\epsilon$ , and an index  $l_0(\epsilon, \alpha)$  such that  $\text{meas}(E_\epsilon) < \epsilon$  and  $|u_{k_l}| \geq |T_\alpha(u_{k_l})| \geq \frac{\alpha}{2}$  on  $(\Omega \setminus E_\epsilon) \setminus M_\alpha$  for all  $l \geq l_0$ . In view of (3.2) this implies  $\text{meas}((\Omega \setminus E_\epsilon) \setminus M_\alpha) \leq \frac{\epsilon}{\alpha^s} \rightarrow 0$  we deduce  $\text{meas}(\Omega \setminus \cup M_\alpha) = \lim_{\alpha \rightarrow \infty} \text{meas}(\Omega \setminus M_\alpha) = 0$   $\square$

**Lemma 3.2** *Let  $\Omega$  be a domain in  $\mathbb{R}^n$  with  $|\Omega| < \infty$  and  $u_k \in W^{1,1}(\Omega, \mathbb{R}^m)$ . Suppose that there exist  $p > 1$  and  $s > 0$  such that:*

$$\sup_k \sum_{i,j} \int_{|u_k| \leq \alpha} \omega_{ij} |D_{ij} u_k|^p dx \leq c(\alpha) < \infty, \quad \forall \alpha > 0, \quad (3.3)$$

and  $\sup_{k \in \mathbb{N}} \int_\Omega |u_k|^s dx \leq c < \infty$ . Then there exist a subsequence  $u_{k_j}$  and a measurable function  $u : \Omega \rightarrow \mathbb{R}^m$  such that  $u_{k_j} \rightarrow u$  in measure. Moreover  $u$  is for almost every  $x \in \Omega$  approximately differentiable, for all  $\eta \in C_0^1(\Omega, \mathbb{R}^m)$  there holds  $\eta \circ u \in W^{1,p}(\Omega, \omega, \mathbb{R}^m)$ . if  $u_k \in W_0^{1,1}(\Omega, \mathbb{R}^m)$  then  $\eta \circ u \in W_0^{1,1}(\Omega, \mathbb{R}^m) \cap W^{1,p}(\Omega, \omega, \mathbb{R}^m)$  provided that  $\eta = id$  on  $B(0, r)$  for some  $r > 0$ .

**Proof**

Choose

$$(u_k)_\alpha = \begin{cases} u_k & \text{if } |u_k| \leq \alpha, \\ 0 & \text{if } |u_k| > \alpha. \end{cases}$$

For the hypotheses:

$$\sum_{i,j} \int_\Omega \omega_{ij} |D_{ij} (u_k)_\alpha|^p dx = \sum_{i,j} \int_{|u_k| \leq \alpha} \omega_{ij} |D_{ij} u_k|^p dx \leq c(\alpha) < \infty.$$

Then,  $(u_k)_\alpha \in W_0^{1,1}(\Omega, \omega, \mathbb{R}^m)$  and for (1.5),  $(H_4)$  and  $|D|u|| \leq |Du|$  we have:

$$\begin{aligned} \int_\Omega |DT_\alpha(|u_k|)|^{p_s} dx &= \int_{|u_k| \leq \alpha} |D|u_k||^{p_s} dx \leq \\ &\sum_{i,j} \int_\Omega \omega_{ij} |D_{ij} (u_k)_\alpha|^p dx \leq c(\alpha) < +\infty \end{aligned}$$

Hence by the compact Sobolev embedding  $W_s^{1,p_s}(\Omega) \hookrightarrow L^{p_s}(\Omega)$ , we have  $\{T_\alpha(|u_k|)\}$  is precompact in  $L^1(\Omega)$ . And, if  $\eta \in C_0^\infty(B(0, 3\alpha), \mathbb{R}^m)$  a symmetric radial such that  $\eta = id$  on  $B(0, 2\alpha)$ , then by (1.3)

and (3.3):  $\sum_{i,j} \int_\Omega \omega_{ij} |D_{ij}(\eta(u_k))|^p dx = \sum_{i,j} \int_{|u_k| \leq \alpha} \omega_{ij} |D_{ij}(u_k)|^p dx + \sum_{i,j} \int_{\alpha < |u_k| \leq 2\alpha} \omega_{ij} |D_{ij}(id)|^p dx +$

$\sum_{i,j} \int_{2\alpha < |u_k| \leq 3\alpha} \omega_{ij} |D_{ij}(\eta(u_k))|^p dx \leq c(\alpha) + c \sum_{i,j} \|\omega_{ij}\|_{L_{loc}^1(\Omega)} + c < \infty$ . Then, by (1.5),  $\eta(u_k)$  is

precompact in  $L^{p_s}(\Omega, \mathbb{R}^m)$ , and as in Lemma 8 [2], there exist a measurable function  $u : \Omega \rightarrow \mathbb{R}^m$  such that  $u_k \rightarrow u$  in measure, with  $u(x) = \langle \vartheta_x, id \rangle$  for almost every  $x \in \Omega$  and  $u$  is approximately differentiable because  $\eta(u_k) \rightarrow \eta(u)$  in  $W^{1,p}(\Omega, \omega, \mathbb{R}^m)$  and  $apDu = ap(\eta \circ u)$ .  $\square$

**Lemma 3.3** *Let  $u_k$  be as in Lemma (3.2) with  $p > 1$ . Then the Young measure  $\vartheta_x$  generated by (a subsequence of)  $Du_k$  has the following properties:*

- (a)  $\vartheta_x$  is a probability measure for almost every  $x \in \Omega$ .
- (b)  $\vartheta_x$  has finite  $p_s$ -th moment for almost every  $x \in \Omega$ , i-e  $\int_{M^{m \times n}} |\lambda|^{p_s} d\vartheta_x(\lambda)$  is finite for almost every  $x \in \Omega$ .
- (c)  $\vartheta_x$  satisfies  $\langle \vartheta_x, id \rangle = apDu(x)$  almost everywhere in  $\Omega$ .

(d)  $\vartheta_x$  is a homogeneous  $W^{1,p_s}$ -gradient young measure for almost every  $x \in \Omega$ .

**Proof** Let  $\widetilde{\vartheta}_x$  denote the Young measure generated by (a subsequence of) the sequence  $\{(u_k, Du_k)\}$ . By Lemma 3.2 we have :

$$\widetilde{\vartheta}_x = \delta_{u(x)} \otimes \vartheta_x.$$

Let  $\eta \in C_0^\infty(B(0, 2\alpha), \mathbb{R}^m)$ ,  $\eta = Id$  on  $B(0, \alpha)$ , and let  $\vartheta^\eta$  be the Young measure generated by

$$D(\eta \circ u_k) = (D\eta)(u_k)Du(x),$$

then  $\vartheta^\eta$  is a probability measure, has finite  $p$ -th moment and

$$\langle \vartheta^\eta, Id \rangle = (D(\eta \circ u))(x) = D\eta(u(x))Du(x).$$

It follows for  $\varphi \in C_0^\infty(M^{m \times n})$ , that:

$$\varphi(D(\eta \circ u_k)) \rightharpoonup \langle \vartheta^\eta, \varphi \rangle = \int_{M^{m \times n}} \varphi(\lambda) d\vartheta_x^\eta(\lambda).$$

Based on the proof (3.2), we have  $\sum_{i,j} \int_{\Omega} |\omega_{ij} D_{ij}(\eta \circ u_k)|^p dx < \infty$ , and by (1.5)

$\sup_{k \in \mathbb{N}} \int_{\Omega} |D(\eta \circ u_k)|^{p_s} dx < \infty$ , and the (Ball's Theorem, proof lemma 9 [2]) we conclude: (a)-(b)-(c)- and (d)  $\square$

#### 4. Approximate Solutions and a Priori Bounds

We introduce the following approximating problems:

$$-div\sigma(x, u_k, Du_k) = f_k + g(x, u_k, Du_k) \text{ in } \Omega. \quad (4.1)$$

$$u_k = 0 \text{ on } \partial\Omega. \quad (4.2)$$

With  $f_k \in W^{-1,p'}(\Omega, \omega^*, \mathbb{R}^m) \cap L^1(\Omega, \mathbb{R}^m)$  and  $f_k \rightharpoonup^* \mu$  in  $M(\Omega, \mathbb{R}^m)$  such that:

$\|f_k\|_{L^1(\Omega, \mathbb{R}^m)} \leq \|\mu\|_{M(\Omega, \mathbb{R}^m)}$ . By [7] and [6], and using the assumptions:  $(H_0)$ ,  $(H_1)$ ,  $(H_2)$ ,  $(H_4)$ ,  $G_0$  and  $G_0$ , the problem (4.1)-(4.2) has a solution  $u_k$  with  $u_k \in W_0^{1,p}(\Omega, \omega, \mathbb{R}^m)$   $u_k$  is the subsequence approximates solutions of (1.1)-(1.2). The results of theorems (2.1) is the consequence of the following proposition:

**Proposition 4.1** *Let,  $f \in L^1(\Omega, \mathbb{R}^m)$  and  $\sigma$  satisfies  $(H_0)$ , the coercivity of  $(H_1) - (H_3)$  and  $g$  satisfies  $(G_0) - (G_1)$ . If  $u \in W_0^{1,p}(\Omega, \omega, \mathbb{R}^m)$  is a solution of:*

$$-div\sigma(x, u, Du) = f + g \quad \text{in } \Omega. \quad (4.3)$$

in the sense of distributions. Then

$$u \in L^{t_{p_s}^*, \infty}(\Omega, \mathbb{R}^m), \quad Du \in L^{t_{p_s}, \infty}(\Omega, \mathbb{R}^m)$$

and

$$\|u\|_{L^{t_{p_s}^*, \infty}(\Omega, \mathbb{R}^m)}^* + \|Du\|_{L^{t_{p_s}, \infty}(\Omega, M^{m \times n})}^*$$

$$\leq C \left( |\Omega|, \|\lambda_1\|_{L^1(\Omega)}, \|\lambda_3\|_{L^{(\frac{p}{2})}'(\Omega)}, \|f\|_{L^1(\Omega; \mathbb{R}^m)}, \|b_1\|_{L^{p'}} \right) \sum_{i,j} \int_{|u| \leq \alpha} \omega_{ij} |D_{ij}u|^p dx \leq M\alpha + L, \quad \forall \alpha > 0,$$

$M$  and  $L$  are the constants depends on:

$$\|\lambda_1\|_{L^1(\Omega)}, \|\lambda_3\|_{L^{(\frac{p}{2})}'(\Omega)}; \|f\|_{L^1(\Omega; \mathbb{R}^m)}, c_2, \|b_1\|_{L^{p'}}$$

**Proof i)-** We suppose the condition of l'angle in  $(H_3)$ . Let  $\alpha > 0$ . Testing  $T_\alpha(u)$  in (4.3) and we use the coercivity condition in  $(H_1)$ , the growth condition in  $(G_1)$ , the Hardy type inequality in  $(H_4)$  and Hölder inequality, we have:

$$\begin{aligned} c_2 \cdot \sum_{i,j} \int_{|u| \leq \alpha} \omega_{ij} |D_{ij} u|^p dx &\leq \alpha \|f\|_{L^1(\Omega, \mathbb{R}^m)} + \alpha \sum_j \left( \int_{|u| \leq \alpha} \omega_{0j} \right)^{\frac{1}{p}} \left( \|b_1\|_{L^{p'}} + \left( \sum_j \int_{|u| \leq \alpha} \gamma_j |u_j|^q dx \right)^{\frac{1}{p'}} \right) \\ &\quad + \|\lambda_2\|_{L^1(\Omega)} + c \|\lambda_3\|_{L(\frac{p}{\theta})}' \left( \sum_{j=1}^m \int_{|u| \leq \alpha} \gamma_j |u_j|^q dx \right)^{\frac{\theta}{p}} \end{aligned} \quad (4.4)$$

Choose :

$$(u)_\alpha = \begin{cases} u & \text{if } |u| \leq \alpha, \\ 0 & \text{if } |u| > \alpha. \end{cases}$$

Then  $u \in W_0^{1,p}(\Omega, \omega, \mathbb{R}^m)$  because  $u \in W_0^{1,p}(\Omega, \omega, \mathbb{R}^m)$  and by Hardy-Type inequality:

$$\begin{aligned} \sum_j \int_{|u| \leq \alpha} \gamma_j |u_j|^q dx &= \sum_j \int_{|u| \leq \alpha} \gamma_j |(u_\alpha)_j|^q dx \\ &\leq c \cdot \left( \sum_{i,j} \int_{\Omega} \omega_{ij} |D_{ij} u_\alpha|^p dx \right)^{\frac{q}{p}} \\ &\leq c \cdot \left( \sum_{i,j} \int_{|u| \leq \alpha} \omega_{ij} |D_{ij} u_\alpha|^p dx \right)^{\frac{q}{p}} \end{aligned}$$

By (4.4)

$$\begin{aligned} c_2 \cdot \left( \sum_{ij} \int_{|u| \leq \alpha} \omega_{ij} |D_{ij} u|^p dx \right) &\leq \alpha \|f\|_{L^1(\Omega, \mathbb{R}^m)} + \alpha \sum_j \left( \int_{|u| \leq \alpha} \omega_{0j} \right)^{\frac{1}{p}} (\|b_1\|_{L^{p'}} + c') \\ &\quad + \|\lambda_2\|_{L^1(\Omega)} + c \|\lambda_3\|_{L(\frac{p}{\theta})}' \cdot \left( \sum_{i,j} \int_{|u| \leq \alpha} \omega_{ij} |D_{ij} u|^p dx \right)^{\frac{\theta q}{p^2}} \end{aligned}$$

and  $\frac{\theta q}{p^2} < 1$ . Then

$$\left( \sum_{ij} \int_{|u| \leq \alpha} \omega_{ij} |D_{ij} u|^p dx \right) \leq c \cdot \left( \alpha \|f\|_{L^1(\Omega, \mathbb{R}^m)} + \alpha \sum_j \left( \int_{|u| \leq \alpha} \omega_{0j} \right)^{\frac{1}{p}} (\|b_1\|_{L^{p'}} + c') + \|\lambda_2\|_{L^1(\Omega)} \right) \leq M\alpha + L, \quad (4.5)$$

with  $L = L(c, \|\lambda_2\|_{L^1(\Omega)}, c \|\lambda_3\|_{L(\frac{p}{\theta})}')$  and  $M = M(c_1, c_2 \|\lambda_3\|, \|f\|_{L^1(\Omega, \mathbb{R}^m)})$ , we choose:

$u^\alpha = \min(|u|, \alpha)$ , then by  $|D|u| \leq |Du|$

$$\begin{aligned} \int_{\Omega} |Du^\alpha|^{p_s} dx &= \int_{|u| \leq \alpha} |D|u||^{p_s} dx + 0 \leq \int_{|u| \leq \alpha} |Du|^{p_s} dx = \int_{\Omega} |Du_\alpha|^{p_s} dx \\ &\leq \left( \sum_{i,j} \int_{\Omega} \omega_{ij} |D_{ij} u_\alpha|^p dx \right)^{\frac{p_s}{p}} = \left( \sum_{i,j} \int_{|u| \leq \alpha} \omega_{ij} |D_{ij} u|^p dx \right)^{\frac{p_s}{p}} \end{aligned} \quad \text{And by (4.5), and } p_s \leq p, \text{ we have:}$$

$$\int_{\Omega} |Du^\alpha|^{p_s} dx \leq c \cdot (\alpha \|f\|_{L^1(\Omega, \mathbb{R}^m)} + \alpha \sum_j \left( \int_{|u| \leq \alpha} \omega_{0j} \right)^{\frac{1}{p}} (\|b_1\|_{L^{p'}} + c') + \|\lambda_2\|_{L^1(\Omega)}. \quad (4.6)$$

By (1.5) and (4.6), we have:

$$\int_{\Omega} |u^{\alpha}|^{p_s^*} dx \leq c. \left( \int_{\Omega} |Du^{\alpha}|^{p_s} dx \right)^{\frac{p_s^*}{p}} \leq c. \left( \alpha \|f\|_{L^1(\Omega; \mathbb{R}^m)} + \alpha \sum_j \left( \int_{|u| \leq \alpha} \omega_{0j} \right)^{\frac{1}{p}} (\|b_1\|_{L^{p'}} + c') + \|\lambda_2\|_{L^1(\Omega)} \right)^{\frac{p_s^*}{p}} \quad (4.7)$$

Then:

$$\begin{aligned} \lambda_{|u|}(\alpha) &= \alpha^{-p_s^*} \int_{|u| > \alpha} \alpha^{p_s^*} dx \leq \alpha^{-p_s^*} \int_{|u| > \alpha} |u^{\alpha}|^{p_s^*} dx \\ &\leq c. \alpha^{-p_s^*} \left( \alpha \|f\|_{L^1(\Omega; \mathbb{R}^m)} + \alpha \sum_j \left( \int_{|u| \leq \alpha} \omega_{0j} \right)^{\frac{1}{p}} (\|b_1\|_{L^{p'}} + c') + \|\lambda_2\|_{L^1(\Omega)} \right)^{\frac{p_s^*}{p}} \end{aligned} \quad (4.8)$$

and we continue in the same way as in a case that is non-degenerated [2] by replacing  $p$  by  $p_s$  as well as

$$\begin{aligned} \|u\|_{L^{t_{p_s}^*, \infty}(\Omega, \mathbb{R}^m)}^* &= \sup_{\alpha > 0} \alpha |\lambda_{|u|}(\alpha)|^{\frac{1}{t_{p_s}^*}} \\ &\leq |\Omega| + \sup_{\alpha > 1} \alpha |\lambda_{|u|}(\alpha)|^{\frac{1}{t_{p_s}^*}} \\ &\leq |\Omega| + c. (\|f\|_{L^1(\Omega; \mathbb{R}^m)}^{\frac{1}{p_s-1}}, \|b_1\|_{L^1}, \|\lambda_2\|_{L^1(\Omega)}^{\frac{1}{p_s-1}}) \end{aligned}$$

i-e:

$$\|u\|_{L^{t_{p_s}^*, \infty}(\Omega, \mathbb{R}^m)}^* \leq c. \left( |\Omega|, \|b_1\|_{L^1}, \|\lambda_2\|_{L^1(\Omega)}, \|\lambda_3\|_{L^{\left(\frac{p}{p_s}\right)'(\Omega)}}, c_2, \|f\|_{L^1(\Omega; \mathbb{R}^m)} \right), \quad (4.9)$$

on the other hen, by using ( $p_s \leq p$ ) and thinks to 1.5, we obtain:

$$\begin{aligned} \lambda_{|Du|}(s) &\leq s^{-p_s} \int_{|u| \leq \alpha} |Du|^{p_s} dx + \lambda_{|u|}(\alpha) \\ &= s^{-p_s} \int_{|u| \leq \alpha} |Du_{\alpha}|^{p_s} dx + \lambda_{|u|}(\alpha) \\ &\leq s^{-p_s} \left( \sum_{i,j} \int_{|u| \leq \alpha} \omega_{ij} |D_{ij} u_{\alpha}|^p dx \right) + \lambda_{|u|}(\alpha) \\ &\leq s^{-p_s} \left( \sum_{i,j} \int_{|u| \leq \alpha} \omega_{ij} |D_{ij} u|^p dx \right) + \lambda_{|u|}(\alpha) \end{aligned}$$

By 4.5 and 4.8:

$$\lambda_{|Du|}(s) \leq c. \left( \max\left(\frac{\alpha}{s^{p_s}}, \frac{1}{s^{p_s}}\right) + \max(\alpha^{-p_s^*}, \alpha^{\frac{p_s^*}{p_s} - p_s^*}) \right)$$

or  $-t_{p_s}^* = \frac{p_s^*}{p_s} - p_s^*$ , so as in [3]

$$\|Du\|_{L^{t_{p_s}^*, \infty}(\Omega, M^{m \times n})}^* \leq c. \left( |\Omega|, \|\lambda_2\|_{L^1(\Omega)}, \|\lambda_3\|_{L^{\left(\frac{p}{p_s}\right)'(\Omega)}}, c_2, \|b_1\|_{L^1}, \|f\|_{L^1(\Omega; \mathbb{R}^m)} \right). \quad (4.10)$$

From (4.5)-(4.9) and (4.10), we obtain the result of the proposition (4.1) in case i). ii)-Suppose the condition (of l'angle in  $H_3$ ): Let  $S_{\alpha}(y) = (T_{\alpha}(y_1); T_{\alpha}(y_2); \dots; T_{\alpha}(y_m))$ ,  $y \in \mathbb{R}^m$ , the cubic truncation, we have  $Ds_{\alpha}(y) = Id$  if  $|y|_{max} = \max_{1 \leq i \leq m} |y_i| \leq \alpha$ , in the same way as in i)- by testing  $S_{\alpha}(u)$  in (4.3).

Then  $\int_{\Omega} \sigma(x, u, Du) : D(S_{\alpha}(u)) dx = \int_{\Omega} f.S_{\alpha}(u) dx + \int_{\Omega} g.S_{\alpha}(u) dx$

or

$$\begin{aligned} \int_{\Omega} \sigma(x, u, Du) : D(S_{\alpha}(u)) dx &= \sum_{i=1}^m \int_{|u_i| \leq \alpha} \sigma_i(x, u, Du) : Du_i dx \\ &\geq \int_{|u| = \max_{1 \leq j \leq m} (|u_j|)} \sum_{i=1}^m \sigma_i(x, u, Du) : Du_i dx \end{aligned}$$

and like  $\sum_{i=1}^m \sigma_i(x, u, Du) : Du_i dx = \sigma(x, u, Du) : Du$ . By the coercivity condition in  $(H_1)$  and the Hölder Inequality we obtain:

$$c_2 \cdot \sum_{i,j} \int_{|u| \leq \alpha} \omega_{ij} |D_{ij} u|^p dx \leq \sqrt{m} \cdot \alpha (\|f\|_{L^1(\Omega; \mathbb{R}^m)} + \alpha \sum_j (\int_{|u| \leq \alpha} \omega_{0j})^{\frac{1}{p}} (\|b_1\|_{L^{p'}} + c') + \|\lambda_2\|_{L^1(\Omega)})$$

$$+ c \cdot \|\lambda_3\|_{L(\frac{p}{\theta})'(\Omega)} \left( \sum_{j=1}^m \int_{|u| \leq \alpha} \gamma_j |u_j|^q dx \right)^{\frac{q}{p}}$$

and we continue in the same way as in i), this completes the proof of the proposition (6)  $\square$

## 5. A Div-Curl Inequality

The result of this section is the key ingredient for the proof that one can pass to the limit in the equation (4.1) for the solution  $\{u_k\}_{k \in \mathbb{N}}$  of approximating problems. Since it is independent of the differential equation we state it a more general form using only the hypotheses (5.1)-(5.8) below:

$$\sigma; \tau : \Omega \times \mathbb{R}^m \times M^{m \times n} \longrightarrow M^{m \times n}, \quad (5.1)$$

is a Carathéodory function.

$$\sigma \text{ and } \tau \text{ satisfying one of the following conditions:} \quad (5.2)$$

- (i)  $\sigma(x, u, F) : MF \geq 0, \tau(x, u, F) : MF \geq 0 ; M = Id - b \otimes b \in M^{m \times n}$ , with  $|b| \leq 1$ .
- (ii)  $\sigma_j(x; u; F) : F_j \geq 0$ , and  $\tau_j(x, u, F) : F_j \geq 0 ; 1 \leq j \leq m$ ,  $\sigma_j, \tau_j$  and  $F_j$  is the  $j^{eme}$  columns of  $\sigma, \tau, F$ .

$$u_k \in W^{1;1}(\Omega, \mathbb{R}^m) \text{ and there exists an } s \geq 0 \text{ such that } \int_{\Omega} |Du_k|^s dx \leq c \text{ uniformly in } k \quad (5.3)$$

$$\text{The sequence } \sigma_k(x) = \sigma(x, u_k, Du_k) \text{ is equiintegrable.} \quad (5.4)$$

$$\text{The sequence } u_k \text{ converges in measure to some function } u, \text{ and } u \text{ is almost everywhere approximately differentiable.} \quad (5.5)$$

$$\text{The sequence } f_k = -div(\sigma_k + \tau_k) - \mu \text{ is bounded in } L^1(\Omega, \mathbb{R}^m). \quad (5.6)$$

$$D_{ij} u_k \in L^r_{loc}(\Omega, \omega_{ij}, M^{m \times n}) \text{ and } (\sigma_k + \tau_k) \in L^r_{loc}(\Omega, \omega^*, M^{m \times n}), \text{ for some} \quad (5.7)$$

$$1 \leq r < \infty \text{ and } 1 \leq i \leq n, 1 \leq j \leq m.$$

$$\text{The sequence } \tau_k(x) = \tau[x](x, u, Du_k) \text{ converges to weakly to 0 in } L^1(\Omega, M^{m \times n}). \quad (5.8)$$

**Lemma 5.1** *Suppose (5.1)-(5.8). Then (after passage to a subsequence) the sequence  $\sigma_k$  converges weakly in  $L^1(\Omega, M^{m \times n})$  and the weak limit  $\bar{\sigma}$  is given by  $\bar{\sigma}(x) = \langle \nu_x; \sigma(x, u(x), \cdot) \rangle$ . Moreover the following inequality holds:*

$$\int_{M^{m \times n}} \sigma(x, u(x), \lambda) : \lambda d\nu_x(\lambda) \leq \bar{\sigma}(x) : apDu(x) \text{ for a.e. } x \in \Omega. \quad (5.9)$$

**Proof** See [3]  $\square$

## 6. Passage to the Limit

**Proposition 6.1** *Suppose that the sequence  $(u_k)_{k \in \mathbb{N}}$  satisfies the hypotheses (5.1)-(5.7),  $(H_2)$  and that the Young measure  $\nu$  generated by the sequence  $(Du_k)_{k \in \mathbb{N}}$  satisfies: a)-c) and d)- in lemma (3.3). Then the sequence  $(\sigma_k)$  is weakly converge in  $L^1(\Omega, M^{m \times n})$ , with  $\bar{\sigma}$  is the limit and  $\bar{\sigma}(x) = \langle \nu_x, u(x), apDu(x) \rangle$ . If in  $H_2$  b)- c)-or d)-holds,  $\sigma(x, u_k, Du_k) \rightarrow \sigma(x, u, apDu)$  strongly in  $L^1(\Omega, M^{m \times n})$ . In the cases (c) and (d) it follows addition that  $Du_k \rightarrow apDu$  in measure.*

**Proof** See [3].

**Proof of the theorem 2.1**

For using the results of proposition (6.1): we assume that (5.1)-(5.7) and the Young measure  $\nu_x$  generated by the sequence  $Du_k$  satisfies (i), (ii) and (iii) in Lemma(3.3), for the approximate systems (4.1)-(4.2). By the proposition 6.1, with  $u_k \in W_0^{1,p}(\Omega, \omega, \mathbb{R}^m)$ , we have:  $\|u_k\|_{L^{t_{ps}^*, \infty}(\Omega, \mathbb{R}^m)} \leq$

$$c \left( |\Omega|, \|\lambda_2\|_{L^1(\Omega)}, \|\lambda_1\|_{L^{p'}}, \|\lambda_3\|_{L^{(\frac{p}{\theta})'}(\Omega)}, c_2, \|\mu\|_{M(\Omega, \omega^*, \mathbb{R}^m)} \right), \text{ and}$$

$$\sum_{i,j} \int_{|u_k| \leq \alpha} \omega_{ij} |D_{ij} u_k|^p dx \leq M\alpha + L < \infty. \quad (6.1)$$

By  $L^{t_{ps}^*, \infty}(\Omega, \mathbb{R}^m) \hookrightarrow L^p(\Omega, \mathbb{R}^m)$  for all  $1 < p < t_{ps}^*$ , then

$$\|u_k\|_{L^p(\Omega, \mathbb{R}^m)} \leq c < \infty. \quad (6.2)$$

Now

- (5.1) is  $(H_0)$
- (5.2) is  $(H_3)$
- (5.3):  $u_k \in W_0^{1,p}(\Omega, \omega, \mathbb{R}^m) \hookrightarrow W_0^{1,p_s}(\Omega, \mathbb{R}^m)$  with  $p_s > 1$ , then  $u_k \in W^{1,1}(\Omega, \mathbb{R}^m)$ . Moreover, by the proposition

$$\|Du_k\|_{L^{t_{ps}^*, \infty}(\Omega, \mathbb{R}^m)} \leq c \left( |\Omega|, \|\lambda_2\|_{L^1(\Omega)}, \lambda_1 \|_{L^{p'}}, \|\lambda_3\|_{L^{(\frac{p}{\theta})'}(\Omega)}, c_2, \|\mu\|_{M(\Omega, \omega^*, \mathbb{R}^m)} \right)$$

hence

$$\|Du_k\|_{L^s(\Omega, M^{m \times n})} \leq c < \infty, \quad \forall 1 < s < t_{ps}$$

with  $\sup_{k \in \mathbb{N}} \int_{\Omega} |Du_k|^s dx < \infty$ .

- (5.4): Let  $A$  a measurable in  $\Omega$ , by  $(H_1)$  and Hölder we have

$$\int_A |\sigma(x, u_k, Du_k)| dx \leq c \left( \sum_{r,s} \int_{\Omega} \omega_{rs} dx \right)^{\frac{1}{p}} \cdot \left\{ \|\lambda_1\|_{L^{p'}(\Omega)} + \left( \sum_{j=1}^m \int_{\Omega} \gamma_j |u_k|_j^q dx \right)^{\frac{1}{p}} + \left( \sum_{i,j} \int_{\Omega} \omega_{ij} |D_{ij} u_k|^p dx \right)^{\frac{1}{p}} \right\},$$

and with (1.5) and (1.3):

$$\int_A |\sigma(x, u_k, Du_k)| dx \leq c \left( \sum_{r,s} \|\omega_{rs}\|_{L^1_{loc}(\Omega)} \right) \times \left\{ \|\lambda_1\|_{L^{p'}(\Omega)} + \|u_k\|_{1,p,\omega}^{\frac{q}{pp'}} + \|u_k\|_{1,p,\omega}^{\frac{p}{p'}} \right\} < \infty.$$

- (5.5): By (6.1) and (6.2) and Lemma (3.2).
- (5.6):  $\|f_k\|_{L^1(\Omega, \mathbb{R}^m)} \leq \|\mu\|_{M(\Omega, \omega^*, \mathbb{R}^m)}$ . And By  $(G_1)$  we have

$$\int_{\Omega} |g(x, u_k, Du_k)| dx \leq \infty \text{ (by Holder and } G_1)$$

- (5.7):  $\forall \varepsilon > 0$  and  $x_0 \in \Omega$   $\int_{B(x_0, \varepsilon)} |D_{ij}u_k|^p \omega_{ij} dx \leq \|u_k\|_{1,p,\omega}^p < \infty$ . And by (H<sub>3</sub>)

$$\begin{aligned} \int_{B(x_0, \varepsilon)} |\sigma_{rs}(x, u_k, Du_k)|^{p'} \omega_{rs}^* dx &= \int_{B(x_0, \varepsilon)} |\sigma_{rs}(x, u_k, Du_k)|^{p'} \omega_{rs}^{1-p'} dx \\ &\leq c \int_{B(x_0, \varepsilon)} w^{1-p'+\frac{p'}{p}} \left[ |\lambda_1|^{p'} + \sum_{j=1}^m \gamma_j |(u_k)_j|^q + \sum_{i,j} \omega_{ij} |D_{ij}u_k|^p \right] dx \\ &\leq c \left( \|\lambda_1\|_{L^{p'}(\Omega)}^{p'} + \|u_k\|_{1,p,\omega}^{\frac{q}{p}} + \|u_k\|_{1,p,\omega}^p \right) < \infty. \end{aligned}$$

Then, by the proposition (6.1)  $\sigma(x, u_k, Du_k) \rightarrow \sigma(x, u, apDu)$  in  $L^1(\Omega, M^{m \times n})$  and  $\forall \varphi \in C_0^\infty(\Omega, \mathbb{R}^m)$ ;  $D\varphi \in L^\infty(\Omega, M^{m \times n})$  hence:

$$\int_{\Omega} \sigma(x, u_k, Du_k) : D\varphi dx \rightarrow \int_{\Omega} \sigma(x, u, apDu) : D\varphi dx$$

i.e:

$$-div\sigma(x, u_k, apDu_k) \rightarrow -div\sigma(x, u, apDu)$$

In the sense of distributions. Moreover, since  $u_k \rightarrow u$  in measure, it follows that (at least for a subsequence)  $u_k \rightarrow u$  almost everywhere and hence that  $g(x, u_k, Du_k) \rightarrow g(x, u, Du)$  almost everywhere from the continuity condition ( $G_0$ ). Since  $g(x, u_k, Du_k)$  is equi-integrable by the growth condition in  $G_1$  and the uniform bounded 6.1- 6.2, we may infer that  $g(x, u_k, Du_k) \rightarrow g(x, u, Du)$  in  $L^1(\Omega, \mathbb{R}^m)$  by the Vitali's converge theorem. On the other hand  $f_k \xrightarrow{*} \mu$  in  $L^1(\Omega, \mathbb{R}^m)$ . Then

$$\begin{aligned} \int_{\Omega} f_k \cdot \varphi dx &\rightarrow \int_{\Omega} \mu \cdot \varphi dx \quad \forall \varphi \in C_0^\infty(\Omega, \mathbb{R}^m) \text{ so } \mu \text{ is the solution in } W_0^{1,p}(\Omega, \omega, \mathbb{R}^m) \text{ of the system:} \\ -div\sigma(x, u, apDu) &= \mu \quad \text{in } \Omega \\ u &= 0 \text{ on } \partial\Omega \end{aligned}$$

to show the estimation (2.1), we take the function  $\eta$  in  $C_0^1(B(0, 2\alpha), \mathbb{R}^m)$ ;  $\eta = Id$  in  $B(0, \alpha)$  and  $|D\eta| \leq c$ , then:

$$\begin{aligned} \sum_{i,j} \int_{\Omega} \omega_{ij} |D_{ij}\eta(u_k)|^p dx &= \sum_{i,j} \int_{\Omega} \omega_{ij} |(D_{ij}\eta)(u_k)|^p |Du_k|^p dx \\ &\leq c^p \cdot \sum_{i,j} \int_{|u_k| \leq \alpha} \omega_{ij} |D_{ij}u_k| dx + c \cdot \sum_{i,j} \int_{|u_k| \leq 2\alpha} \omega_{ij} |D_{ij}u_k|^p dx \\ &\leq c \cdot c(\alpha) + c \cdot c(2\alpha) < \infty, \end{aligned}$$

thanks to (6.1).

Now, we have  $\eta(u_k) \rightarrow \eta(u)$ , for every  $x \in \Omega$  because  $\eta$  is  $C^\infty$ . Then  $\eta(u_k) \rightarrow \eta(u)$ , in  $W_0^{1,p}(\Omega, \omega, \mathbb{R}^m)$ , and  $apDu = apD(\eta \circ u)$  on  $\{|u| < \alpha\}$ . Hence,

$$\begin{aligned} \sum_{i,j} \int_{\Omega} \omega_{ij} |D_{ij}(\eta \circ u)|^p dx &\leq \liminf_{k \rightarrow \infty} \int_{\Omega} \omega |D(\eta \circ u_k)|^p dx \\ &\leq \liminf_{k \rightarrow \infty} \sum_{i,j} \int_{|u_k| \leq 2\alpha} |D_{ij}\eta(u_k)|^p |D_{ij}u_k| \omega_{ij} dx \\ &\leq \leq c \cdot \liminf_{k \rightarrow \infty} \int_{|u_k| \leq 2\alpha} \omega_{ij} |D_{ij}u_k|^p dx \\ &\leq c \cdot c(2\alpha) < \infty. \end{aligned}$$

Then:

$$\sum_{i,j} \int_{|u| \leq \alpha} \omega_{ij} |apDu|^p dx = \sum_{i,j} \int_{|u_k| \leq 2\alpha} \omega_{ij} |D(\eta \circ u)|^p dx < \infty,$$

in the same as in the proof of the proposition (6) by replacing  $u_k$  by  $u$  and  $f_k$  by  $\mu$ , we obtain the estimation (2.1) and this completes the proof of the theorem 2.1  $\square$

**Case:**  $0 < \theta < \frac{n(p_s-1)}{n-1}$  (the general case) The idea is to consider the regularized problems:

$$-div\phi_\varepsilon(x, u_\varepsilon, Du_\varepsilon) = \mu \text{ in } \Omega, \quad (6.3)$$

$$u_\varepsilon = 0 \text{ on } \partial\Omega \quad (6.4)$$

With

$$\phi_{\varepsilon,r,s}(x, u, F) = \sigma_{rs}(x, u, F) + \varepsilon\beta \left( \sum_{ij} \omega_{ij}^{\frac{1}{p'}} |F_{ij}|^{s-2} \right) \omega_{rs}^{\frac{1}{p}} F_{rs},$$

$\forall 1 \leq r \leq n, \forall 1 \leq s \leq m$  with  $s > n + 1$ , and  $\varepsilon < \frac{1}{2}$ , we have  $p < s$ , then  $s' < p'$ , and

$(\frac{s}{\theta})' < (\frac{p}{\theta})'$ . Moreover  $\exists c > 0$  which doesn't depend on  $p, s$ , such that  $\omega_{rs}^{\frac{1}{p}} \leq c \cdot \omega_{rs}^{\frac{1}{s}}$

$\forall 1 \leq r \leq n$  and  $1 \leq s \leq m$ .

By  $(H_1)$  for  $\sigma$ , we obtain

$$\begin{aligned} |\phi_{\varepsilon,r,s}(x, u, F)| &\leq \beta' \cdot |\omega_{rs}|^{\frac{1}{p}} \left[ \lambda_1 + \sum_{j=1}^m \gamma_j^{\frac{1}{p'}} |u_j|^{\frac{q}{p'}} + \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |F_{ij}|^\theta \right] \\ &\quad + \varepsilon\beta \omega_{rs}^{\frac{1}{p}} \left( \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |F_{ij}|^{s-1} \right) \left( \theta < \frac{n(p_s-1)}{n-1} < n(s-1) \right) \\ &\leq \leq \beta' \omega_{rs}^{\frac{1}{p}} \left[ \lambda_1 + \sum_{j=1}^m \gamma_j^{\frac{1}{p'}} |u_j|^{\frac{q}{p'}} + \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |F_{ij}|^{s-1} \right]. \end{aligned}$$

And  $p < s$ , then  $\frac{1}{p'} < \frac{1}{s'}$  and like  $\omega_{rs}^{\frac{1}{p}} \leq c \cdot \omega_{rs}^{\frac{1}{s}}$ , then:

$$|\phi_{\varepsilon,r,s}(x, u, F)| \leq \beta' \cdot |\omega_{rs}|^{\frac{1}{s}} \left[ \lambda_1 + \sum_{j=1}^m \gamma_j^{\frac{1}{s'}} |u_j|^{\frac{q}{s'}} + \sum_{ij} \omega_{ij}^{\frac{1}{s'}} |F_{ij}|^{s-1} \right], \text{ and by } (H_3), \text{ we conclude that}$$

$$\phi_\varepsilon(x, u, F) : F = \sigma(x, u, F) : F + \varepsilon \sum_{i,j,r,s} \omega_{ij}^{\frac{1}{p'}} \omega_{rs}^{\frac{1}{p}} |F_{ij}|^{s-2} F_{ij} \cdot F_{rs}$$

$$\geq -\lambda_2 - \sum_{j=1}^m \lambda_3 \gamma_j^{\frac{q}{s}} \cdot |u_j|^{\frac{q\alpha}{s}} + \varepsilon \sum_{ij} \omega_{ij} |F_{ij}|^s.$$

On the other hand,  $0 < \alpha < p - 1 < s - 1, 1 < q < \frac{p^2}{\alpha} < \frac{s^2}{\alpha'}$ ,  $\lambda_1 \in L^{p'}(\Omega) \hookrightarrow L^{s'}(\Omega)$ , and  $\lambda_3 \in L^{(\frac{p}{\alpha})'}(\Omega) \hookrightarrow L^{(\frac{s}{\alpha})'}(\Omega)$  and as  $\sigma_\varepsilon$  verifies the conditions of the structures (of l'angle and sign), the strict monotony, the s-quasi monotonous with regard to  $F$  is a  $C^1$  monotony in relation with  $F$  or accepting a

convex potential because:  $F \rightarrow \varepsilon\beta \left( \sum_{ij} \omega_{ij}^{\frac{1}{p'}} |F_{ij}|^{s-2} \right) \omega_{rs}^{\frac{1}{p}} F_{rs}$  verify them as well, hence  $\sigma_\varepsilon$  verifies the

hypotheses  $(H_0) - (H_5)$ , for the regularized problems (6.3)-(6.4), thus for the previous case,  $\theta = s - 1$ , of theorem 2.1, there exists a solution,  $u_\varepsilon \in W_0^{1,s}(\Omega, \omega, \mathbb{R}^m)$ , of the system (6.3)-(6.4). Now showing that the conditions: i), ii) and iii), of lemma (3.3), and the hypotheses (5.1)-(5.8) of the div-curl inequality are verified for  $u_\varepsilon$  with order  $s$  in the place of  $p$ .

We suppose the condition of l'angle verifying that  $\phi_\varepsilon$  by testing,  $T_\alpha(u_\varepsilon) \alpha \succ 0$  in (5.3)-(5.4), we get:

$$\int_\Omega \phi_\varepsilon(x, u_\varepsilon, Du_\varepsilon) : DT_\alpha(u_\varepsilon) dx = \int_\Omega f \cdot T_\alpha(u_\varepsilon) dx, \text{ so}$$

$$\begin{aligned} &\int_{|u_\varepsilon| \leq \alpha} \sigma(x, u_\varepsilon, Du_\varepsilon) : Du_\varepsilon dx + \int_{|u_\varepsilon| > \alpha} \frac{\alpha}{|u_\varepsilon|} \sigma_\varepsilon(x, u_\varepsilon, Du_\varepsilon) : \left( Id - \frac{u_\varepsilon}{|u_\varepsilon|} \otimes \frac{u_\varepsilon}{|u_\varepsilon|} \right) Du_\varepsilon dx \\ &\quad + \varepsilon\beta \int_{|u_\varepsilon| \leq \alpha} \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |D_{ij}u_\varepsilon|^{s-2} \sum_{r,s} \omega_{rs}^{\frac{1}{p}} |D_{rs}u_\varepsilon|^2 dx \\ &\quad + \varepsilon\beta \int_{|u_\varepsilon| > \alpha} \sum_{ij} \omega_{ij}^{\frac{1}{p'}} |D_{ij}u_\varepsilon|^{s-2} \sum_{r,s} \omega_{rs} D_{rs}u_\varepsilon \left( Id - \frac{u_\varepsilon}{|u_\varepsilon|} \otimes \frac{u_\varepsilon}{|u_\varepsilon|} \right) \end{aligned}$$

$$\leq \alpha \cdot \|f\|_{L^1(\Omega, \mathbb{R}^m)}.$$

since

$$\sum_{rs} |D_{rs} u_\varepsilon|^{s-2} D_{rs} u_\varepsilon \left( Id - \frac{\alpha}{|u_\varepsilon|} \left( \frac{u_\varepsilon}{|u_\varepsilon|} \otimes \frac{u_\varepsilon}{|u_\varepsilon|} \right) \right) \geq 0$$

so

$$\int_{|u_\varepsilon| \leq \alpha} \sigma(x, u_\varepsilon, Du_\varepsilon) : Du_\varepsilon dx \leq \alpha \|f\|_{L^1(\Omega, \mathbb{R}^m)}.$$

And by the coercivity condition of  $\sigma$  in  $(H_1)$  and Hölder inequality, we get as in the proof of the proposition

$$\sum_{ij} \int_{|u_\varepsilon| \leq \alpha} \omega_{ij} |D_{ij} u_\varepsilon|^p dx \leq M' \alpha + L', \quad (6.5)$$

And the following a priori estimation:

$$\|u_\varepsilon\|_{L^{t_{ps}^*}(\Omega, \mathbb{R}^m)}^* + \|Du_\varepsilon\|_{L^{t_{ps}}(\Omega, \mathbb{R}^{m \times n})}^* < c < \infty, \quad (6.6)$$

and by the injection  $L^{\beta', \infty} \hookrightarrow L^{\alpha'}$ ,  $\forall 0 < \alpha' < \beta'$ , then  $\forall, 0 < r < t_{ps}^*, \forall 0 < p < t_{ps}$

$$\|u_\varepsilon\|_{L^r(\Omega, \mathbb{R}^m)} + \|Du_\varepsilon\|_{L^p(\Omega, \mathbb{M}^{m \times n})} + \|Du_\varepsilon\|_{L^{t_{ps}}(\Omega, \mathbb{M}^{m \times n})}^* < \infty. \quad (6.7)$$

We suppose that the condition of the sign is verify.

As in the same way in the proof of the proposition (6), we test  $S_\alpha(u_\varepsilon)$  in (6.3)-(6.4), we obtain (6.5) and (6.7).

Starting with verifying that i), ii) et iii) of lemma (3.3) and the hypotheses (5.1) and (5.7) for  $\sigma_\varepsilon$ . By (6.5) and (6.7), the points i), ii) et iii) are a direct consequence of lemma (3.2) and lemma (3.3). On the other hand:

-(5.1): for  $\sigma$  is  $(H_0)$  and  $\tau_{rs}(x, u, F) = \varepsilon \beta \left( \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |F_{ij}|^{s-2} \right) \cdot \omega_{rs}^{\frac{1}{p}} F_{rs}$  is a Carathéodory function, because  $x \mapsto \omega_{ij}(x)$ , is measurable, so  $\sigma_\varepsilon$  is a Cathéodory function.

-(5.2)

$$(i) \quad \phi_\varepsilon(x, u, F) : MF = \sigma(x, u, F) : MF + \left( \sum_{rs} \left( \varepsilon \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |F_{ij}|^{s-2} \right) \omega_{rs}^{\frac{1}{p}} F_{rs} \right) (MF)_{rs} \geq 0,$$

with  $M = Id - a \otimes a$  and  $|a| \leq 1$ .

(ii)

$$\begin{aligned} \phi_{rs}(x, u, F) \cdot F_j &= \sigma_j(x, u, F) : F_j + \tau_j(x, u, F) \cdot F_j \\ &= \sigma_j(x, u, F) : F_j + \sum_{l=1}^m \varepsilon \beta \left( \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |F_{ij}|^{s-1} \right) \cdot \omega_{lj}^{\frac{1}{p}} |F_{lj}|^2 \geq 0, \end{aligned}$$

$\forall 1 \leq j \leq m$ .

-(5,3):  $u_\varepsilon \in W_0^{1,s}(\Omega, \omega, \mathbb{R}^m) \hookrightarrow W_0^{1,s_s}(\Omega, \mathbb{R}^m)$ ,  $s_s > 1$ , so  $u_\varepsilon \in W^{1,1}(\Omega, \mathbb{R}^m)$ , and by (6.7)

$\sup_{\varepsilon > 0} \int_{\Omega} |Du_\varepsilon|^p dx < \infty$ ,  $\forall, 0 < p < t_{ps}$ .

(4.5):  $\sigma(x, u_\varepsilon, Du_\varepsilon)$  is equi-integrable as previously  $\forall \Omega' \subset \Omega$ , measurable, we have:

$$\begin{aligned} &\int_{\Omega'} \left| \sum_{i,j} \left( \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\varepsilon|^{s-2} \right) \omega_{rs}^{\frac{1}{p}} D_{rs} u_\varepsilon \right| dx \\ &\leq \left( \sum_{i,j} \int_{\Omega'} \omega_{ij} |D_{ij} u_\varepsilon|^{s-1} dx \right) \\ &\leq c \cdot \sum_{ij} \int_{\Omega'} \omega_{ij} |D_{ij} u_\varepsilon|^s dx \leq c \cdot \|u_\varepsilon\|_{1,s,w}^s. \end{aligned}$$

-(5.5): by (6.7) and the lemma (3.2).

-(5.6): by (6.3),  $-div(\sigma_l + \tau_k) - \mu = 0$ , with  $\mu \in M(\Omega, \mathbb{R}^m)$  is bounded in  $L^1(\Omega, \mathbb{R}^m)$ .

-(5.7):  $\forall \varepsilon > 0$  and  $x_0 \in \Omega$ , by the growth condition of  $\sigma_\varepsilon$  and previously with  $s$  in the place of  $p$ ,

$$\int_{B(x_{\bar{A}}, \varepsilon)} |\sigma_\varepsilon(x, u_\varepsilon, Du_\varepsilon)|^s \omega_{rs}^* dx < \infty$$

and

$$-(5.8): \int_{B(x_{\bar{A}}, \varepsilon)} |D_{ij} u_\varepsilon|^s \omega_{rs} dx < \|u_\varepsilon\|_{1,s,w}^\varepsilon < \infty.$$

Testing that  $u_\varepsilon$  in (6.3)-(6.4)

$$\begin{aligned} \varepsilon \beta \int_{\Omega} \left( \sum_{ij} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\varepsilon|^{s-2} \right) \left( \sum_{rs} \omega_{rs}^{\frac{1}{p}} |D_{rs} u_\varepsilon|^2 \right) dx \\ \leq \|u_\varepsilon\|_{L^\infty(\Omega, \mathbb{R}^m)} \|\mu\|_{M(\Omega, \omega^*, \mathbb{R}^m)} \end{aligned} \quad (6.8)$$

We have  $W_0^{1,s}(\Omega, w, \mathbb{R}^m) \hookrightarrow W_0^{1,s_s}(\Omega, \mathbb{R}^m) \hookrightarrow L^\infty(\Omega, \mathbb{R}^m)$ . Then

$$\begin{aligned} \|u_\varepsilon\|_{L^\infty(\Omega, \mathbb{R}^m)} &\leq c \cdot \left( \sum_{ij} \int_{\Omega} \omega_{ij} |D_{ij} u_\varepsilon|^s dx \right)^{\frac{1}{s}} \\ &\leq c \cdot \left( \sum_{ij} \int_{\Omega} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\varepsilon|^{s-2} \omega_{ij}^{\frac{1}{p}} |D_{ij} u_\varepsilon|^2 dx \right)^{\frac{1}{s}} \\ &\leq c \left( \int_{\Omega} \left( \sum_{ij} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\varepsilon|^{s-2} \right) \cdot \left( \sum_{rs} \omega_{rs}^{\frac{1}{p}} |D_{rs} u_\varepsilon|^2 dx \right)^{\frac{1}{s}} \right)^{\frac{1}{s}}. \end{aligned} \quad (6.9)$$

Thanks to (6.8) and (6.9), we have

$$\begin{aligned} &\int_{\Omega} \sum_{ij} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\varepsilon|^{s-2} \sum_{rs} \omega_{rs}^{\frac{1}{p}} |D_{rs} u_\varepsilon|^2 dx \\ &\leq \frac{c \cdot \|\mu\|_{M(\Omega, \omega^*, \mathbb{R}^m)}}{\varepsilon} \left( \int_{\Omega} \left( \sum_{ij} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\varepsilon|^{s-2} \right) \cdot \left( \sum_{rs} \omega_{rs}^{\frac{1}{p}} |D_{rs} u_\varepsilon|^2 dx \right) \right) \text{ So:} \\ &\left( \int_{\Omega} \left( \sum_{ij} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\varepsilon|^{s-2} \right) \cdot \left( \sum_{rs} \omega_{rs}^{\frac{1}{p}} |D_{rs} u_\varepsilon|^2 dx \right) \right)^{\frac{s-1}{s}} \leq \frac{c \|\mu\|_M}{\varepsilon}, \end{aligned}$$

which mean that

$$\left( \int_{\Omega} \left( \sum_{ij} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\varepsilon|^{s-2} \right) \cdot \left( \sum_{rs} \omega_{rs}^{\frac{1}{p}} |D_{rs} u_\varepsilon|^2 dx \right) \right)^{\frac{1}{s}} \leq \frac{c \|\mu\|_M}{\varepsilon}, \quad (6.10)$$

and

$$\|u_\varepsilon\|_{L^\infty(\Omega, \mathbb{R}^m)} \leq c \cdot \left( \frac{c \|\mu\|_M}{\varepsilon} \right)^{\frac{1}{s-1}}. \quad (6.11)$$

On the other hand and  $\forall 1 < p < \frac{s}{s-1}$ , can write

$$\begin{aligned}
& \left\| \epsilon \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\epsilon|^{s-2} \omega_{rs}^{\frac{1}{p}} |F_{rs}| \right\|_{L^{\frac{s}{s-1}}(\Omega, M^{m \times n})} \\
& \leq \epsilon^{\frac{s}{s-1}} \left( \int_{\Omega} \left| \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\epsilon|^{s-1} \omega_{rs}^{\frac{1}{p}} \right|^{\frac{s-1}{s}} dx \right)^{\frac{s-1}{s}} \\
& \leq c \epsilon^{\frac{s}{s-1}} \left( \int_{\Omega} \left| \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\epsilon|^{s-1} \omega_{rs}^{\frac{1}{p}} \right|^{\frac{s-1}{s}} dx \right)^{\frac{s-1}{s}} \\
& \leq c \epsilon^{\frac{s}{s-1}} \left( \sum_{i,j} \int_{\Omega} \omega_{ij} |D_{ij} u_\epsilon|^{s-2} \sum_{r,s} \omega_{rs}^{(s-1)p} |D_{rs} u_\epsilon|^2 dx \right) < \infty.
\end{aligned}$$

thanks to (6.10). Now, since  $u_\epsilon \in W_0^{1,s}(\Omega, \omega, \mathbb{R}^m) \hookrightarrow W_0^{1,s_s}(\Omega, \mathbb{R}^m) \hookrightarrow W_0^{1,p_s}(\Omega, \mathbb{R}^m)$ , so by testing  $T_\alpha(u_\epsilon)$  in (6.3)-(6.4), we obtain as in the proof of the proposition (4.1)

$$\|Du_\epsilon\|_{L^{\frac{n(p_s-1)}{n-1}, \infty}(\Omega, M^{m \times n})}^* \leq c. \quad (6.12)$$

By the Hölder inequality for the exponent  $a$  with  $a$  and  $\xi$  are the solutions of systems:

$$\begin{cases} a'\xi = \tau > \frac{n(p_s-1)}{n-1} \\ a((s-1)\rho - \xi) = s \end{cases}$$

a given system accepting the solution when  $\rho < \frac{s}{s-1}$ . So

$$\begin{aligned}
& \int_{\Omega} |\epsilon \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\epsilon|^{s-1} \omega_{ij}^{\frac{1}{p}}|^\rho dx \\
& \leq c \int_{\Omega} \epsilon^\rho \left( \sum_{i,j} \omega_{ij}^{\frac{\rho}{p'}} |D_{ij} u_\epsilon|^{(s-1)\rho - \xi} \omega_{ij}^{\frac{\rho}{p}} |D_{ij} u_\epsilon|^\xi \right)^\rho dx \\
& \leq c \epsilon^\rho \left( \sum_{i,j} \int_{\Omega} \omega_{ij}^{a\rho} |D_{ij} u_\epsilon|^{a((s-1)\rho - \xi)} dx \right)^{\frac{1}{a}} \cdot \left( \int_{\Omega} |Du_\epsilon|^{a'\xi} dx \right)^{\frac{1}{a'}} \\
& \leq c \epsilon^\rho \left( \sum_{i,j} \int_{\Omega} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\epsilon|^{s-2} \sum_{r,s} \omega_{rs}^{\frac{1}{p}} |D_{rs} u_\epsilon|^2 \right)^{\frac{1}{a}} \cdot \|Du_\epsilon\|_{L^\tau(\Omega, M^{m \times n})}^{\frac{\tau}{a}}.
\end{aligned}$$

And by the injection:  $L^{\frac{n(p_s-1)}{n-1}} \hookrightarrow L^\tau \quad \forall \tau > \frac{n(p_s-1)}{n-1}$  and thanks to (6.10)-(6.12), we get:

$$\begin{aligned}
\int_{\Omega} |\epsilon \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\epsilon|^{s-1} \omega_{ij}^{\frac{1}{p}}|^\rho dx & \leq c \epsilon^\rho \left( \frac{c \|\mu\|_M}{\epsilon} \right)^{\frac{s}{(s-1)a}} \cdot c^{\frac{\tau}{a}} \\
& \leq c \cdot c^{\frac{\tau}{a}} \epsilon^{\frac{a((s-1)\rho - s)}{a(s-1)}} \\
& \leq c \cdot c^{\frac{\tau}{a}} \epsilon^{\frac{a\xi}{a(s-1)}} \\
& \leq c \cdot c^{\frac{\tau}{a}} \cdot \epsilon^{\frac{\xi}{s-1}}
\end{aligned}$$

with  $\frac{\xi}{s-1} > 0$ . Hence

$$\lim_{\epsilon \rightarrow 0} \left\| \epsilon \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_\epsilon|^{s-1} \omega_{rs}^{\frac{1}{p}} D_{rs} u_\epsilon \right\|_{L^p(\Omega, M^{m \times n})} = 0, \quad \forall \rho < \frac{s}{s-1}.$$

In particular for  $\rho = 1$

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} |\varepsilon \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_{\varepsilon}|^{s-1} \omega_{rs}^{\frac{1}{p}} D_{rs} u_{\varepsilon}| dx = 0,$$

which mean that

$$\tau[\varepsilon](x, u_{\varepsilon}, Du_{\varepsilon}) = \varepsilon \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_{\varepsilon}|^{s-2} \omega_{rs}^{\frac{1}{p}} D_{rs} u_{\varepsilon} \rightharpoonup 0$$

in  $L^1(\Omega, \mathbb{M}^{m \times n})$ .

As well as by the proposition 6.1,  $div\sigma(x, u_{\varepsilon}, Du_{\varepsilon})$  converges to  $div\sigma(x, u, apDu)$ , in the sense of the distributions, and as

$$\tau[\varepsilon](x, u_{\varepsilon}, Du_{\varepsilon}) = \varepsilon \sum_{i,j} \omega_{ij}^{\frac{1}{p'}} |D_{ij} u_{\varepsilon}|^{s-2} \omega_{rs}^{\frac{1}{p}} D_{rs} u_{\varepsilon} \rightharpoonup 0,$$

in  $L^1(\Omega, \mathbb{M}^{m \times n})$ . Then  $div\sigma_{\varepsilon}(x, u_{\varepsilon}, Du_{\varepsilon})$  converge to  $div\sigma(x, u, apDu)$  in the sense of distributions, i-e:  $u$  is the solution of the system

$$\begin{cases} -div\sigma(x, u, apDu) & = \mu & \in \Omega \\ u & = 0, & \text{on } \partial\Omega. \end{cases}$$

In the same way as in the case of  $\theta = p - 1$ , we have

$$\int_{|u| \leq \alpha} |apDu|^s dx < c(\alpha) < \infty \text{ and } p < s.$$

So we conclude as in the proof of the proposition 6.1, in order to get the estimation of theorem (2.1). This completes the proof of the theorem.

## References

1. Hungerbuhler. N, *Quasilinear elliptic systems in divergence form with weak monotonicity*, New York J. Math. 5 (1999), 83–90.
2. Hungerbuhler. N, G. Dolzmann, S. Muller, *Nonlinear elliptic systems with measure-valued right hand side*, Math. Z. 226 (1997), 545–574.
3. Fabien Augsburguer, *Young measures and quasilinear systems in divergence form with weak monotonicity*, Thesis n° 1448, University Press, Fribourg, 2004.
4. P. Lions, F. Murat, *Solutions renormalisées d'équations elliptiques*, to appear.
5. P. Bénilan, L. Boccardo, T. Gallouët, R. Gariepy, M. Pierre, J. Vasquez, *An  $L^1$  theory on existence and uniqueness of solutions of nonlinear elliptic equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire 37 (1995), 16–26.
6. Rami. E, Azroul. E, Ellekhlifi. M, *Quasilinear degenerated elliptic system in divergence form with mild monotonicity in weighted Sobolev spaces*, Afrika Matematika (2019), 1–16.
7. Rami. E, Barbara. A, Azroul. E, *Existence of a weak solution of some quasilinear elliptic system in a weighted Sobolev space*, International Journal of Mathematics Trends and Technology (IJMTT), 66(2), Feb 2020.

*El Houcine Rami,*  
*Laboratory L2MASI, Department of Mathematics,*  
*Faculty of Sciences Dhar El Mahraz,*  
*Sidi Mohamed Ben Abdellah University,*  
*B.P 1796 Atlas, Fez, Morocco.*  
*E-mail address: ramielhoucine@gmail.com*

and

*Abdelkrim Barbara,*  
*Laboratory L2MASI, Department of Mathematics,*

*Faculty of Sciences Dhar El Mahraz,  
Sidi Mohamed Ben Abdellah University,  
B.P 1796 Atlas, Fez, Morocco.  
E-mail address: [abdelkrim.barbara@usmba.ac.ma](mailto:abdelkrim.barbara@usmba.ac.ma)*

*and*

*Elhoussine Azroul,  
Laboratory L2MASI, Department of Mathematics,  
Faculty of Sciences Dhar El Mahraz,  
Sidi Mohamed Ben Abdellah University,  
B.P 1796 Atlas, Fez, Morocco.  
E-mail address: [elhoussine.azroul@gmail.com](mailto:elhoussine.azroul@gmail.com)*